



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 24, 2026 – 09:14 PM EDT

PDB ID : 1AED / pdb_00001aed
Title : SPECIFICITY OF LIGAND BINDING TO A BURIED POLAR CAVITY AT THE ACTIVE SITE OF CYTOCHROME C PEROXIDASE (3,4-DIMETHYLTHIAZOLE)
Authors : Musah, R.A.; Jensen, G.M.; Fitzgerald, M.M.; Mcree, D.E.; Goodin, D.B.
Deposited on : 1997-02-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	294	 76% 19%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2999 atoms, of which 511 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	291	2841	1492	503	388	452	6	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	THR	variant	UNP P00431
A	152	GLY	ASP	variant	UNP P00431
A	191	GLY	TRP	engineered mutation	UNP P00431
A	272	ASP	ASN	conflict	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0

- Molecule 3 is 3,4-DIMETHYLTHIAZOLIUM ION (CCD ID: DTI) (formula: C₅H₈NS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	S		
3	A	1	15	5	8	1	1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	100	100	100	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYTOCHROME C PEROXIDASE

Chain A:  76% 19%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	108.00Å 77.30Å 51.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 2.10 7.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	79.0 (7.00-2.10) 68.9 (7.00-2.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.74 (at 2.01Å)	Xtrriage
Refinement program	XTALVIEW	Depositor
R, R_{free}	(Not available) , (Not available) 0.209 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtrriage
Anisotropy	0.240	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.55 , 104.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	2999	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, DTI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.97	8/2402 (0.3%)	1.88	69/3250 (2.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	47	VAL	CA-CB	6.71	1.63	1.54
1	A	181	HIS	CD2-NE2	-6.62	1.30	1.37
1	A	6	HIS	CD2-NE2	-6.49	1.30	1.37
1	A	175	HIS	CD2-NE2	-6.34	1.30	1.37
1	A	96	HIS	CD2-NE2	-6.33	1.30	1.37
1	A	60	HIS	CD2-NE2	-6.14	1.31	1.37
1	A	52	HIS	CD2-NE2	-5.25	1.32	1.37
1	A	175	HIS	CG-ND1	-5.02	1.32	1.38

All (69) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	255	GLN	OE1-CD-NE2	-12.38	110.22	122.60
1	A	159	GLN	OE1-CD-NE2	-10.43	112.17	122.60
1	A	120	GLN	N-CA-C	9.12	123.94	111.54
1	A	253	ASN	OD1-CG-ND2	-9.01	113.59	122.60
1	A	255	GLN	CG-CD-NE2	8.59	129.28	116.40
1	A	20	GLN	OE1-CD-NE2	-8.43	114.17	122.60
1	A	195	ASN	OD1-CG-ND2	-8.33	114.27	122.60
1	A	240	GLN	OE1-CD-NE2	-8.29	114.31	122.60
1	A	127	ARG	CB-CG-CD	-8.20	92.44	111.30
1	A	78	ASN	OD1-CG-ND2	-8.18	114.42	122.60
1	A	205	ASN	OD1-CG-ND2	-8.13	114.47	122.60
1	A	162	ASN	N-CA-C	8.11	128.06	110.80
1	A	136	ASP	CA-CB-CG	8.03	120.63	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	62	ASN	OD1-CG-ND2	-7.92	114.69	122.60
1	A	159	GLN	CG-CD-NE2	7.60	127.81	116.40
1	A	117	GLN	OE1-CD-NE2	-7.41	115.19	122.60
1	A	162	ASN	CA-C-O	7.10	130.66	120.51
1	A	220	ASN	OD1-CG-ND2	-7.10	115.50	122.60
1	A	24	ASN	OD1-CG-ND2	-7.01	115.59	122.60
1	A	82	ASN	OD1-CG-ND2	-6.94	115.66	122.60
1	A	208	ASN	OD1-CG-ND2	-6.76	115.84	122.60
1	A	222	GLN	OE1-CD-NE2	-6.66	115.94	122.60
1	A	128	CYS	N-CA-C	6.63	120.21	110.59
1	A	42	TYR	N-CA-C	6.59	121.15	113.18
1	A	195	ASN	N-CA-C	6.56	120.20	112.72
1	A	34	ASP	N-CA-C	6.49	120.94	112.89
1	A	216	ASN	OD1-CG-ND2	-6.48	116.12	122.60
1	A	164	ASN	OD1-CG-ND2	-6.44	116.16	122.60
1	A	254	ASP	CA-CB-CG	6.42	119.02	112.60
1	A	90	LYS	CA-CB-CG	-6.20	101.70	114.10
1	A	51	TRP	CB-CG-CD1	-6.19	117.61	126.90
1	A	12	LYS	N-CA-C	6.12	119.48	110.30
1	A	51	TRP	CG-CD2-CE3	6.08	139.99	133.90
1	A	210	ASP	CA-CB-CG	6.03	118.63	112.60
1	A	181	HIS	CB-CG-CD2	-5.95	123.46	131.20
1	A	195	ASN	CB-CG-ND2	5.85	125.18	116.40
1	A	130	ARG	CG-CD-NE	-5.85	99.13	112.00
1	A	38	ASN	CA-CB-CG	-5.83	106.77	112.60
1	A	162	ASN	CA-CB-CG	5.80	118.40	112.60
1	A	87	ASN	OD1-CG-ND2	-5.80	116.80	122.60
1	A	175	HIS	CB-CG-ND1	5.79	131.39	122.70
1	A	24	ASN	CB-CG-ND2	5.64	124.87	116.40
1	A	67	TYR	N-CA-C	5.61	117.84	111.11
1	A	117	GLN	CG-CD-NE2	5.61	124.81	116.40
1	A	133	THR	CA-C-N	5.60	125.58	120.03
1	A	133	THR	C-N-CA	5.60	125.58	120.03
1	A	175	HIS	CB-CG-CD2	-5.59	123.93	131.20
1	A	60	HIS	CB-CG-CD2	-5.59	123.93	131.20
1	A	276	PHE	CA-C-N	5.55	125.56	119.89
1	A	276	PHE	C-N-CA	5.55	125.56	119.89
1	A	109	SER	N-CA-C	5.54	117.40	111.36
1	A	119	MET	CA-C-N	5.53	129.98	122.07
1	A	119	MET	C-N-CA	5.53	129.98	122.07
1	A	240	GLN	CG-CD-NE2	5.49	124.64	116.40
1	A	151	ALA	N-CA-C	5.48	117.96	111.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	78	ASN	CB-CG-ND2	5.43	124.55	116.40
1	A	70	THR	N-CA-C	5.41	119.50	113.01
1	A	184	ASN	OD1-CG-ND2	-5.41	117.19	122.60
1	A	120	GLN	CB-CA-C	-5.40	103.92	111.80
1	A	166	ARG	N-CA-CB	-5.39	102.20	110.12
1	A	272	ASP	CA-CB-CG	5.38	117.97	112.60
1	A	217	ASP	N-CA-C	5.27	118.81	112.38
1	A	126	TRP	CG-CD2-CE3	5.19	139.09	133.90
1	A	166	ARG	CG-CD-NE	-5.08	100.82	112.00
1	A	253	ASN	N-CA-C	5.06	119.43	113.16
1	A	253	ASN	CA-CB-CG	-5.05	107.55	112.60
1	A	200	ASN	OD1-CG-ND2	-5.04	117.56	122.60
1	A	66	SER	N-CA-C	5.02	118.67	112.54
1	A	166	ARG	NE-CZ-NH2	-5.02	114.68	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2338	503	2209	16	0
2	A	43	0	30	0	0
3	A	7	8	8	0	0
4	A	100	0	0	5	0
All	All	2488	511	2247	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:CYS:HA	4:A:326:HOH:O	1.62	0.98
1:A:130:ARG:NE	4:A:442:HOH:O	2.37	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:20:GLN:HE22	1:A:287:LYS:H	1.51	0.57
1:A:195:ASN:HD22	1:A:195:ASN:H	1.54	0.54
1:A:130:ARG:CZ	4:A:442:HOH:O	2.56	0.53
1:A:67:TYR:HA	1:A:130:ARG:HG2	1.90	0.53
1:A:119:MET:O	1:A:120:GLN:HG2	2.11	0.49
1:A:130:ARG:NH2	4:A:442:HOH:O	2.45	0.49
1:A:93:GLU:HB3	1:A:94:PRO:HD2	1.96	0.46
1:A:195:ASN:H	1:A:195:ASN:ND2	2.14	0.44
1:A:216:ASN:HD21	1:A:220:ASN:HB2	1.83	0.43
1:A:119:MET:C	1:A:120:GLN:HG2	2.43	0.43
1:A:23:TYR:HD1	1:A:24:ASN:HD22	1.66	0.43
1:A:127:ARG:HH11	1:A:127:ARG:HD3	1.77	0.41
1:A:166:ARG:HH22	1:A:250:GLU:CD	2.27	0.41
1:A:106:ASP:CG	4:A:326:HOH:O	2.64	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/294 (98%)	283 (98%)	5 (2%)	1 (0%)	36 36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	162	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	246/251 (98%)	233 (95%)	13 (5%)	20	19

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	LEU
1	A	35	GLU
1	A	47	VAL
1	A	53	ILE
1	A	72	ARG
1	A	74	LYS
1	A	120	GLN
1	A	171	LEU
1	A	195	ASN
1	A	206	LEU
1	A	212	LYS
1	A	219	ASN
1	A	278	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	78	ASN
1	A	195	ASN
1	A	208	ASN
1	A	240	GLN
1	A	292	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	A	295	4,1	50,50,50	2.40	28 (56%)	67,82,82	1.19	6 (8%)
3	DTI	A	296	-	6,7,7	0.71	0	7,9,9	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	295	4,1	-	2/14/54/54	-
3	DTI	A	296	-	-	-	0/1/1/1

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	295	HEM	C4D-ND	-4.30	1.32	1.40
2	A	295	HEM	FE-NB	4.08	2.07	1.94
2	A	295	HEM	FE-ND	4.03	2.07	1.94
2	A	295	HEM	C1B-NB	-3.92	1.33	1.40
2	A	295	HEM	C4A-NA	-3.90	1.32	1.39
2	A	295	HEM	FE-NA	3.81	2.07	1.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	295	HEM	C3C-C4C	-3.80	1.39	1.46
2	A	295	HEM	FE-NC	3.52	2.06	1.95
2	A	295	HEM	CBC-CAC	3.39	1.46	1.30
2	A	295	HEM	CBB-CAB	3.24	1.45	1.30
2	A	295	HEM	C1B-C2B	-3.15	1.38	1.44
2	A	295	HEM	C4C-NC	-3.03	1.33	1.39
2	A	295	HEM	C1A-NA	-2.94	1.34	1.39
2	A	295	HEM	C1C-C2C	-2.94	1.39	1.45
2	A	295	HEM	C1A-C2A	-2.90	1.38	1.44
2	A	295	HEM	C4D-C3D	-2.90	1.40	1.45
2	A	295	HEM	O2A-CGA	-2.76	1.21	1.30
2	A	295	HEM	C3C-C2C	2.71	1.42	1.37
2	A	295	HEM	CAB-C3B	-2.63	1.40	1.47
2	A	295	HEM	C1D-C2D	-2.61	1.39	1.44
2	A	295	HEM	C3B-C4B	-2.59	1.39	1.44
2	A	295	HEM	O2D-CGD	-2.56	1.22	1.30
2	A	295	HEM	C4B-NB	-2.41	1.34	1.38
2	A	295	HEM	C1C-NC	-2.30	1.35	1.39
2	A	295	HEM	CAC-C3C	-2.23	1.41	1.47
2	A	295	HEM	C4A-C3A	-2.19	1.38	1.43
2	A	295	HEM	C3B-C2B	2.19	1.41	1.37
2	A	295	HEM	C1D-ND	-2.15	1.34	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	295	HEM	C3B-C2B-C1B	-2.68	104.40	106.41
2	A	295	HEM	C3C-C2C-C1C	-2.64	104.55	107.05
2	A	295	HEM	O2D-CGD-CBD	2.33	121.35	114.00
2	A	295	HEM	C2B-C1B-NB	2.12	112.28	109.84
2	A	295	HEM	O2A-CGA-CBA	2.12	120.71	114.00
2	A	295	HEM	O2A-CGA-O1A	-2.07	118.02	123.33

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	295	HEM	CAA-CBA-CGA-O2A
2	A	295	HEM	CAA-CBA-CGA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	291/294 (98%)	0.05	1 (0%) 90 91	4, 13, 28, 43	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	226	LYS	3.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DTI	A	296	7/7	0.91	0.12	15,15,15,15	0
2	HEM	A	295	43/43	0.95	0.07	3,9,12,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.