



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:41 PM UTC

PDB ID : 1AEI / pdb_00001aei
Title : CRYSTAL STRUCTURE OF THE ANNEXIN XII HEXAMER
Authors : Luecke, H.; Chang, B.T.; Mailliard, W.S.; Schlaepfer, D.D.; Haigler, H.T.
Deposited on : 1995-09-23
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	
1	C	315	
1	D	315	
1	E	315	
1	F	315	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 18252 atoms, of which 3456 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ANNEXIN XII.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
1	A	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				
1	B	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				
1	C	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				
1	D	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				
1	E	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				
1	F	315	Total	C	H	N	O	S		0	0	0
			3038	1543	576	428	484	7				

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Ca	0	0
			4	4		
2	B	4	Total	Ca	0	0
			4	4		
2	C	4	Total	Ca	0	0
			4	4		
2	D	4	Total	Ca	0	0
			4	4		
2	E	4	Total	Ca	0	0
			4	4		
2	F	4	Total	Ca	0	0
			4	4		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: ANNEXIN XII




• Molecule 1: ANNEXIN XII




• Molecule 1: ANNEXIN XII






• Molecule 1: ANNEXIN XII

Chain D: 64% 29% 6% •




• Molecule 1: ANNEXIN XII

Chain E: 66% 28% 5% •



• Molecule 1: ANNEXIN XII

Chain F: 64% 29% 5% •





4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.20Å 179.20Å 100.70Å 90.00° 97.60° 90.00°	Depositor
Resolution (Å)	10.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.80)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.236 , 0.278	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	18252	wwPDB-VP
Average B, all atoms (Å ²)	9.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.38	10/2498 (0.4%)	1.97	76/3361 (2.3%)
1	B	1.37	18/2498 (0.7%)	2.01	69/3361 (2.1%)
1	C	1.44	19/2498 (0.8%)	2.09	100/3361 (3.0%)
1	D	1.36	16/2498 (0.6%)	2.02	79/3361 (2.4%)
1	E	1.36	14/2498 (0.6%)	2.03	88/3361 (2.6%)
1	F	1.41	15/2498 (0.6%)	2.05	74/3361 (2.2%)
All	All	1.39	92/14988 (0.6%)	2.03	486/20166 (2.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 92 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	254	HIS	CD2-NE2	-8.08	1.28	1.37
1	D	62	HIS	CD2-NE2	-7.66	1.29	1.37
1	E	95	HIS	CD2-NE2	-7.48	1.29	1.37
1	D	169	HIS	CD2-NE2	-7.37	1.29	1.37
1	A	164	PRO	CA-CB	-7.37	1.45	1.54

The worst 5 of 486 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	203	GLN	OE1-CD-NE2	-14.70	107.90	122.60
1	B	203	GLN	OE1-CD-NE2	-12.63	109.97	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	203	GLN	OE1-CD-NE2	-10.77	111.83	122.60
1	F	304	GLY	O-C-N	10.68	134.03	122.60
1	C	106	ASN	OD1-CG-ND2	-10.39	112.21	122.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	258	LYS	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2462	576	2450	36	0
1	B	2462	576	2450	28	0
1	C	2462	576	2450	37	0
1	D	2462	576	2450	32	0
1	E	2462	576	2450	26	0
1	F	2462	576	2450	39	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	4	0	0	0	0
2	E	4	0	0	0	0
2	F	4	0	0	0	0
All	All	14796	3456	14700	194	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 194 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:293:SER:HB3	1:F:296:GLU:HG3	1.67	0.76

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:160:LYS:HD3	1:D:163:GLU:HB2	1.69	0.73
1:B:265:LYS:HD2	1:B:265:LYS:H	1.54	0.72
1:C:160:LYS:HD3	1:C:163:GLU:HB2	1.72	0.71
1:D:165:VAL:HA	1:D:203:GLN:HE21	1.55	0.70

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/315 (99%)	302 (96%)	9 (3%)	2 (1%)	21	51
1	B	313/315 (99%)	304 (97%)	7 (2%)	2 (1%)	21	51
1	C	313/315 (99%)	302 (96%)	9 (3%)	2 (1%)	21	51
1	D	313/315 (99%)	305 (97%)	7 (2%)	1 (0%)	36	66
1	E	313/315 (99%)	301 (96%)	11 (4%)	1 (0%)	36	66
1	F	313/315 (99%)	302 (96%)	9 (3%)	2 (1%)	21	51
All	All	1878/1890 (99%)	1816 (97%)	52 (3%)	10 (0%)	24	55

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLY
1	B	186	GLY
1	B	132	LYS
1	C	186	GLY
1	D	132	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/260 (100%)	238 (92%)	22 (8%)	10	31
1	B	260/260 (100%)	232 (89%)	28 (11%)	6	21
1	C	260/260 (100%)	235 (90%)	25 (10%)	8	26
1	D	260/260 (100%)	232 (89%)	28 (11%)	6	21
1	E	260/260 (100%)	235 (90%)	25 (10%)	8	26
1	F	260/260 (100%)	234 (90%)	26 (10%)	7	24
All	All	1560/1560 (100%)	1406 (90%)	154 (10%)	7	24

5 of 154 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	150	LEU
1	F	137	LYS
1	E	220	LEU
1	F	24	LYS
1	F	258	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	14	ASN
1	E	225	ASN
1	E	45	ASN
1	E	106	ASN
1	E	316	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 24 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.