



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:35 PM UTC

PDB ID : 3AE7 / pdb_00003ae7
Title : Crystal structure of porcine heart mitochondrial complex II bound with 2-Iodo-N-(3-isopropoxy-phenyl)-benzamide
Authors : Harada, S.; Sasaki, T.; Shindo, M.; Kido, Y.; Inaoka, D.K.; Omori, J.; Osanai, A.; Sakamoto, K.; Mao, J.; Matsuoka, S.; Inoue, M.; Honma, T.; Tanaka, A.; Kita, K.
Deposited on : 2010-02-04
Resolution : 3.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1062 (3.72-3.52)
Clashscore	190562	1092 (3.72-3.52)
Ramachandran outliers	187476	1057 (3.72-3.52)
Sidechain outliers	187428	1055 (3.72-3.52)
RSRZ outliers	180081	1060 (3.72-3.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	622	81% 16% ..
2	B	252	80% 13% 5%
3	C	140	90% 8% ..
4	D	103	95% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	F3S	B	304	-	-	X	-

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 8615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	613	Total	C	N	O	S	0	0	0
			4729	2954	848	895	32			

- Molecule 2 is a protein called Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	239	Total	C	N	O	S	0	0	0
			1922	1214	326	360	22			


- Molecule 3 is a protein called Succinate dehydrogenase cytochrome b560 subunit, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	138	Total	C	N	O	S	0	0	0
			1064	695	179	183	7			

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	102	Total	C	N	O	S	0	0	0
			765	499	128	133	5			

- Molecule 5 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 6 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe₂S₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	Fe	S	0	0
			4	2	2		

- Molecule 7 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 8 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	Fe	S	0	0
			7	3	4		

- Molecule 9 is 2-iodo-N-[3-(1-methylethoxy)phenyl]benzamide (CCD ID: 12J) (formula: $\text{C}_{16}\text{H}_{16}\text{INO}_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	B	1	Total	C	I	N	O	0	0
			20	16	1	1	2		

- Molecule 10 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial



- Molecule 2: Succinate dehydrogenase [ubiquinone] iron-sulfur subunit, mitochondrial



- Molecule 3: Succinate dehydrogenase cytochrome b560 subunit, mitochondrial



- Molecule 4: Succinate dehydrogenase [ubiquinone] cytochrome b small subunit, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.17Å 83.99Å 293.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.13 – 3.62 48.13 – 3.62	Depositor EDS
% Data completeness (in resolution range)	96.0 (48.13-3.62) 96.1 (48.13-3.62)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 3.57Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.256 , 0.305 0.260 , 0.307	Depositor DCC
R_{free} test set	1046 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å ²)	122.3	Xtriage
Anisotropy	0.180	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 71.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	8615	wwPDB-VP
Average B, all atoms (Å ²)	145.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: F3S, FAD, HEM, 12J, SF4, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/4828	0.79	5/6531 (0.1%)
2	B	0.41	0/1964	0.74	2/2648 (0.1%)
3	C	0.46	1/1091 (0.1%)	0.80	0/1483
4	D	0.38	0/784	0.77	0/1066
All	All	0.42	1/8667 (0.0%)	0.78	7/11728 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	119	ILE	CA-CB	6.14	1.57	1.54

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	434	ARG	N-CA-C	10.01	125.40	110.39
1	A	58	THR	N-CA-C	-6.79	104.10	112.38
1	A	432	SER	CB-CA-C	6.46	121.00	109.38
1	A	434	ARG	N-CA-CB	-6.43	100.91	109.78
1	A	57	HIS	N-CA-C	6.02	118.80	111.82
2	B	180	GLY	CA-C-N	5.83	125.30	119.24
2	B	180	GLY	C-N-CA	5.83	125.30	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4729	0	4618	62	0
2	B	1922	0	1904	21	0
3	C	1064	0	1104	5	0
4	D	765	0	773	2	0
5	A	53	0	31	12	0
6	B	4	0	0	0	0
7	B	8	0	0	0	0
8	B	7	0	0	2	0
9	B	20	0	16	2	0
10	C	43	0	30	2	0
All	All	8615	0	8476	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:57:HIS:NE2	5:A:700:FAD:HM82	1.17	1.45
2:B:219:MET:HE1	2:B:232:GLY:HA3	1.60	0.81
1:A:113:ASN:HD22	2:B:138:GLY:H	1.27	0.80
1:A:150:VAL:H	1:A:154:THR:HG22	1.52	0.74
1:A:61:ALA:HB3	1:A:155:GLY:HA3	1.70	0.74
1:A:415:LEU:HG	5:A:700:FAD:C2	2.19	0.73
1:A:59:VAL:HG23	1:A:159:LEU:HD23	1.71	0.72
1:A:61:ALA:HA	5:A:700:FAD:N5	2.07	0.68
2:B:168:CYS:HB3	8:B:304:F3S:S3	2.33	0.68
9:B:1201:12J:O1	9:B:1201:12J:I	2.82	0.67
2:B:51:ILE:HD11	2:B:59:LEU:HD22	1.75	0.67
1:A:57:HIS:CD2	5:A:700:FAD:HM82	2.22	0.67
2:B:177:LYS:HB3	2:B:210:PHE:HB3	1.76	0.67
1:A:58:THR:HG22	5:A:700:FAD:O4'	1.99	0.61
2:B:168:CYS:CB	8:B:304:F3S:S3	2.90	0.60
1:A:342:SER:HA	1:A:353:VAL:HG11	1.85	0.58
1:A:556:LYS:H	1:A:556:LYS:HD2	1.69	0.58
1:A:57:HIS:NE2	5:A:700:FAD:C8	2.67	0.57
1:A:415:LEU:HG	5:A:700:FAD:O2	2.05	0.57
1:A:49:THR:OG1	5:A:700:FAD:O2B	2.22	0.57
1:A:322:VAL:HG12	1:A:323:TYR:H	1.69	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:ALA:N	5:A:700:FAD:N1A	2.40	0.56
1:A:104:ALA:HA	1:A:416:LEU:HD11	1.88	0.55
1:A:246:GLN:HB2	1:A:372:PRO:HG3	1.87	0.55
1:A:111:LEU:HD11	1:A:419:VAL:HG21	1.88	0.54
1:A:57:HIS:CE1	5:A:700:FAD:HM82	2.22	0.53
1:A:354:THR:HG23	1:A:355:LYS:HG2	1.91	0.53
1:A:67:ALA:HB2	1:A:108:VAL:HG21	1.92	0.52
3:C:29:HIS:HD2	3:C:30:ILE:HG23	1.73	0.52
10:C:1305:HEM:CHA	10:C:1305:HEM:HBD1	2.39	0.51
2:B:164:CYS:HB2	2:B:221:CYS:HB2	1.92	0.51
2:B:102:LEU:HD22	2:B:166:THR:HG21	1.93	0.51
1:A:563:ASP:H	1:A:571:GLN:HE22	1.58	0.51
3:C:96:PHE:HB3	3:C:97:PRO:HD3	1.93	0.50
1:A:100:MET:HA	1:A:420:VAL:HG11	1.94	0.50
1:A:51:LEU:HD21	1:A:229:THR:HG21	1.93	0.50
10:C:1305:HEM:HBD1	10:C:1305:HEM:HHA	1.94	0.49
1:A:115:GLY:HA2	2:B:129:LEU:HD12	1.95	0.49
1:A:258:ILE:HD13	1:A:263:CYS:HB2	1.95	0.49
1:A:368:MET:HE3	1:A:408:ASN:HA	1.95	0.49
1:A:10:SER:C	1:A:12:GLN:H	2.21	0.49
2:B:116:SER:HB2	3:C:15:PHE:HD1	1.78	0.48
1:A:118:PHE:HA	1:A:150:VAL:HG22	1.95	0.48
1:A:516:TRP:HB3	2:B:60:THR:HG21	1.94	0.48
4:D:72:LEU:O	4:D:76:LEU:HB2	2.14	0.48
1:A:323:TYR:HD2	1:A:357:PRO:HB2	1.79	0.47
1:A:116:MET:HA	1:A:161:THR:HG21	1.95	0.47
1:A:463:THR:HG23	1:A:464:ILE:HG13	1.97	0.47
1:A:429:ILE:O	1:A:432:SER:O	2.33	0.47
1:A:104:ALA:HB3	1:A:105:PRO:HD3	1.96	0.47
1:A:289:ALA:HB1	1:A:292:ALA:HB3	1.97	0.47
2:B:43:MET:HA	2:B:88:ARG:HA	1.97	0.46
1:A:57:HIS:C	1:A:59:VAL:N	2.70	0.46
2:B:155:LEU:HD23	2:B:185:MET:HE2	1.97	0.46
2:B:45:LEU:HA	2:B:48:LEU:HD12	1.99	0.45
2:B:216:HIS:CD2	9:B:1201:12J:H6	2.51	0.45
1:A:326:LEU:HA	1:A:329:LEU:HD23	1.99	0.45
1:A:518:THR:HA	1:A:521:VAL:HG22	1.97	0.45
1:A:195:LEU:HD22	1:A:454:LEU:HD21	1.99	0.44
2:B:108:ILE:HD12	2:B:112:VAL:HG12	1.99	0.44
1:A:61:ALA:HB1	5:A:700:FAD:C4	2.47	0.44
1:A:59:VAL:HG22	1:A:156:HIS:HA	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:SER:HB3	1:A:337:ARG:HH12	1.83	0.44
1:A:306:THR:HA	1:A:309:ILE:HD12	1.99	0.44
1:A:514:MET:HA	1:A:514:MET:HE2	1.99	0.44
3:C:71:GLU:HG2	4:D:132:MET:HE1	2.00	0.44
1:A:381:ARG:HG3	1:A:388:GLN:HB3	2.00	0.43
2:B:219:MET:HE2	2:B:219:MET:HA	1.98	0.43
1:A:270:ARG:HD3	1:A:276:LEU:HD13	1.99	0.43
1:A:299:ASP:O	1:A:303:ARG:HB2	2.17	0.43
1:A:443:LYS:HB2	1:A:446:ALA:HB2	2.00	0.43
1:A:49:THR:HA	5:A:700:FAD:N3A	2.33	0.43
1:A:13:TYR:HA	1:A:14:PRO:HD3	1.92	0.43
1:A:65:ILE:HD12	1:A:108:VAL:HG22	2.01	0.43
1:A:88:SER:HB2	1:A:406:GLY:HA3	2.00	0.42
2:B:60:THR:O	2:B:110:ASP:O	2.38	0.42
1:A:258:ILE:HG13	1:A:265:ILE:HD11	2.02	0.42
1:A:390:VAL:HA	1:A:391:PRO:HD3	1.91	0.41
1:A:72:MET:HE1	1:A:121:THR:HG21	2.02	0.41
1:A:72:MET:HE3	1:A:128:GLN:H	1.86	0.41
1:A:18:HIS:HB2	1:A:205:ILE:HA	2.02	0.41
2:B:233:LYS:HA	3:C:117:LEU:HD13	2.01	0.41
2:B:235:ILE:HA	2:B:238:ILE:HD12	2.03	0.41
1:A:191:GLY:HA3	1:A:206:ARG:HA	2.02	0.41
1:A:414:SER:O	1:A:418:LEU:HD13	2.21	0.41
2:B:225:CYS:HA	2:B:226:PRO:HD3	1.87	0.40
1:A:129:ARG:O	1:A:147:CYS:HB3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	611/622 (98%)	567 (93%)	39 (6%)	5 (1%)	16 46

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	237/252 (94%)	223 (94%)	13 (6%)	1 (0%)	30	59
3	C	136/140 (97%)	132 (97%)	3 (2%)	1 (1%)	18	49
4	D	100/103 (97%)	97 (97%)	3 (3%)	0	100	100
All	All	1084/1117 (97%)	1019 (94%)	58 (5%)	7 (1%)	21	51

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	GLY
1	A	290	PRO
2	B	110	ASP
1	A	433	CYS
1	A	569	GLN
3	C	82	LEU
1	A	260	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	499/506 (99%)	483 (97%)	16 (3%)	34	54
2	B	214/220 (97%)	205 (96%)	9 (4%)	26	49
3	C	117/118 (99%)	113 (97%)	4 (3%)	32	53
4	D	76/76 (100%)	75 (99%)	1 (1%)	61	69
All	All	906/920 (98%)	876 (97%)	30 (3%)	33	54

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	49	THR
1	A	59	VAL
1	A	65	ILE
1	A	109	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	165	ARG
1	A	167	LEU
1	A	276	LEU
1	A	277	ILE
1	A	294	ASP
1	A	322	VAL
1	A	507	LEU
1	A	534	LEU
1	A	556	LYS
1	A	561	GLU
1	A	568	ILE
1	A	597	LEU
2	B	51	ILE
2	B	69	ILE
2	B	177	LYS
2	B	189	ARG
2	B	192	ILE
2	B	214	ARG
2	B	217	THR
2	B	220	ASN
2	B	235	ILE
3	C	30	ILE
3	C	64	LEU
3	C	80	LEU
3	C	122	LEU
4	D	78	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	44	ASN
1	A	62	GLN
1	A	113	ASN
1	A	128	GLN
1	A	143	GLN
1	A	156	HIS
1	A	278	ASN
1	A	280	GLN
1	A	384	ASN
1	A	408	ASN
1	A	461	ASN
1	A	491	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	571	GLN
2	B	121	GLN
2	B	144	GLN
2	B	174	ASN
2	B	220	ASN
2	B	230	ASN
3	C	17	ASN
3	C	29	HIS
3	C	104	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
9	12J	B	1201	-	21,21,21	1.00	2 (9%)	28,28,28	0.76	1 (3%)
6	FES	B	302	2	0,4,4	-	-	-		
10	HEM	C	1305	3,4	50,50,50	1.94	11 (22%)	67,82,82	1.37	4 (5%)
8	F3S	B	304	2	0,9,9	-	-	-		

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FAD	A	700	1	58,58,58	1.12	5 (8%)	85,89,89	1.60	16 (18%)
7	SF4	B	303	2	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	12J	B	1201	-	-	4/12/12/12	0/2/2/2
6	FES	B	302	2	-	-	0/1/1/1
10	HEM	C	1305	3,4	-	11/14/54/54	-
8	F3S	B	304	2	-	-	0/3/3/3
5	FAD	A	700	1	-	3/34/50/50	0/6/6/6
7	SF4	B	303	2	-	-	0/6/5/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	C	1305	HEM	C3D-C2D	8.12	1.54	1.36
10	C	1305	HEM	FE-ND	4.45	2.08	1.94
10	C	1305	HEM	FE-NB	4.42	2.08	1.94
10	C	1305	HEM	FE-NA	3.87	2.07	1.95
5	A	700	FAD	C4X-N5	3.75	1.38	1.30
5	A	700	FAD	C10-N1	2.91	1.39	1.33
10	C	1305	HEM	CAC-C3C	2.90	1.55	1.47
9	B	1201	12J	C8-N	-2.90	1.35	1.41
10	C	1305	HEM	FE-NC	2.82	2.04	1.95
10	C	1305	HEM	CAB-C3B	2.77	1.54	1.47
9	B	1201	12J	C2-C1	2.69	1.49	1.39
5	A	700	FAD	C2A-N1A	2.64	1.38	1.33
5	A	700	FAD	C2A-N3A	2.52	1.38	1.33
10	C	1305	HEM	CMA-C3A	2.16	1.55	1.50
10	C	1305	HEM	CMB-C2B	2.15	1.55	1.50
10	C	1305	HEM	CMC-C2C	2.12	1.55	1.50
5	A	700	FAD	C8A-N7A	2.08	1.35	1.31
10	C	1305	HEM	CMD-C2D	2.05	1.55	1.50

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	700	FAD	N3A-C2A-N1A	-5.98	119.53	128.58
10	C	1305	HEM	C4D-ND-C1D	5.96	112.27	105.21
5	A	700	FAD	C5A-C4A-N3A	-4.23	120.89	126.72
5	A	700	FAD	N9A-C8A-N7A	-3.86	108.47	113.94
5	A	700	FAD	C2A-N3A-C4A	3.44	120.23	111.83
5	A	700	FAD	C5A-N7A-C8A	3.36	108.73	103.45
5	A	700	FAD	C4-N3-C2	-3.22	119.92	125.64
10	C	1305	HEM	CAD-C3D-C4D	3.14	130.18	124.70
5	A	700	FAD	N3A-C4A-N9A	2.83	131.97	127.17
5	A	700	FAD	C4X-C4-N3	2.80	120.38	113.25
10	C	1305	HEM	C1B-NB-C4B	2.50	108.16	105.21
5	A	700	FAD	C10-C4X-N5	-2.45	119.81	124.81
5	A	700	FAD	O4-C4-C4X	-2.41	120.17	126.53
5	A	700	FAD	C9A-C5X-N5	-2.38	119.93	122.45
5	A	700	FAD	C4X-C10-N10	2.37	119.87	116.48
10	C	1305	HEM	C3B-C4B-NB	-2.34	107.79	109.47
5	A	700	FAD	C4A-C5A-N7A	-2.32	107.92	110.58
9	B	1201	12J	C10-O2-C14	2.27	122.42	119.43
5	A	700	FAD	C4X-C10-N1	-2.21	119.17	124.59
5	A	700	FAD	C4'-C3'-C2'	-2.18	109.94	113.57
5	A	700	FAD	C5X-C9A-N10	2.08	119.85	117.97

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	700	FAD	N10-C1'-C2'-O2'
10	C	1305	HEM	C2B-C3B-CAB-CBB
10	C	1305	HEM	C4B-C3B-CAB-CBB
10	C	1305	HEM	C2C-C3C-CAC-CBC
10	C	1305	HEM	C4C-C3C-CAC-CBC
10	C	1305	HEM	C2D-C3D-CAD-CBD
10	C	1305	HEM	C4D-C3D-CAD-CBD
10	C	1305	HEM	C3D-CAD-CBD-CGD
9	B	1201	12J	C9-C8-N-C7
9	B	1201	12J	C13-C8-N-C7
9	B	1201	12J	C9-C10-O2-C14
5	A	700	FAD	O4B-C4B-C5B-O5B
9	B	1201	12J	C11-C10-O2-C14
10	C	1305	HEM	CAA-CBA-CGA-O2A
10	C	1305	HEM	CAD-CBD-CGD-O1D
10	C	1305	HEM	CAA-CBA-CGA-O1A
10	C	1305	HEM	CAD-CBD-CGD-O2D

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	700	FAD	C3B-C4B-C5B-O5B

There are no ring outliers.


4 monomers are involved in 18 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	1201	12J	2	0
10	C	1305	HEM	2	0
8	B	304	F3S	2	0
5	A	700	FAD	12	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand HEM C 1305





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	613/622 (98%)	0.07	8 (1%) 75 49	110, 139, 186, 210	0
2	B	239/252 (94%)	0.09	4 (1%) 69 43	114, 134, 171, 195	0
3	C	138/140 (98%)	0.04	2 (1%) 73 47	125, 153, 187, 198	0
4	D	102/103 (99%)	0.01	2 (1%) 65 40	118, 149, 174, 188	0
All	All	1092/1117 (97%)	0.06	16 (1%) 72 46	110, 141, 184, 210	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	GLY	4.7
4	D	133	LEU	3.3
1	A	419	VAL	2.6
1	A	160	HIS	2.5
3	C	77	VAL	2.5
2	B	173	TRP	2.4
2	B	122	TYR	2.4
1	A	264	LEU	2.3
4	D	132	MET	2.3
1	A	132	GLY	2.3
2	B	234	ALA	2.3
2	B	160	LEU	2.2
1	A	296	ALA	2.2
3	C	38	PRO	2.1
1	A	205	ILE	2.1
1	A	410	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	FAD	A	700	53/53	0.82	0.16	127,127,127,127	0
8	F3S	B	304	7/7	0.93	0.10	142,142,142,142	0
9	12J	B	1201	20/20	0.97	0.14	136,137,138,138	0
10	HEM	C	1305	43/43	0.97	0.11	134,134,135,135	0
7	SF4	B	303	8/8	0.99	0.03	118,118,118,118	0
6	FES	B	302	4/4	0.99	0.05	105,106,106,106	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 12J B 1201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around HEM C 1305:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.