



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:36 PM UTC

PDB ID : 5AEG / pdb_00005aeg
Title : A bacterial protein structure in glycoside hydrolase family 31.
Authors : Jin, Y.; Speciale, G.; Davies, G.J.; Williams, S.J.; Goddard-Borger, E.D.
Deposited on : 2015-08-30
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	2% 90% 7% .
1	B	686	3% 87% 9% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALPHA-GLUCOSIDASE YIHQ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	667	Total	C	N	O	S	0	1	0
			5371	3445	894	1004	28			
1	B	670	Total	C	N	O	S	0	1	0
			5398	3458	902	1010	28			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	679	LEU	-	expression tag	UNP P32138
A	680	GLU	-	expression tag	UNP P32138
A	681	HIS	-	expression tag	UNP P32138
A	682	HIS	-	expression tag	UNP P32138
A	683	HIS	-	expression tag	UNP P32138
A	684	HIS	-	expression tag	UNP P32138
A	685	HIS	-	expression tag	UNP P32138
A	686	HIS	-	expression tag	UNP P32138
B	679	LEU	-	expression tag	UNP P32138
B	680	GLU	-	expression tag	UNP P32138
B	681	HIS	-	expression tag	UNP P32138
B	682	HIS	-	expression tag	UNP P32138
B	683	HIS	-	expression tag	UNP P32138
B	684	HIS	-	expression tag	UNP P32138
B	685	HIS	-	expression tag	UNP P32138
B	686	HIS	-	expression tag	UNP P32138


- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		
2	B	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			8	6	2		
4	A	1	Total	C	O	0	0
			8	6	2		
4	B	1	Total	C	O	0	0
			8	6	2		

- Molecule 5 is 5-fluoro-alpha-L-idopyranose (CCD ID: B9D) (formula: C₆H₁₁FO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	F	O	0	0
			12	6	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	465	Total	O	0	0
			465	465		
6	B	259	Total	O	0	0
			259	259		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ALPHA-GLUCOSIDASE YIHQ



• Molecule 1: ALPHA-GLUCOSIDASE YIHQ



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.67Å 112.64Å 111.88Å 90.00° 109.52° 90.00°	Depositor
Resolution (Å)	105.45 – 1.85 105.45 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.3 (105.45-1.85) 98.4 (105.45-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.57 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0095	Depositor
R, R_{free}	0.160 , 0.193 0.172 , 0.200	Depositor DCC
R_{free} test set	7578 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	27.1	Xtriage
Anisotropy	0.048	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11534	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MPD, B9D, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	17/5533 (0.3%)	1.13	7/7519 (0.1%)
1	B	1.14	9/5562 (0.2%)	1.04	8/7560 (0.1%)
All	All	1.29	26/11095 (0.2%)	1.09	15/15079 (0.1%)

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	541	ARG	N-CA	7.08	1.55	1.46
1	A	551	ASP	C-O	6.65	1.30	1.24
1	A	484	ALA	CA-CB	-6.31	1.43	1.53
1	A	84	SER	N-CA	-6.28	1.38	1.46
1	A	499	GLY	C-O	-6.24	1.19	1.24
1	B	462	GLN	CA-C	6.12	1.60	1.52
1	A	631	PRO	CA-C	6.08	1.59	1.52
1	A	122	ALA	CA-C	6.02	1.60	1.52
1	A	472	ASP	CG-OD2	5.80	1.36	1.25
1	B	647	GLY	C-O	-5.79	1.18	1.23
1	B	469	TRP	CA-C	5.58	1.59	1.52
1	B	70	ALA	CA-C	5.57	1.59	1.52
1	A	418	ASN	C-O	-5.47	1.16	1.24
1	A	611	ARG	CZ-NH1	5.47	1.40	1.32
1	B	399	CYS	N-CA	5.46	1.53	1.46
1	B	676	LYS	CA-C	5.43	1.60	1.52
1	A	376	ASP	CA-C	-5.39	1.46	1.52
1	B	405	ASP	CG-OD2	5.29	1.35	1.25
1	A	421	SER	CA-C	5.25	1.59	1.52
1	A	457	GLY	N-CA	-5.24	1.40	1.45
1	B	525	CYS	C-O	5.18	1.30	1.24
1	A	13	PHE	N-CA	5.07	1.55	1.46
1	A	226	PHE	C-O	5.07	1.30	1.23
1	A	111	ASN	C-O	5.04	1.30	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	261	ILE	C-O	-5.02	1.19	1.24
1	B	657	LYS	N-CA	-5.01	1.41	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	512	PHE	CA-C-N	-7.37	113.40	126.45
1	B	512	PHE	C-N-CA	-7.37	113.40	126.45
1	B	50	MET	CG-SD-CE	-6.33	86.98	100.90
1	A	497	GLY	N-CA-C	6.02	123.92	115.43
1	A	653	ALA	CA-C-N	-5.97	114.12	120.03
1	A	653	ALA	C-N-CA	-5.97	114.12	120.03
1	B	79	TRP	N-CA-C	5.78	119.03	110.48
1	B	343	ASP	CB-CA-C	-5.75	101.29	110.84
1	B	15	GLN	N-CA-C	5.62	117.41	111.28
1	A	295	MET	CG-SD-CE	-5.53	88.75	100.90
1	A	343	ASP	CB-CA-C	-5.46	102.31	110.88
1	A	170	LYS	N-CA-CB	-5.41	101.77	110.69
1	B	464	TYR	N-CA-C	5.28	119.77	113.23
1	B	343	ASP	N-CA-C	5.15	119.81	111.37
1	A	19	SER	N-CA-C	5.01	118.29	110.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5371	0	5090	8	0
1	B	5398	0	5107	31	0
2	A	2	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	16	0	28	0	0
4	B	8	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	12	0	8	1	0
6	A	465	0	0	3	0
6	B	259	0	0	1	0
All	All	11534	0	10247	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:512:PHE:O	1:B:513:GLU:CB	2.37	0.71
1:B:13:PHE:CD1	1:B:72:VAL:HG21	2.29	0.68
1:B:13:PHE:CD1	1:B:72:VAL:CG2	2.77	0.68
1:B:13:PHE:CE1	1:B:72:VAL:HG23	2.31	0.66
1:B:301:ARG:NH1	6:B:2163:HOH:O	2.27	0.63
1:B:251:ASP:OD1	1:B:573:LYS:NZ	2.28	0.62
1:B:512:PHE:O	1:B:513:GLU:HB2	2.00	0.60
1:B:13:PHE:CG	1:B:72:VAL:HG21	2.37	0.59
1:A:21:THR:HG22	1:A:31:THR:HG22	1.88	0.56
1:B:295:MET:HE1	1:B:300:LYS:HG2	1.87	0.56
1:B:57:ILE:HD11	1:B:149:TRP:CD1	2.41	0.55
1:A:52:ARG:HD2	1:A:171:GLU:OE2	2.08	0.53
1:A:619:HIS:HD2	6:A:2412:HOH:O	1.92	0.53
1:A:648:GLU:HG2	6:A:2197:HOH:O	2.08	0.52
1:B:13:PHE:CZ	1:B:72:VAL:HG23	2.45	0.52
1:B:331:LEU:HA	1:B:401:GLY:O	2.10	0.51
1:B:275:MET:HE3	1:B:280:VAL:HG21	1.93	0.49
1:B:494:ALA:HB1	1:B:585:PRO:HG3	1.95	0.48
1:B:50:MET:HE1	1:B:409:TYR:O	2.14	0.47
1:B:301:ARG:NH2	1:B:367:PHE:O	2.47	0.47
1:A:64:LYS:HE3	1:A:118:TRP:CE2	2.50	0.47
1:B:13:PHE:CE1	1:B:72:VAL:CG2	2.98	0.46
1:B:13:PHE:CD1	1:B:72:VAL:HG23	2.48	0.46
1:B:515:LYS:HG2	1:B:544:ASP:HB3	1.98	0.46
1:B:13:PHE:CG	1:B:72:VAL:CG2	2.99	0.45
1:B:388:GLU:OE1	1:B:392:LYS:HE3	2.17	0.44
1:A:366:GLU:HG2	1:A:371:TYR:CE2	2.52	0.44
1:B:185:PRO:HG3	1:B:203:MET:HE1	2.00	0.44
1:B:557:HIS:CE1	1:B:655:ILE:HG22	2.52	0.44
1:A:300:LYS:HE3	6:A:2263:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:508:TYR:OH	5:B:1681:B9D:H6A	2.19	0.43
1:B:506:GLY:H	1:B:524:TRP:CG	2.37	0.43
1:A:8:LEU:HD23	1:A:9:LEU:HD12	2.01	0.42
1:B:194:TYR:CZ	1:B:227:GLU:HB2	2.55	0.42
1:B:597:ASP:OD2	1:B:629:TYR:OH	2.28	0.42
1:B:453:PHE:HA	1:B:467:MET:O	2.21	0.41
1:B:468:MET:HG2	1:B:498:HIS:CE1	2.56	0.41
1:B:304:TRP:HA	1:B:306:TRP:CH2	2.56	0.40
1:B:72:VAL:HG12	1:B:73:SER:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	664/686 (97%)	646 (97%)	18 (3%)	0	100	100
1	B	669/686 (98%)	644 (96%)	23 (3%)	2 (0%)	36	25
All	All	1333/1372 (97%)	1290 (97%)	41 (3%)	2 (0%)	43	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	171	GLU
1	B	475	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	562/580 (97%)	546 (97%)	16 (3%)	38	23
1	B	565/580 (97%)	549 (97%)	16 (3%)	38	23
All	All	1127/1160 (97%)	1095 (97%)	32 (3%)	38	23

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	LEU
1	A	9	LEU
1	A	13	PHE
1	A	26	GLN
1	A	34	LYS
1	A	94	ASN
1	A	107	LEU
1	A	181	PHE
1	A	301	ARG
1	A	307	LYS
1	A	369	GLU
1	A	392	LYS
1	A	412	THR
1	A	469	TRP
1	A	628	LEU
1	A	678	ILE
1	B	9	LEU
1	B	31	THR
1	B	58	LYS
1	B	94	ASN
1	B	99	ASP
1	B	221	LYS
1	B	276	ARG
1	B	307	LYS
1	B	392	LYS
1	B	393	ASN
1	B	432	LEU
1	B	469	TRP
1	B	513	GLU
1	B	581	LYS
1	B	596	ASP
1	B	628	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	GLN
1	A	62	GLN
1	A	82	HIS
1	A	417	HIS
1	A	473	GLN
1	B	12	GLN
1	B	15	GLN
1	B	36	ASN
1	B	54	ASN
1	B	109	ASN
1	B	166	GLN
1	B	262	GLN
1	B	288	GLN
1	B	579	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 5 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	MPD	B	1680	-	7,7,7	0.58	0	9,10,10	0.95	0
4	MPD	A	1683	-	7,7,7	0.38	0	9,10,10	1.32	1 (11%)
4	MPD	A	1682	-	7,7,7	0.47	0	9,10,10	0.94	0
5	B9D	B	1681	1	9,12,13	1.85	1 (11%)	14,18,20	1.47	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MPD	B	1680	-	-	0/5/5/5	-
4	MPD	A	1683	-	-	3/5/5/5	-
4	MPD	A	1682	-	-	0/5/5/5	-
5	B9D	B	1681	1	-	0/2/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1681	B9D	O5-C1	4.99	1.47	1.37

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1681	B9D	O3-C3-C4	2.41	114.97	110.05
5	B	1681	B9D	C1-C2-C3	-2.21	108.28	110.71
5	B	1681	B9D	O5-C5-C4	-2.17	107.60	111.06
4	A	1683	MPD	O2-C2-CM	2.16	114.72	107.99

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1683	MPD	C2-C3-C4-C5
4	A	1683	MPD	O2-C2-C3-C4
4	A	1683	MPD	CM-C2-C3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1681	B9D	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	667/686 (97%)	-0.25	13 (1%) 66 70	15, 26, 51, 87	2 (0%)
1	B	670/686 (97%)	0.36	20 (2%) 52 56	19, 41, 64, 97	1 (0%)
All	All	1337/1372 (97%)	0.06	33 (2%) 58 62	15, 33, 61, 97	3 (0%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	13	PHE	12.7
1	B	298	PHE	6.3
1	B	8	LEU	4.9
1	B	13	PHE	4.4
1	B	677	SER	4.0
1	A	8	LEU	3.8
1	A	97	ALA	3.4
1	A	298	PHE	3.4
1	A	18	ASP	3.0
1	A	678	ILE	2.8
1	A	76	PRO	2.5
1	B	76	PRO	2.5
1	A	9	LEU	2.5
1	B	649	VAL	2.5
1	B	664	ALA	2.5
1	B	14	HIS	2.4
1	A	170	LYS	2.3
1	A	19	SER	2.3
1	B	384	ALA	2.3
1	B	299	GLY	2.3
1	B	70	ALA	2.2
1	B	316	LEU	2.2
1	B	367	PHE	2.2
1	A	359	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	74	GLN	2.2
1	B	15	GLN	2.2
1	B	72	VAL	2.2
1	B	16	ASN	2.1
1	B	307	LYS	2.1
1	B	86	GLY	2.1
1	A	370	PHE	2.0
1	B	297	SER	2.0
1	B	395	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MPD	B	1680	8/8	0.84	0.18	68,72,76,77	0
4	MPD	A	1683	8/8	0.87	0.21	59,65,66,75	0
4	MPD	A	1682	8/8	0.93	0.13	45,53,62,69	0
5	B9D	B	1681	12/13	0.93	0.09	32,38,46,50	0
2	CA	A	1679	1/1	0.98	0.09	33,33,33,33	0
3	CL	A	1681	1/1	0.99	0.06	28,28,28,28	0
3	CL	B	1679	1/1	0.99	0.07	39,39,39,39	0
2	CA	A	1680	1/1	0.99	0.02	18,18,18,18	0
2	CA	B	1678	1/1	1.00	0.01	27,27,27,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.