



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 14, 2026 – 04:53 PM UTC

PDB ID : 5AEX / pdb_00005aex
Title : Crystal structure of *Saccharomyces cerevisiae* Mep2
Authors : Rutherford, J.C.; Chembath, A.; van den Berg, B.
Deposited on : 2015-01-12
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	505	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 72%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">2% 72% 15% • 10%</p>
1	B	505	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 65%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">2% 65% 22% • 10%</p>
1	C	505	<div style="display: flex; align-items: center;"> <div style="width: 6%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 68%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 19%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">6% 68% 19% • 10%</p>
1	D	505	<div style="display: flex; align-items: center;"> <div style="width: 2%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 70%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">2% 70% 17% • 10%</p>
1	E	505	<div style="display: flex; align-items: center;"> <div style="width: 5%; height: 10px; background-color: red; margin-right: 2px;"></div> <div style="width: 66%; height: 10px; background-color: green; margin-right: 2px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: orange; margin-right: 2px;"></div> <div style="width: 10%; height: 10px; background-color: grey;"></div> </div> <p style="text-align: center;">5% 66% 22% • 10%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	505	<p>3% 69% 19% 10%</p>
1	H	505	<p>4% 67% 17% 13%</p>
1	I	505	<p>6% 69% 18% 11%</p>
1	J	505	<p>8% 66% 17% 15%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	I	1455	-	-	X	-

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 30576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMMONIUM TRANSPORTER MEP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	456	3437	2242	561	608	26	41	0	0
1	B	456	3437	2242	561	608	26	65	0	0
1	C	456	3437	2242	561	608	26	69	0	0
1	D	455	3431	2239	560	606	26	60	0	0
1	E	456	3437	2242	561	608	26	95	0	0
1	F	456	3437	2242	561	608	26	57	0	0
1	H	440	3320	2175	541	578	26	117	0	0
1	I	447	3375	2206	549	594	26	204	0	0
1	J	427	3220	2115	519	561	25	258	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	500	HIS	-	expression tag	UNP P41948
A	501	HIS	-	expression tag	UNP P41948
A	502	HIS	-	expression tag	UNP P41948
A	503	HIS	-	expression tag	UNP P41948
A	504	HIS	-	expression tag	UNP P41948
A	505	HIS	-	expression tag	UNP P41948
B	500	HIS	-	expression tag	UNP P41948
B	501	HIS	-	expression tag	UNP P41948
B	502	HIS	-	expression tag	UNP P41948
B	503	HIS	-	expression tag	UNP P41948
B	504	HIS	-	expression tag	UNP P41948

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	505	HIS	-	expression tag	UNP P41948
C	500	HIS	-	expression tag	UNP P41948
C	501	HIS	-	expression tag	UNP P41948
C	502	HIS	-	expression tag	UNP P41948
C	503	HIS	-	expression tag	UNP P41948
C	504	HIS	-	expression tag	UNP P41948
C	505	HIS	-	expression tag	UNP P41948
D	500	HIS	-	expression tag	UNP P41948
D	501	HIS	-	expression tag	UNP P41948
D	502	HIS	-	expression tag	UNP P41948
D	503	HIS	-	expression tag	UNP P41948
D	504	HIS	-	expression tag	UNP P41948
D	505	HIS	-	expression tag	UNP P41948
E	500	HIS	-	expression tag	UNP P41948
E	501	HIS	-	expression tag	UNP P41948
E	502	HIS	-	expression tag	UNP P41948
E	503	HIS	-	expression tag	UNP P41948
E	504	HIS	-	expression tag	UNP P41948
E	505	HIS	-	expression tag	UNP P41948
F	500	HIS	-	expression tag	UNP P41948
F	501	HIS	-	expression tag	UNP P41948
F	502	HIS	-	expression tag	UNP P41948
F	503	HIS	-	expression tag	UNP P41948
F	504	HIS	-	expression tag	UNP P41948
F	505	HIS	-	expression tag	UNP P41948
H	500	HIS	-	expression tag	UNP P41948
H	501	HIS	-	expression tag	UNP P41948
H	502	HIS	-	expression tag	UNP P41948
H	503	HIS	-	expression tag	UNP P41948
H	504	HIS	-	expression tag	UNP P41948
H	505	HIS	-	expression tag	UNP P41948
I	500	HIS	-	expression tag	UNP P41948
I	501	HIS	-	expression tag	UNP P41948
I	502	HIS	-	expression tag	UNP P41948
I	503	HIS	-	expression tag	UNP P41948
I	504	HIS	-	expression tag	UNP P41948
I	505	HIS	-	expression tag	UNP P41948
J	500	HIS	-	expression tag	UNP P41948
J	501	HIS	-	expression tag	UNP P41948
J	502	HIS	-	expression tag	UNP P41948
J	503	HIS	-	expression tag	UNP P41948
J	504	HIS	-	expression tag	UNP P41948

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	505	HIS	-	expression tag	UNP P41948


- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
2	B	1	5	4	1	0	0
2	C	1	5	4	1	0	0
2	D	1	5	4	1	0	0
2	E	1	5	4	1	0	0
2	F	1	5	4	1	0	0
2	H	1	5	4	1	0	0
2	I	1	5	4	1	0	0
2	J	1	5	4	1	0	0




• Molecule 1: AMMONIUM TRANSPORTER MEP2




• Molecule 1: AMMONIUM TRANSPORTER MEP2







• Molecule 1: AMMONIUM TRANSPORTER MEP2




• Molecule 1: AMMONIUM TRANSPORTER MEP2



● Molecule 1: AMMONIUM TRANSPORTER MEP2



● Molecule 1: AMMONIUM TRANSPORTER MEP2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	103.69Å 232.35Å 279.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.28 – 3.20 49.28 – 3.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.28-3.20) 95.6 (49.28-3.20)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 3.19Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.189 , 0.241 0.199 , 0.252	Depositor DCC
R_{free} test set	2453 reflections (2.19%)	wwPDB-VP
Wilson B-factor (Å ²)	85.6	Xtrriage
Anisotropy	0.267	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	30576	wwPDB-VP
Average B, all atoms (Å ²)	84.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/3534	0.94	3/4818 (0.1%)
1	B	0.83	0/3534	0.93	8/4818 (0.2%)
1	C	0.73	0/3534	0.89	7/4818 (0.1%)
1	D	0.85	0/3528	0.89	5/4810 (0.1%)
1	E	0.74	0/3534	0.90	2/4818 (0.0%)
1	F	0.90	0/3534	0.93	6/4818 (0.1%)
1	H	0.69	0/3415	0.86	7/4654 (0.2%)
1	I	0.56	0/3470	0.84	4/4729 (0.1%)
1	J	0.50	0/3313	0.83	5/4518 (0.1%)
All	All	0.76	0/31396	0.89	47/42801 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 47 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	340	ASP	N-CA-C	10.56	125.98	112.89
1	C	304	THR	CA-C-N	-7.90	111.66	119.87
1	C	304	THR	C-N-CA	-7.90	111.66	119.87
1	I	240	PHE	N-CA-C	7.23	118.81	111.07
1	F	367	VAL	N-CA-C	-7.01	106.22	111.90

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	240	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3437	0	3401	44	0
1	B	3437	0	3401	75	0
1	C	3437	0	3401	62	1
1	D	3431	0	3396	53	0
1	E	3437	0	3401	67	0
1	F	3437	0	3401	53	0
1	H	3320	0	3295	51	0
1	I	3375	0	3337	57	1
1	J	3220	0	3188	54	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	1	0
2	F	5	0	0	0	0
2	H	5	0	0	0	0
2	I	5	0	0	2	0
2	J	5	0	0	0	0
All	All	30576	0	30221	495	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 495 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:140:MET:HE1	1:B:200:GLY:HA3	1.55	0.88
1:J:10:THR:HG22	1:J:12:GLU:H	1.37	0.88
1:F:256:ARG:NH1	1:F:308:GLY:O	2.10	0.84

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:THR:HG22	1:A:12:GLU:H	1.47	0.78
1:C:400:ALA:O	1:C:404:THR:OG1	2.01	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:12:GLU:OE2	1:I:387:TYR:OH[4_545]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	454/505 (90%)	429 (94%)	21 (5%)	4 (1%)	14 47
1	B	454/505 (90%)	422 (93%)	29 (6%)	3 (1%)	18 52
1	C	454/505 (90%)	421 (93%)	29 (6%)	4 (1%)	14 47
1	D	453/505 (90%)	419 (92%)	29 (6%)	5 (1%)	11 43
1	E	454/505 (90%)	421 (93%)	28 (6%)	5 (1%)	11 43
1	F	454/505 (90%)	429 (94%)	24 (5%)	1 (0%)	43 73
1	H	434/505 (86%)	406 (94%)	20 (5%)	8 (2%)	6 34
1	I	441/505 (87%)	402 (91%)	34 (8%)	5 (1%)	11 43
1	J	421/505 (83%)	387 (92%)	32 (8%)	2 (0%)	24 59
All	All	4019/4545 (88%)	3736 (93%)	246 (6%)	37 (1%)	14 47

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	112	GLY
1	D	435	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	I	112	GLY
1	A	95	GLY
1	B	95	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	353/398 (89%)	321 (91%)	32 (9%)	9	34
1	B	353/398 (89%)	329 (93%)	24 (7%)	14	46
1	C	353/398 (89%)	327 (93%)	26 (7%)	13	43
1	D	352/398 (88%)	324 (92%)	28 (8%)	11	40
1	E	353/398 (89%)	328 (93%)	25 (7%)	13	44
1	F	353/398 (89%)	326 (92%)	27 (8%)	12	42
1	H	341/398 (86%)	317 (93%)	24 (7%)	14	45
1	I	346/398 (87%)	327 (94%)	19 (6%)	19	52
1	J	330/398 (83%)	315 (96%)	15 (4%)	24	58
All	All	3134/3582 (88%)	2914 (93%)	220 (7%)	14	45

5 of 220 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	73	SER
1	F	101	THR
1	J	447	SER
1	I	242	TRP
1	E	117	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	416	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	246	ASN
1	F	265	ASN
1	H	386	HIS
1	C	199	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	1458	-	4,4,4	1.24	0	6,6,6	0.60	0
2	PO4	C	1458	-	4,4,4	1.12	0	6,6,6	0.46	0
2	PO4	J	1455	-	4,4,4	0.99	0	6,6,6	0.51	0
2	PO4	B	1458	-	4,4,4	1.06	0	6,6,6	0.53	0
2	PO4	F	1458	-	4,4,4	1.20	0	6,6,6	0.60	0
2	PO4	H	1457	-	4,4,4	1.00	0	6,6,6	0.42	0
2	PO4	I	1455	-	4,4,4	1.00	0	6,6,6	0.43	0
2	PO4	E	1458	-	4,4,4	1.17	0	6,6,6	0.42	0
2	PO4	D	1458	-	4,4,4	1.09	0	6,6,6	0.49	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	1455	PO4	2	0
2	E	1458	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/505 (90%)	-0.27	10 (2%) 62 42	27, 62, 90, 173	11 (2%)
1	B	456/505 (90%)	-0.09	12 (2%) 57 37	38, 71, 102, 170	18 (3%)
1	C	456/505 (90%)	0.07	29 (6%) 25 16	32, 79, 119, 155	19 (4%)
1	D	455/505 (90%)	-0.17	8 (1%) 67 48	33, 69, 102, 158	16 (3%)
1	E	456/505 (90%)	0.04	26 (5%) 29 19	36, 74, 107, 207	25 (5%)
1	F	456/505 (90%)	-0.14	14 (3%) 51 32	31, 63, 93, 158	15 (3%)
1	H	440/505 (87%)	0.14	21 (4%) 35 22	36, 81, 130, 160	31 (7%)
1	I	447/505 (88%)	0.24	28 (6%) 26 17	54, 98, 142, 188	55 (12%)
1	J	426/505 (84%)	0.42	38 (8%) 15 10	42, 110, 150, 167	71 (16%)
All	All	4048/4545 (89%)	0.02	186 (4%) 37 23	27, 75, 130, 207	261 (6%)

The worst 5 of 186 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	334	SER	10.7
1	E	376	SER	7.1
1	C	368	ASN	6.2
1	E	377	PRO	6.1
1	A	447	SER	6.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	PO4	H	1457	5/5	0.70	0.10	200,202,202,204	0
2	PO4	I	1455	5/5	0.76	0.10	178,181,181,182	0
2	PO4	J	1455	5/5	0.84	0.07	163,163,163,163	0
2	PO4	A	1458	5/5	0.91	0.09	98,101,104,105	0
2	PO4	D	1458	5/5	0.92	0.12	142,143,144,147	0
2	PO4	B	1458	5/5	0.92	0.09	99,100,101,102	0
2	PO4	F	1458	5/5	0.94	0.10	97,97,100,104	0
2	PO4	C	1458	5/5	0.96	0.06	110,110,111,112	0
2	PO4	E	1458	5/5	0.96	0.08	113,113,114,114	0

6.5 Other polymers [i](#)

There are no such residues in this entry.