



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:37 PM UTC

PDB ID : 8AE4 / pdb_00008ae4
Title : Crystal structure of human legumain in complex with Clitocypin 2
Authors : Elamin, T.; Brandstetter, H.; Dall, E.
Deposited on : 2022-07-12
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	
2	B	160	
3	C	2	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	SCH	A	189	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Legumain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	20	0	0
			2116	1336	361	403	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	SNN	ASP	modified residue	UNP Q99538
A	263	GLN	ASN	conflict	UNP Q99538

- Molecule 2 is a protein called Clitocypin-2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	151	Total	C	N	O	2	1	0
			1199	766	204	229			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	153	LEU	-	expression tag	UNP Q3Y9I4
B	154	GLU	-	expression tag	UNP Q3Y9I4
B	155	HIS	-	expression tag	UNP Q3Y9I4
B	156	HIS	-	expression tag	UNP Q3Y9I4
B	157	HIS	-	expression tag	UNP Q3Y9I4
B	158	HIS	-	expression tag	UNP Q3Y9I4
B	159	HIS	-	expression tag	UNP Q3Y9I4
B	160	HIS	-	expression tag	UNP Q3Y9I4

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	201	Total	O	0	0
			201	201		
5	B	122	Total	O	0	0
			122	122		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.47Å 63.63Å 85.53Å 90.00° 102.06° 90.00°	Depositor
Resolution (Å)	41.53 – 1.79 41.53 – 1.79	Depositor EDS
% Data completeness (in resolution range)	97.0 (41.53-1.79) 96.7 (41.53-1.79)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874, PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.192 , 0.208 0.192 , 0.208	Depositor DCC
R_{free} test set	2001 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	22.9	Xtriage
Anisotropy	0.260	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3694	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SNN, SCH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2155	0.57	0/2919
2	B	0.36	0/1236	0.59	0/1689
All	All	0.36	0/3391	0.57	0/4608

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2116	0	1997	32	0
2	B	1199	0	1152	34	0
3	C	28	0	25	2	0
4	A	28	0	26	0	0
5	A	201	0	0	3	0
5	B	122	0	0	3	0
All	All	3694	0	3200	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:ASP:O	1:A:235:VAL:HG22	1.78	0.83
2:B:10:ARG:HH12	2:B:47:PHE:HB2	1.45	0.81
1:A:189:SCH:HE3	1:A:216:SER:H	1.47	0.77
1:A:189:SCH:HE1	2:B:70:ASN:H	1.52	0.73
1:A:259:LYS:HE3	5:A:482:HOH:O	1.89	0.72
2:B:53[B]:GLN:OE1	2:B:67:GLN:NE2	2.23	0.71
2:B:106:ASP:OD2	2:B:148:ARG:NH2	2.23	0.71
3:C:1:NAG:H62	3:C:2:NAG:H61	1.74	0.69
2:B:44:PRO:HA	2:B:47:PHE:CE1	2.28	0.68
2:B:38:LYS:NZ	5:B:201:HOH:O	2.23	0.68
1:A:189:SCH:SD	1:A:215:SER:HB3	2.39	0.63
2:B:150:THR:HG23	5:B:212:HOH:O	2.01	0.61
1:A:189:SCH:HE1	2:B:70:ASN:N	2.16	0.61
2:B:44:PRO:HA	2:B:47:PHE:CZ	2.36	0.60
2:B:104:THR:CB	2:B:107:VAL:HG13	2.31	0.60
1:A:189:SCH:HE2	2:B:70:ASN:C	2.28	0.58
1:A:189:SCH:SD	1:A:215:SER:CB	2.93	0.57
2:B:8:ILE:CD1	2:B:53[A]:GLN:NE2	2.70	0.54
2:B:8:ILE:CD1	2:B:53[A]:GLN:HE21	2.21	0.53
2:B:43:THR:HB	2:B:44:PRO:CD	2.38	0.53
1:A:189:SCH:HE3	1:A:216:SER:N	2.22	0.52
1:A:189:SCH:HB3	2:B:70:ASN:O	2.10	0.52
2:B:10:ARG:NH1	2:B:47:PHE:HB2	2.21	0.51
2:B:104:THR:OG1	2:B:107:VAL:HG13	2.11	0.51
1:A:207:THR:HA	1:A:268:MET:O	2.12	0.50
1:A:219:CYS:HB3	1:A:232:TRP:CD1	2.48	0.49
1:A:65:GLN:HE22	1:A:286:MET:HE1	1.77	0.49
2:B:102:PRO:O	2:B:103:SER:HB3	2.13	0.48
1:A:223:GLU:HG2	2:B:152:GLU:O	2.13	0.48
2:B:150:THR:CG2	5:B:212:HOH:O	2.59	0.47
1:A:131:LYS:NZ	5:A:408:HOH:O	2.47	0.46
2:B:120:ASP:OD2	2:B:122:GLU:OE2	2.32	0.46
1:A:111:GLN:H	1:A:111:GLN:CD	2.24	0.46
2:B:8:ILE:HD13	2:B:53[A]:GLN:NE2	2.31	0.45
1:A:249:GLU:HG3	1:A:253:LYS:HD3	1.99	0.44
2:B:23:VAL:HG22	2:B:24:GLY:N	2.32	0.44
2:B:8:ILE:HD13	2:B:53[A]:GLN:HE21	1.82	0.44
1:A:189:SCH:HE2	2:B:70:ASN:O	2.17	0.44
2:B:44:PRO:N	2:B:45:PRO:CD	2.81	0.43
1:A:259:LYS:CE	5:A:482:HOH:O	2.58	0.43
1:A:85:ILE:HD12	1:A:94:ASP:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:189:SCH:HE2	1:A:189:SCH:HB3	1.69	0.43
2:B:71:THR:HG21	2:B:75:TYR:CE2	2.53	0.43
2:B:8:ILE:HD11	2:B:53[A]:GLN:NE2	2.33	0.43
1:A:27:GLY:HA3	1:A:65:GLN:OE1	2.19	0.43
2:B:114:ILE:HD13	2:B:121:VAL:HG23	2.01	0.43
3:C:2:NAG:O7	3:C:2:NAG:O3	2.30	0.43
1:A:221:TYR:CG	2:B:51:ILE:HD13	2.53	0.42
1:A:147:SNN:O5	1:A:148:HIS:HB2	2.19	0.42
2:B:8:ILE:HD11	2:B:53[A]:GLN:HE21	1.85	0.42
2:B:12:ARG:HB2	2:B:27:TYR:CE2	2.54	0.42
1:A:217:TYR:CD2	1:A:264:THR:HB	2.54	0.42
1:A:53:TYR:CZ	1:A:100:PRO:HD3	2.55	0.42
1:A:222:ASP:OD2	1:A:225:ARG:HD2	2.19	0.42
1:A:67:VAL:HG22	1:A:131:LYS:HD2	2.03	0.41
1:A:56:ILE:HG23	1:A:61:ILE:HD12	2.01	0.41
2:B:59:ASP:OD2	2:B:95:GLU:OE1	2.39	0.41
1:A:281:MET:O	1:A:281:MET:HG3	2.21	0.41
1:A:196:ASN:HA	1:A:270:TYR:CD1	2.55	0.40
1:A:70:MET:O	1:A:104:THR:HA	2.21	0.40
2:B:23:VAL:HG22	2:B:24:GLY:H	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	258/263 (98%)	256 (99%)	2 (1%)	0	100	100
2	B	150/160 (94%)	145 (97%)	5 (3%)	0	100	100
All	All	408/423 (96%)	401 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/228 (100%)	227 (100%)	0	100	100
2	B	126/134 (94%)	124 (98%)	2 (2%)	55	47
All	All	353/362 (98%)	351 (99%)	2 (1%)	78	77

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	104	THR
2	B	107	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	GLN
1	A	88	ASN
1	A	176	HIS
1	A	254	GLN
1	A	284	GLN
2	B	67	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SCH	A	189	1	6,7,8	0.66	0	4,7,9	0.95	0
1	SNN	A	147	1	7,8,8	1.89	1 (14%)	10,11,11	2.35	5 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SCH	A	189	1	-	1/2/6/8	-
1	SNN	A	147	1	-	-	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	147	SNN	C-N1	-4.44	1.31	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	147	SNN	O5-C5-C4	-3.64	121.84	126.48
1	A	147	SNN	O-C-CA	-3.34	123.79	126.19
1	A	147	SNN	CA-C-N1	3.09	109.72	107.30
1	A	147	SNN	C4-C5-N1	2.47	110.95	108.02
1	A	147	SNN	CA-C4-C5	-2.38	100.76	103.55

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	189	SCH	N-CA-CB-SG

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	189	SCH	10	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	147	SNN	1	0

5.5 Carbohydrates

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAG	C	1	3,1	14,14,15	0.55	0	17,19,21	1.28	2 (11%)
3	NAG	C	2	3	14,14,15	0.69	0	17,19,21	1.79	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	4/6/23/26	0/1/1/1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	2	NAG	C1-O5-C5	5.36	119.37	112.19
3	C	1	NAG	C4-C3-C2	-3.31	106.16	111.02
3	C	2	NAG	C1-C2-N2	-2.68	106.20	110.43
3	C	1	NAG	O4-C4-C3	2.27	115.72	110.38
3	C	2	NAG	C2-N2-C7	-2.24	119.90	122.90

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	C4-C5-C6-O6
3	C	2	NAG	O7-C7-N2-C2
3	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	NAG	2	0
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	301	1	14,14,15	0.71	0	17,19,21	0.82	0
4	NAG	A	302	1	14,14,15	0.54	0	17,19,21	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	301	1	-	2/6/23/26	0/1/1/1
4	NAG	A	302	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	NAG	C8-C7-N2-C2
4	A	301	NAG	O7-C7-N2-C2
4	A	302	NAG	O5-C5-C6-O6
4	A	302	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	259/263 (98%)	0.10	6 (2%) 61 61	15, 24, 37, 60	1 (0%)
2	B	151/160 (94%)	0.33	7 (4%) 37 36	15, 25, 43, 52	3 (1%)
All	All	410/423 (96%)	0.18	13 (3%) 50 50	15, 24, 39, 60	4 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	GLY	6.3
2	B	23	VAL	4.1
2	B	24	GLY	3.7
1	A	41	TYR	2.9
1	A	287	LYS	2.9
2	B	2	ALA	2.8
2	B	152	GLU	2.7
2	B	142	GLY	2.7
2	B	3	SER	2.6
2	B	44	PRO	2.5
1	A	286	MET	2.5
1	A	28	LYS	2.1
1	A	235	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SCH	A	189	8/9	0.79	0.18	20,23,45,48	0

Continued on next page...

Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	SNN	A	147	8/8	0.96	0.06	15,18,19,19	0

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	C	2	14/15	0.71	0.19	40,47,52,57	0
3	NAG	C	1	14/15	0.91	0.10	21,25,37,43	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	302	14/15	0.84	0.10	35,42,47,49	0
4	NAG	A	301	14/15	0.89	0.10	31,42,48,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.