



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:36 AM UTC

PDB ID : 2AGW / pdb_00002agw
Title : Crystal structure of tryptamine-reduced aromatic amine dehydrogenase (AADH) from *Alcaligenes faecalis* in complex with tryptamine
Authors : Masgrau, L.; Roujeinikova, A.; Johannissen, L.O.; Hothi, P.; Basran, J.; Ranaghan, K.E.; Mulholland, A.J.; Sutcliffe, M.J.; Scrutton, N.S.; Leys, D.
Deposited on : 2005-07-27
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1756 (1.46-1.46)
Clashscore	190562	1795 (1.46-1.46)
Ramachandran outliers	187476	1776 (1.46-1.46)
Sidechain outliers	187428	1776 (1.46-1.46)
RSRZ outliers	180081	1756 (1.46-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	135	% 70% 10% • 19%
1	H	135	4% 73% 14% • 12%
2	A	361	% 91% 9% •
2	B	361	% 93% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8603 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aromatic amine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	109	Total	C	N	O	S	0	0	0
			836	510	148	163	15			
1	H	119	Total	C	N	O	S	0	5	0
			925	565	162	183	15			

- Molecule 2 is a protein called Aromatic amine dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	360	Total	C	N	O	S	0	2	0
			2811	1773	491	533	14			
2	B	361	Total	C	N	O	S	0	2	0
			2799	1764	488	533	14			

- Molecule 3 is 2-(1H-INDOL-3-YL)ETHANAMINE (CCD ID: TSS) (formula: C₁₀H₁₂N₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	N	0	0
			12	10	2		
3	H	1	Total	C	N	0	0
			12	10	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	128	Total	O	0	0
			128	128		
4	H	114	Total	O	0	0
			114	114		
4	A	534	Total	O	0	0
			534	534		
4	B	432	Total	O	0	0
			432	432		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aromatic amine dehydrogenase



- Molecule 1: Aromatic amine dehydrogenase



- Molecule 2: Aromatic amine dehydrogenase



- Molecule 2: Aromatic amine dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.00Å 88.75Å 81.15Å 90.00° 89.99° 90.00°	Depositor
Resolution (Å)	15.00 – 1.45 15.00 – 1.45	Depositor EDS
% Data completeness (in resolution range)	95.6 (15.00-1.45) 95.5 (15.00-1.45)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.45Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.163 , 0.195 0.165 , 0.197	Depositor DCC
R_{free} test set	8497 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.422	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 49.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8603	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TQQ, TSS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.73	0/842	0.87	0/1146
1	H	0.70	0/944	0.87	0/1289
2	A	0.78	1/2883 (0.0%)	0.84	0/3907
2	B	0.74	0/2872	0.83	0/3895
All	All	0.75	1/7541 (0.0%)	0.84	0/10237

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	172	PRO	CA-C	6.12	1.55	1.51

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	172[A]	THR	Mainchain
1	H	172[B]	THR	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	836	0	733	13	0
1	H	925	0	811	24	0
2	A	2811	0	2752	29	0
2	B	2799	0	2713	25	0
3	D	12	0	12	1	0
3	H	12	0	12	0	0
4	A	534	0	0	4	0
4	B	432	0	0	3	0
4	D	128	0	0	4	0
4	H	114	0	0	6	0
All	All	8603	0	7033	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:173[A]:THR:HA	4:H:2109:HOH:O	1.70	0.92
1:H:128:ASP:OD1	1:H:172[B]:THR:HG23	1.80	0.81
1:H:80:HIS:O	1:H:142[A]:THR:HG21	1.85	0.75
2:A:104:ARG:HH11	2:A:106:HIS:HE1	1.35	0.74
1:D:179:LEU:C	4:D:3050:HOH:O	2.31	0.73
2:B:104:ARG:HH11	2:B:106:HIS:HE1	1.38	0.72
2:A:124:ASN:HD21	2:A:178:GLY:H	1.35	0.71
2:B:124:ASN:HD21	2:B:178:GLY:H	1.37	0.71
1:H:105:SER:HA	1:H:164:ASN:HD21	1.55	0.71
4:H:2051:HOH:O	2:B:118[B]:MET:SD	2.49	0.70
1:D:71:GLU:N	4:D:3097:HOH:O	2.24	0.69
1:D:105:SER:HA	1:D:164:ASN:HD21	1.57	0.69
2:B:91:TYR:OH	2:B:428:HIS:HD2	1.78	0.66
2:A:91:TYR:OH	2:A:428:HIS:HD2	1.81	0.64
2:B:197:GLN:NE2	2:B:227:SER:H	1.98	0.61
2:A:248:LEU:HD13	2:A:263:ARG:HD3	1.81	0.61
2:A:197:GLN:NE2	2:A:227:SER:H	1.99	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:167:SER:HB2	2:A:177:GLN:HE22	1.66	0.60
1:H:176:LEU:HD22	2:B:73:ARG:HD3	1.83	0.60
1:H:80:HIS:O	1:H:142[A]:THR:CG2	2.48	0.60
2:A:335:ARG:H	2:A:386:GLN:HE22	1.48	0.59
1:H:173[B]:THR:OG1	2:B:76:LEU:HD22	2.02	0.59
1:H:159:ASN:HD22	2:A:177:GLN:NE2	2.02	0.57
2:B:335:ARG:H	2:B:386:GLN:HE22	1.52	0.57
1:D:167:SER:HB2	2:B:177:GLN:HE22	1.69	0.56
2:B:106:HIS:HD2	2:B:418:ALA:O	1.89	0.56
2:A:197:GLN:HE21	2:A:227:SER:H	1.52	0.55
2:B:335:ARG:H	2:B:386:GLN:NE2	2.05	0.55
2:A:335:ARG:H	2:A:386:GLN:NE2	2.03	0.55
1:H:173[B]:THR:HG21	2:B:76:LEU:CD1	2.37	0.54
1:H:81:CYS:C	1:H:142[A]:THR:HG23	2.34	0.52
2:A:91:TYR:OH	2:A:428:HIS:CD2	2.62	0.52
2:B:197:GLN:HE21	2:B:227:SER:H	1.58	0.52
3:D:3001:TSS:HA2	4:D:3128:HOH:O	2.09	0.52
2:A:143:HIS:HD2	4:A:606:HOH:O	1.93	0.51
2:A:106:HIS:HD2	2:A:418:ALA:O	1.94	0.51
1:D:109:TQQ:HB2	1:D:160:TRP:NE1	2.25	0.51
1:D:159:ASN:HD22	2:B:177:GLN:NE2	2.10	0.50
1:H:146:GLU:CD	2:A:354:LYS:HZ1	2.19	0.50
2:A:124:ASN:ND2	2:A:178:GLY:H	2.07	0.49
1:D:146:GLU:CD	2:B:354:LYS:HZ1	2.21	0.49
1:H:159:ASN:HD22	2:A:177:GLN:HE21	1.58	0.49
2:A:128:GLN:HE22	2:A:186:ARG:C	2.21	0.49
1:H:109:TQQ:HB2	1:H:160:TRP:NE1	2.30	0.47
1:H:172[A]:THR:OG1	4:H:2036:HOH:O	2.20	0.47
1:H:80:HIS:O	1:H:83[B]:VAL:HG12	2.15	0.47
2:A:271:LYS:NZ	4:A:824:HOH:O	2.47	0.46
2:B:128:GLN:HE22	2:B:186:ARG:C	2.23	0.46
2:A:133:GLY:O	2:A:428:HIS:HE1	1.98	0.46
2:A:179:LEU:HD21	2:A:328:TYR:CE2	2.51	0.45
2:B:124:ASN:ND2	2:B:178:GLY:H	2.10	0.45
2:A:200:SER:N	2:A:201:PRO:HA	2.32	0.45
2:A:275:ILE:HA	2:A:290:SER:HA	1.99	0.45
1:H:155:HIS:HE1	4:A:436:HOH:O	1.99	0.45
2:B:143:HIS:HD2	4:B:503:HOH:O	2.00	0.44
2:B:321:LYS:HD3	4:B:839:HOH:O	2.18	0.44
1:H:173[B]:THR:HG21	2:B:76:LEU:HD13	1.99	0.44
1:D:112:THR:HB	1:D:145:ARG:HD3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:172[B]:THR:CG2	4:H:2036:HOH:O	2.66	0.43
4:D:3025:HOH:O	2:B:143:HIS:HE1	2.02	0.43
1:H:173[B]:THR:HG21	2:B:76:LEU:HD11	2.00	0.43
2:B:73:ARG:HG2	2:B:74:GLU:HG3	2.01	0.43
2:A:236:ARG:HD3	4:A:741:HOH:O	2.19	0.43
2:A:200:SER:N	2:A:201:PRO:CA	2.82	0.42
2:A:231:GLN:NE2	2:A:234:ARG:HH11	2.17	0.42
2:B:275:ILE:HA	2:B:290:SER:HA	2.01	0.42
1:D:155:HIS:HE1	4:B:441:HOH:O	2.02	0.42
1:H:112:THR:HG22	1:H:123:LEU:HD23	2.02	0.42
1:D:127:HIS:HB2	2:A:76:LEU:HD22	2.02	0.41
2:A:131:ASN:H	2:A:187:GLN:HE22	1.68	0.41
1:D:173:THR:HG21	2:A:76:LEU:CD1	2.51	0.41
1:D:157:ASP:CG	2:B:354:LYS:HZ3	2.29	0.41
1:H:77:TYR:CE1	1:H:79:ARG:HG2	2.55	0.41
1:H:172[B]:THR:HG21	4:H:2036:HOH:O	2.20	0.41
1:H:139:GLN:NE2	4:H:2073:HOH:O	2.51	0.40
1:D:173:THR:HG21	2:A:76:LEU:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	106/135 (78%)	104 (98%)	2 (2%)	0	100	100
1	H	121/135 (90%)	118 (98%)	3 (2%)	0	100	100
2	A	360/361 (100%)	348 (97%)	11 (3%)	1 (0%)	36	16
2	B	360/361 (100%)	347 (96%)	12 (3%)	1 (0%)	36	16
All	All	947/992 (96%)	917 (97%)	28 (3%)	2 (0%)	43	21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	216	VAL
2	A	200	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	95/112 (85%)	95 (100%)	0	100	100
1	H	106/112 (95%)	106 (100%)	0	100	100
2	A	305/305 (100%)	304 (100%)	1 (0%)	86	74
2	B	302/305 (99%)	301 (100%)	1 (0%)	86	74
All	All	808/834 (97%)	806 (100%)	2 (0%)	86	76

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	179	LEU
2	B	349	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (36) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	139	GLN
1	D	143	GLN
1	D	155	HIS
1	D	164	ASN
1	H	139	GLN
1	H	143	GLN
1	H	155	HIS
1	H	164	ASN
2	A	86	GLN
2	A	106	HIS
2	A	124	ASN
2	A	128	GLN
2	A	143	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	A	177	GLN
2	A	180	ASN
2	A	187	GLN
2	A	197	GLN
2	A	231	GLN
2	A	251	ASN
2	A	386	GLN
2	A	424	GLN
2	A	426	GLN
2	A	428	HIS
2	B	106	HIS
2	B	124	ASN
2	B	128	GLN
2	B	143	HIS
2	B	177	GLN
2	B	187	GLN
2	B	197	GLN
2	B	231	GLN
2	B	251	ASN
2	B	386	GLN
2	B	424	GLN
2	B	426	GLN
2	B	428	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TQQ	D	109	1	16,17,18	4.35	7 (43%)	14,24,26	0.81	0
1	TQQ	H	109	1	16,17,18	4.25	6 (37%)	14,24,26	1.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TQQ	D	109	1	-	0/5/19/21	0/2/2/2
1	TQQ	H	109	1	-	0/5/19/21	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	109	TQQ	CH2-N2	12.59	1.48	1.28
1	H	109	TQQ	CH2-N2	12.36	1.48	1.28
1	D	109	TQQ	O2-CZ2	9.51	1.44	1.23
1	H	109	TQQ	O2-CZ2	9.00	1.43	1.23
1	D	109	TQQ	CH2-CZ2	-4.82	1.35	1.48
1	H	109	TQQ	CH2-CZ2	-4.60	1.36	1.48
1	H	109	TQQ	CE2-CZ2	-3.31	1.38	1.50
1	D	109	TQQ	CE2-CZ2	-3.29	1.38	1.50
1	H	109	TQQ	CD2-CG	2.76	1.47	1.42
1	D	109	TQQ	CD1-CG	-2.37	1.35	1.38
1	D	109	TQQ	CD2-CG	2.17	1.46	1.42
1	H	109	TQQ	CZ3-CE3	2.16	1.40	1.35
1	D	109	TQQ	CZ3-CH2	-2.00	1.41	1.44

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	109	TQQ	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	H	109	TQQ	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TSS	H	2001	-	13,13,13	1.35	1 (7%)	17,17,17	1.79	3 (17%)
3	TSS	D	3001	-	13,13,13	1.37	2 (15%)	17,17,17	1.93	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TSS	H	2001	-	-	2/3/3/3	0/2/2/2
3	TSS	D	3001	-	-	2/3/3/3	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	2001	TSS	CD2-CG	-4.09	1.37	1.44
3	D	3001	TSS	CD2-CG	-3.73	1.37	1.44
3	D	3001	TSS	CD1-CG	2.10	1.39	1.36

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	3001	TSS	CE3-CD2-CE2	4.46	123.46	118.84
3	H	2001	TSS	CE3-CD2-CE2	4.35	123.34	118.84
3	D	3001	TSS	CZ2-CE2-CD2	-3.98	118.33	122.19
3	H	2001	TSS	CE3-CD2-CG	-3.51	128.62	133.85
3	D	3001	TSS	CE3-CD2-CG	-3.46	128.69	133.85
3	H	2001	TSS	CZ2-CE2-CD2	-3.06	119.22	122.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	3001	TSS	CA-CB-CG-CD2
3	H	2001	TSS	CA-CB-CG-CD2
3	D	3001	TSS	CA-CB-CG-CD1
3	H	2001	TSS	CA-CB-CG-CD1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	3001	TSS	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	108/135 (80%)	-0.00	1 (0%) 81 84	10, 16, 26, 33	0
1	H	118/135 (87%)	0.21	6 (5%) 33 34	7, 17, 39, 50	5 (4%)
2	A	360/361 (99%)	-0.42	3 (0%) 82 86	7, 13, 25, 56	2 (0%)
2	B	361/361 (100%)	0.00	3 (0%) 82 86	8, 18, 30, 48	2 (0%)
All	All	947/992 (95%)	-0.13	13 (1%) 73 77	7, 15, 30, 56	9 (0%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	179	LEU	3.4
2	A	76	LEU	3.1
2	A	75	VAL	2.8
1	H	62	LEU	2.8
2	A	73	ARG	2.7
2	B	316	ASP	2.7
2	B	317	GLU	2.1
1	D	179	LEU	2.1
1	H	111	GLY	2.1
1	H	123	LEU	2.1
2	B	73	ARG	2.0
1	H	69	GLU	2.0
1	H	70	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TQQ	D	109	16/17	0.95	0.06	13,15,19,21	0
1	TQQ	H	109	16/17	0.97	0.05	11,13,17,23	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TSS	D	3001	12/12	0.87	0.10	24,26,29,31	0
3	TSS	H	2001	12/12	0.89	0.09	20,23,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.