



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:54 PM UTC

PDB ID : 3AGX / pdb_00003agx
Title : Crystal structure of human Hsp40 Hdj1 peptide-binding domain
Authors : Suzuki, H.; Noguchi, S.; Satow, Y.
Deposited on : 2010-04-12
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	181	10% 83% 13% ••
1	B	181	16% 69% 13% • 18%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2831 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DnaJ homolog subfamily B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	176	Total	C	N	O	S	0	5	0
			1416	898	254	259	5			
1	B	148	Total	C	N	O	S	0	5	0
			1200	774	212	209	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	MET	-	expression tag	UNP P25685
B	160	MET	-	expression tag	UNP P25685


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	116	Total	O	0	0
			116	116		
2	B	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DnaJ homolog subfamily B member 1



- Molecule 1: DnaJ homolog subfamily B member 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	103.25Å 40.92Å 62.65Å 90.00° 96.62° 90.00°	Depositor
Resolution (Å)	42.03 – 1.85 42.03 – 1.85	Depositor EDS
% Data completeness (in resolution range)	94.3 (42.03-1.85) 94.3 (42.03-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.5.0089	Depositor
R, R_{free}	0.228 , 0.247 0.225 , 0.224	Depositor DCC
R_{free} test set	2156 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.603	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2831	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/1472	0.93	0/1984
1	B	0.98	0/1248	1.01	4/1678 (0.2%)
All	All	0.92	0/2720	0.97	4/3662 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	297	VAL	N-CA-C	-5.97	102.24	108.15
1	B	322	PHE	CA-C-N	-5.22	114.40	119.78
1	B	322	PHE	C-N-CA	-5.22	114.40	119.78
1	B	326	ILE	CB-CA-C	-5.22	104.79	110.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1416	0	1478	21	0
1	B	1200	0	1292	17	0
2	A	116	0	0	0	0
2	B	99	0	0	2	0
All	All	2831	0	2770	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 7.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:ARG:H	1:A:293[A]:MET:HE3	0.92	1.07
1:A:290:ARG:N	1:A:293[A]:MET:HE3	1.76	1.01
1:A:290:ARG:H	1:A:293[A]:MET:CE	1.79	0.94
1:A:290:ARG:CZ	1:A:293[A]:MET:HE2	2.02	0.90
1:A:326:ILE:HG22	1:A:331:ARG:HG3	1.85	0.58
1:A:197:ILE:HD13	1:A:227:THR:HG22	1.87	0.56
1:A:172:LEU:HD21	1:A:249:ARG:HB2	1.90	0.52
1:B:257:PRO:HD2	2:B:47:HOH:O	2.09	0.52
1:A:262:LEU:HD21	1:B:266:LEU:HD11	1.93	0.50
1:B:206:ILE:CG2	1:B:218:ILE:HD13	2.43	0.48
1:B:327:PRO:HB2	1:B:329:THR:HG22	1.96	0.48
1:B:206:ILE:HG23	1:B:218:ILE:HG21	1.95	0.48
1:B:185:ILE:HG12	1:B:204:LEU:HG	1.97	0.47
1:B:329:THR:HG23	1:B:330:SER:N	2.30	0.47
1:B:314:LEU:C	1:B:314:LEU:HD23	2.41	0.46
1:A:294:ARG:HA	1:A:316:ILE:O	2.16	0.45
1:A:290:ARG:N	1:A:293[A]:MET:CE	2.56	0.45
1:B:170:VAL:HG12	1:B:174:GLU:HB2	1.98	0.45
1:A:221:PRO:HA	1:A:234:ASP:OD1	2.18	0.44
1:A:226:GLN:HB2	1:A:232:PRO:HB3	2.00	0.44
1:B:331[B]:ARG:NH2	2:B:146:HOH:O	2.51	0.43
1:B:206:ILE:HG21	1:B:218:ILE:HD13	2.01	0.43
1:A:323:PRO:HG2	1:A:326:ILE:HD11	2.01	0.43
1:A:340:ILE:HD12	1:A:340:ILE:HG21	1.81	0.43
1:B:222:LYS:HE2	1:B:222:LYS:HB3	1.76	0.43
1:B:169:ARG:HB3	1:B:242[A]:LYS:HG2	2.01	0.43
1:B:294:ARG:HA	1:B:316:ILE:O	2.19	0.42
1:A:340:ILE:HG23	1:A:340:ILE:HD13	1.70	0.42
1:A:222:LYS:HB3	1:A:232:PRO:HG2	2.02	0.41
1:B:181:LYS:HA	1:B:181:LYS:HD2	1.88	0.41
1:A:264:GLU:HB3	1:A:269:CYS:HB3	2.01	0.41
1:B:326:ILE:HG22	1:B:330:SER:HB2	2.02	0.41
1:A:168:LEU:HD11	1:A:181:LYS:HD3	2.02	0.40
1:A:172:LEU:CD1	1:A:247:PHE:HB2	2.52	0.40
1:A:231:ILE:HD12	1:A:231:ILE:H	1.84	0.40
1:A:328:GLN:OE1	1:A:331:ARG:NH1	2.44	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	179/181 (99%)	173 (97%)	6 (3%)	0	100	100
1	B	147/181 (81%)	147 (100%)	0	0	100	100
All	All	326/362 (90%)	320 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	166/166 (100%)	165 (99%)	1 (1%)	78	73
1	B	140/166 (84%)	138 (99%)	2 (1%)	59	49
All	All	306/332 (92%)	303 (99%)	3 (1%)	76	60

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	340	ILE
1	B	248[A]	LYS
1	B	248[B]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	230	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	176/181 (97%)	0.66	19 (10%)	11 10	19, 38, 70, 82	5 (2%)
1	B	148/181 (81%)	0.89	29 (19%)	3 2	18, 36, 84, 90	5 (3%)
All	All	324/362 (89%)	0.77	48 (14%)	5 5	18, 37, 79, 90	10 (3%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	236	VAL	7.2
1	B	185	ILE	6.5
1	B	233	ALA	6.2
1	A	231	ILE	6.2
1	B	235	ILE	5.3
1	B	206	ILE	4.7
1	A	340	ILE	4.6
1	B	222	LYS	4.6
1	B	237	PHE	4.1
1	B	183	MET	4.1
1	B	166	HIS	4.0
1	B	220	PHE	4.0
1	B	204	LEU	4.0
1	A	233	ALA	4.0
1	B	218	ILE	3.7
1	A	190	LEU	3.6
1	B	238	VAL	3.6
1	B	239	LEU	3.5
1	B	205	THR	3.4
1	A	232	PRO	3.4
1	A	235	ILE	3.4
1	A	229[A]	ASN	3.4
1	B	221	PRO	3.3
1	B	208	VAL	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	227	THR	3.1
1	B	340	ILE	3.0
1	B	168	LEU	2.9
1	A	165	THR	2.8
1	B	219	THR	2.7
1	B	329	THR	2.7
1	A	220	PHE	2.6
1	B	182	LYS	2.6
1	B	209[A]	LYS	2.6
1	A	221	PRO	2.6
1	B	165	THR	2.6
1	B	212	TRP	2.5
1	A	197	ILE	2.5
1	A	222	LYS	2.4
1	A	230	ASN	2.3
1	B	179	CYS	2.3
1	A	306	LYS	2.3
1	A	203	ILE	2.2
1	A	166	HIS	2.2
1	B	181	LYS	2.2
1	B	234	ASP	2.2
1	B	170	VAL	2.1
1	A	236	VAL	2.1
1	A	219	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.