



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:32 AM UTC

PDB ID : 3AHE / pdb_00003ahe
Title : Phosphoketolase from Bifidobacterium Breve complexed with dihydroxyethyl thiamine diphosphate
Authors : Suzuki, R.; Katayama, T.; Kim, B.-J.; Wakagi, T.; Shoun, H.; Ashida, H.; Yamamoto, K.; Fushinobu, S.
Deposited on : 2010-04-22
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	 81% 12% • 5%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7280 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	802	Total	C	N	O	S	0	0	0
			6379	4053	1085	1217	24			


There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP D6PAH1
A	-18	GLY	-	expression tag	UNP D6PAH1
A	-17	SER	-	expression tag	UNP D6PAH1
A	-16	SER	-	expression tag	UNP D6PAH1
A	-15	HIS	-	expression tag	UNP D6PAH1
A	-14	HIS	-	expression tag	UNP D6PAH1
A	-13	HIS	-	expression tag	UNP D6PAH1
A	-12	HIS	-	expression tag	UNP D6PAH1
A	-11	HIS	-	expression tag	UNP D6PAH1
A	-10	HIS	-	expression tag	UNP D6PAH1
A	-9	SER	-	expression tag	UNP D6PAH1
A	-8	SER	-	expression tag	UNP D6PAH1
A	-7	GLY	-	expression tag	UNP D6PAH1
A	-6	LEU	-	expression tag	UNP D6PAH1
A	-5	VAL	-	expression tag	UNP D6PAH1
A	-4	PRO	-	expression tag	UNP D6PAH1
A	-3	ARG	-	expression tag	UNP D6PAH1
A	-2	GLY	-	expression tag	UNP D6PAH1
A	-1	SER	-	expression tag	UNP D6PAH1
A	0	HIS	-	expression tag	UNP D6PAH1

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is 2-[3-[(4-AMINO-2-METHYL-5-PYRIMIDINYL)METHYL]-2-(1,2-DIHYDROXYETHYL)-4-METHYL-1,3-THIAZOL-3-IUM-5-YL]ETHYL TRIHYDROGEN DIPHOSPHATE (CCD ID: THD) (formula: $C_{14}H_{22}N_4O_9P_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			30	14	4	9	2	1		

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is NONAETHYLENE GLYCOL (CCD ID: 2PE) (formula: $C_{18}H_{38}O_{10}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			28	18	10		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	797	Total	O	0	0
			797	797		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	173.53Å 173.53Å 163.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.07 – 2.10 49.07 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.07-2.10) 99.7 (49.07-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.75 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.161 , 0.206 (Not available) , 0.175	Depositor DCC
R_{free} test set	3637 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7280	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO, THD, NA, 2PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.57	41/6553 (0.6%)	1.24	18/8909 (0.2%)

All (41) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	617	ALA	CA-CB	8.05	1.66	1.53
1	A	749	VAL	CA-CB	7.49	1.63	1.54
1	A	382	ILE	CA-CB	7.41	1.64	1.54
1	A	185	ALA	CA-CB	7.02	1.65	1.53
1	A	164	ALA	N-CA	6.79	1.54	1.46
1	A	575	ILE	C-O	6.79	1.31	1.24
1	A	5	VAL	N-CA	6.64	1.58	1.46
1	A	701	ALA	CA-CB	6.55	1.64	1.53
1	A	697	GLU	C-O	6.46	1.31	1.24
1	A	515	HIS	CA-C	6.40	1.61	1.52
1	A	696	THR	CA-CB	-5.96	1.44	1.53
1	A	119	ILE	CA-C	5.95	1.58	1.53
1	A	640	VAL	CA-CB	5.88	1.60	1.54
1	A	219	ILE	CA-CB	5.79	1.61	1.54
1	A	779	ALA	CA-CB	5.76	1.62	1.53
1	A	208	VAL	N-CA	5.75	1.53	1.46
1	A	230	ASP	CG-OD1	5.73	1.36	1.25
1	A	50	GLU	CA-C	5.67	1.59	1.53
1	A	195	GLN	CG-CD	5.59	1.66	1.52
1	A	159	TYR	CA-CB	5.59	1.62	1.53
1	A	692	ASP	N-CA	5.57	1.53	1.46
1	A	414	ALA	CA-CB	5.56	1.62	1.53
1	A	572	VAL	N-CA	5.49	1.53	1.46
1	A	133	PHE	C-O	5.49	1.30	1.24
1	A	89	THR	CA-C	5.48	1.59	1.52
1	A	735	LYS	CA-CB	5.46	1.60	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	298	THR	CA-C	5.43	1.57	1.52
1	A	786	GLN	CA-C	-5.43	1.45	1.52
1	A	746	MET	CA-C	5.42	1.59	1.52
1	A	754	ARG	CD-NE	-5.34	1.38	1.46
1	A	626	ALA	C-O	5.33	1.30	1.23
1	A	753	ASP	CA-C	5.27	1.60	1.53
1	A	555	ASP	C-N	5.25	1.37	1.33
1	A	146	GLU	CA-C	5.19	1.60	1.52
1	A	533	SER	C-O	5.18	1.30	1.24
1	A	176	VAL	CA-CB	5.17	1.60	1.54
1	A	134	SER	CB-OG	-5.12	1.31	1.42
1	A	204	THR	N-CA	5.09	1.52	1.46
1	A	760	ALA	C-O	-5.03	1.18	1.24
1	A	558	VAL	CA-C	5.03	1.59	1.52
1	A	757	LEU	CA-C	5.02	1.59	1.52

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	549	ASN	N-CA-C	7.28	123.06	114.04
1	A	382	ILE	CB-CA-C	-6.04	101.39	111.29
1	A	561	LEU	N-CA-CB	-5.83	101.62	110.07
1	A	68	THR	CA-C-N	-5.81	113.70	119.56
1	A	68	THR	C-N-CA	-5.81	113.70	119.56
1	A	508	ILE	N-CA-C	5.55	117.54	111.77
1	A	550	GLY	N-CA-C	5.54	122.28	112.53
1	A	153	GLU	N-CA-C	5.38	117.23	111.36
1	A	456	ASN	CA-C-N	-5.34	115.27	120.34
1	A	456	ASN	C-N-CA	-5.34	115.27	120.34
1	A	436	ASP	N-CA-C	-5.29	106.12	112.58
1	A	126	LEU	N-CA-C	-5.29	105.19	111.69
1	A	752	MET	N-CA-C	-5.17	107.00	113.72
1	A	132	GLN	N-CA-C	5.16	118.83	112.54
1	A	388	LEU	CA-C-N	-5.12	114.96	120.03
1	A	388	LEU	C-N-CA	-5.12	114.96	120.03
1	A	759	ALA	N-CA-C	5.10	117.22	111.11
1	A	120	THR	N-CA-C	-5.02	103.31	110.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6379	0	6085	43	0
2	A	1	0	0	0	0
3	A	30	0	19	1	0
4	A	1	0	0	0	0
5	A	44	0	66	1	0
6	A	28	0	38	0	0
7	A	797	0	0	14	0
All	All	7280	0	6208	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:735:LYS:HD3	7:A:1498:HOH:O	1.92	0.70
1:A:43:ARG:NH2	1:A:135:TYR:OH	2.29	0.66
1:A:331:GLU:HB2	7:A:1508:HOH:O	1.99	0.63
1:A:664:LYS:NZ	7:A:1344:HOH:O	2.32	0.62
1:A:286:MET:HE3	7:A:1515:HOH:O	2.00	0.61
1:A:170:ASN:HD21	1:A:431:ARG:HH11	1.47	0.61
1:A:27:LYS:HE2	7:A:950:HOH:O	2.05	0.54
1:A:435:PRO:HA	1:A:476:GLN:O	2.07	0.54
1:A:635:GLU:OE1	1:A:635:GLU:HA	2.08	0.54
1:A:637:ASN:HB3	5:A:829:EDO:H22	1.90	0.54
1:A:60:ARG:NH1	7:A:1336:HOH:O	2.37	0.54
1:A:368:GLU:OE1	7:A:1307:HOH:O	2.18	0.53
1:A:594:SER:HB2	1:A:597:LYS:HD2	1.92	0.51
1:A:703:LYS:NZ	7:A:1605:HOH:O	2.37	0.51
1:A:273:GLU:OE1	7:A:1291:HOH:O	2.19	0.50
1:A:96:GLY:HA3	1:A:153:GLU:O	2.12	0.50
1:A:432:ILE:CD1	1:A:446:THR:HG21	2.42	0.50
1:A:78:ILE:O	1:A:82:ILE:HG13	2.13	0.49
1:A:756:ALA:HB2	1:A:781:ARG:CZ	2.43	0.49
1:A:775:ASP:HA	7:A:1136:HOH:O	2.12	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:144:ALA:HB1	1:A:145:PRO:HD2	1.95	0.48
1:A:142:HIS:CD2	1:A:155:GLY:HA2	2.49	0.48
1:A:217:TYR:HB2	1:A:300:LYS:HD3	1.96	0.47
1:A:65:TRP:HE1	1:A:304:CYS:HB3	1.79	0.47
1:A:65:TRP:NE1	1:A:304:CYS:HB3	2.31	0.46
1:A:527:PRO:HG2	7:A:1386:HOH:O	2.15	0.46
1:A:43:ARG:HD3	7:A:1487:HOH:O	2.15	0.46
1:A:646:SER:HB3	1:A:655:LEU:HD22	1.97	0.46
1:A:618:ARG:HD3	7:A:1233:HOH:O	2.16	0.45
3:A:827:THD:C2	3:A:827:THD:HA4'2	2.29	0.45
1:A:326:SER:HB2	1:A:333:HIS:ND1	2.31	0.45
1:A:583:MET:HA	1:A:617:ALA:HB1	2.00	0.44
1:A:748:ARG:HA	1:A:753:ASP:OD2	2.18	0.43
1:A:657:ALA:HA	1:A:801:TRP:CZ2	2.54	0.43
1:A:554:GLN:HE22	1:A:738:GLY:HA3	1.84	0.43
1:A:635:GLU:HB3	1:A:636:ASN:ND2	2.34	0.42
1:A:602:PHE:CD2	1:A:602:PHE:N	2.88	0.42
1:A:28:TYR:CD2	1:A:76:ALA:HB2	2.55	0.41
1:A:60:ARG:HH11	1:A:60:ARG:HD3	1.64	0.41
1:A:170:ASN:ND2	1:A:431:ARG:HH11	2.17	0.41
1:A:635:GLU:HB2	1:A:639:GLU:OE2	2.20	0.41
1:A:435:PRO:O	1:A:436:ASP:HB3	2.22	0.40
1:A:28:TYR:CD2	1:A:28:TYR:C	2.99	0.40
1:A:85:HIS:HB3	7:A:857:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	800/845 (95%)	767 (96%)	32 (4%)	1 (0%)	48 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	677/712 (95%)	665 (98%)	12 (2%)	51 60

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	A	134	SER
1	A	147	THR
1	A	267	PHE
1	A	387	LYS
1	A	391	LEU
1	A	560	SER
1	A	561	LEU
1	A	605	LYS
1	A	635	GLU
1	A	684	GLU
1	A	754	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	79	ASN
1	A	105	GLN
1	A	170	ASN
1	A	408	GLN
1	A	441	ASN
1	A	472	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	549	ASN
1	A	554	GLN
1	A	574	ASN
1	A	673	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 15 ligands modelled in this entry, 2 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	2PE	A	840	-	27,27,27	1.34	1 (3%)	26,26,26	1.34	3 (11%)
5	EDO	A	830	-	3,3,3	0.52	0	2,2,2	0.41	0
5	EDO	A	833	-	3,3,3	0.54	0	2,2,2	0.95	0
5	EDO	A	831	-	3,3,3	0.54	0	2,2,2	0.41	0
5	EDO	A	834	-	3,3,3	0.36	0	2,2,2	0.98	0
5	EDO	A	837	-	3,3,3	0.10	0	2,2,2	1.27	0
5	EDO	A	836	-	3,3,3	0.70	0	2,2,2	0.25	0
5	EDO	A	839	-	3,3,3	0.36	0	2,2,2	0.73	0
5	EDO	A	838	-	3,3,3	0.71	0	2,2,2	0.67	0
5	EDO	A	832	-	3,3,3	0.45	0	2,2,2	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	835	-	3,3,3	0.48	0	2,2,2	0.91	0
3	THD	A	827	2	29,31,31	2.30	9 (31%)	36,46,46	2.34	12 (33%)
5	EDO	A	829	-	3,3,3	0.60	0	2,2,2	0.20	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	2PE	A	840	-	-	13/25/25/25	-
5	EDO	A	830	-	-	1/1/1/1	-
5	EDO	A	833	-	-	0/1/1/1	-
5	EDO	A	831	-	-	1/1/1/1	-
5	EDO	A	834	-	-	1/1/1/1	-
5	EDO	A	837	-	-	1/1/1/1	-
5	EDO	A	836	-	-	0/1/1/1	-
5	EDO	A	839	-	-	1/1/1/1	-
5	EDO	A	838	-	-	0/1/1/1	-
5	EDO	A	832	-	-	0/1/1/1	-
5	EDO	A	835	-	-	0/1/1/1	-
3	THD	A	827	2	-	4/18/23/23	0/2/2/2
5	EDO	A	829	-	-	1/1/1/1	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	827	THD	C2-S1	6.04	1.86	1.74
3	A	827	THD	C7'-C5'	5.65	1.61	1.51
3	A	827	THD	C4'-N3'	4.15	1.40	1.35
3	A	827	THD	C5-C4	3.58	1.42	1.35
3	A	827	THD	O9-C8	3.14	1.41	1.32
3	A	827	THD	C9-C8	2.79	1.57	1.49
3	A	827	THD	CM4-C4	2.70	1.53	1.49
3	A	827	THD	C5-S1	-2.68	1.68	1.74
3	A	827	THD	C7'-N3	2.42	1.51	1.47
6	A	840	2PE	O7-C6	2.22	1.51	1.42

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	827	THD	C5-S1-C2	5.32	93.87	90.80
3	A	827	THD	C7'-N3-C4	-5.31	118.24	124.98
3	A	827	THD	C7'-C5'-C4'	4.86	127.49	122.56
3	A	827	THD	C6'-N1'-C2'	4.78	123.93	116.07
3	A	827	THD	N1'-C2'-N3'	-3.41	119.85	125.53
6	A	840	2PE	C17-O16-C15	3.27	127.59	113.26
3	A	827	THD	C6-C5-C4	3.12	135.85	128.20
3	A	827	THD	CM2-C2'-N1'	2.96	120.35	117.20
3	A	827	THD	O10-C9-C8	2.72	115.53	110.36
6	A	840	2PE	O4-C5-C6	2.67	122.52	110.35
3	A	827	THD	C7'-N3-C2	2.63	129.39	126.13
3	A	827	THD	O1A-PA-O3A	-2.07	101.67	107.27
3	A	827	THD	C5'-C7'-N3	-2.04	110.67	113.14
3	A	827	THD	C4-C5-S1	-2.04	108.78	110.63
6	A	840	2PE	O10-C9-C8	2.02	119.58	110.35

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	827	THD	O9-C8-C9-O10
6	A	840	2PE	O19-C20-C21-O22
6	A	840	2PE	O4-C5-C6-O7
6	A	840	2PE	O22-C23-C24-O25
6	A	840	2PE	O25-C26-C27-O28
5	A	831	EDO	O1-C1-C2-O2
5	A	834	EDO	O1-C1-C2-O2
5	A	837	EDO	O1-C1-C2-O2
6	A	840	2PE	O7-C8-C9-O10
6	A	840	2PE	O13-C14-C15-O16
6	A	840	2PE	O1-C2-C3-O4
6	A	840	2PE	O10-C11-C12-O13
6	A	840	2PE	C6-C5-O4-C3
3	A	827	THD	PA-O3A-PB-O1B
6	A	840	2PE	C14-C15-O16-C17
6	A	840	2PE	C24-C23-O22-C21
6	A	840	2PE	C17-C18-O19-C20
5	A	839	EDO	O1-C1-C2-O2
6	A	840	2PE	C2-C3-O4-C5
3	A	827	THD	PA-O3A-PB-O2B
3	A	827	THD	PA-O3A-PB-O3B
5	A	830	EDO	O1-C1-C2-O2
5	A	829	EDO	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	827	THD	1	0
5	A	829	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	802/845 (94%)	-0.52	0 100 100	11, 19, 33, 50	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	2PE	A	840	28/28	0.66	0.25	36,50,59,61	0
5	EDO	A	836	4/4	0.77	0.24	51,52,56,57	0
5	EDO	A	830	4/4	0.84	0.14	40,40,43,45	0
5	EDO	A	831	4/4	0.85	0.18	51,53,53,54	0
5	EDO	A	829	4/4	0.88	0.12	33,36,39,42	0
5	EDO	A	837	4/4	0.89	0.13	38,39,40,45	0
5	EDO	A	832	4/4	0.91	0.14	47,48,48,51	0
5	EDO	A	833	4/4	0.91	0.11	32,33,33,36	0
5	EDO	A	834	4/4	0.91	0.11	34,34,37,44	0

Continued on next page...




Continued from previous page...


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	A	835	4/4	0.92	0.13	36,38,38,40	0
5	EDO	A	839	4/4	0.96	0.07	26,29,29,30	0
4	NA	A	828	1/1	0.96	0.10	18,18,18,18	1
5	EDO	A	838	4/4	0.98	0.07	17,17,18,18	0
2	MG	A	826	1/1	0.98	0.03	15,15,15,15	0
3	THD	A	827	30/30	0.98	0.06	12,17,25,31	4

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2PE A 840:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.