



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:48 PM EDT

PDB ID : 3AHF / pdb_00003ahf
Title : Phosphoketolase from Bifidobacterium Breve complexed with inorganic phosphate
Authors : Suzuki, R.; Katayama, T.; Kim, B.-J.; Wakagi, T.; Shoun, H.; Ashida, H.; Yamamoto, K.; Fushinobu, S.
Deposited on : 2010-04-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	829	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6947 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	802	Total	C	N	O	S	0	0	0
			6379	4053	1085	1217	24			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP D6PAH1
A	-18	GLY	-	expression tag	UNP D6PAH1
A	-17	SER	-	expression tag	UNP D6PAH1
A	-16	SER	-	expression tag	UNP D6PAH1
A	-15	HIS	-	expression tag	UNP D6PAH1
A	-14	HIS	-	expression tag	UNP D6PAH1
A	-13	HIS	-	expression tag	UNP D6PAH1
A	-12	HIS	-	expression tag	UNP D6PAH1
A	-11	HIS	-	expression tag	UNP D6PAH1
A	-10	HIS	-	expression tag	UNP D6PAH1
A	-9	SER	-	expression tag	UNP D6PAH1
A	-8	SER	-	expression tag	UNP D6PAH1
A	-7	GLY	-	expression tag	UNP D6PAH1
A	-6	LEU	-	expression tag	UNP D6PAH1
A	-5	VAL	-	expression tag	UNP D6PAH1
A	-4	PRO	-	expression tag	UNP D6PAH1
A	-3	ARG	-	expression tag	UNP D6PAH1
A	-2	GLY	-	expression tag	UNP D6PAH1
A	-1	SER	-	expression tag	UNP D6PAH1
A	0	HIS	-	expression tag	UNP D6PAH1

- Molecule 2 is THIAMINE DIPHOSPHATE (CCD ID: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	518	Total	O	0	0
			518	518		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	173.91Å 173.91Å 163.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.88 – 2.30 40.88 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (40.88-2.30) 99.7 (40.88-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.14 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.177 , 0.226 0.175 , 0.225	Depositor DCC
R_{free} test set	2814 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	26.6	Xtriage
Anisotropy	0.088	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6947	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPP, PO4, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.31	16/6553 (0.2%)	1.22	22/8909 (0.2%)

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	749	VAL	CA-CB	10.00	1.66	1.54
1	A	176	VAL	CA-CB	9.40	1.62	1.54
1	A	180	ILE	CA-CB	-7.10	1.46	1.54
1	A	382	ILE	CA-CB	6.48	1.62	1.54
1	A	488	ALA	CA-CB	6.28	1.63	1.53
1	A	185	ALA	CA-CB	6.08	1.64	1.53
1	A	219	ILE	CA-CB	6.05	1.61	1.54
1	A	298	THR	CA-C	5.58	1.57	1.52
1	A	701	ALA	CA-CB	5.55	1.62	1.53
1	A	803	TYR	N-CA	5.39	1.51	1.45
1	A	640	VAL	CA-CB	5.38	1.60	1.54
1	A	586	ALA	CA-CB	5.22	1.61	1.53
1	A	15	ARG	CA-C	5.17	1.57	1.52
1	A	565	LYS	CG-CD	-5.11	1.37	1.52
1	A	280	ALA	CA-CB	5.11	1.61	1.53
1	A	654	GLU	CA-C	5.05	1.59	1.52

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	374	ASN	CA-C-N	9.04	128.69	119.56
1	A	374	ASN	C-N-CA	9.04	128.69	119.56
1	A	543	VAL	CB-CA-C	-9.03	96.48	111.29
1	A	262	ARG	NE-CZ-NH2	-6.58	113.28	119.20
1	A	642	VAL	CB-CA-C	-6.36	99.70	110.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	176	VAL	N-CA-C	5.86	113.95	108.15
1	A	442	ARG	N-CA-C	5.78	119.66	111.92
1	A	219	ILE	N-CA-C	5.77	115.89	110.30
1	A	36	SER	N-CA-C	-5.66	104.73	111.69
1	A	650	VAL	N-CA-CB	5.65	119.12	111.21
1	A	201	ASN	CA-C-N	-5.40	114.21	120.04
1	A	201	ASN	C-N-CA	-5.40	114.21	120.04
1	A	493	GLY	N-CA-C	5.37	123.27	115.32
1	A	741	THR	N-CA-C	5.35	115.60	108.38
1	A	22	ILE	CB-CA-C	-5.27	105.03	112.14
1	A	423	ILE	N-CA-C	-5.21	105.63	110.53
1	A	267	PHE	N-CA-C	-5.17	105.82	111.82
1	A	396	VAL	CB-CA-C	5.15	117.11	111.08
1	A	786	GLN	N-CA-C	-5.12	105.70	111.28
1	A	467	MET	CG-SD-CE	-5.10	89.69	100.90
1	A	492	THR	N-CA-C	-5.04	106.97	112.72
1	A	262	ARG	CG-CD-NE	-5.03	100.94	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6379	0	6085	75	0
2	A	26	0	16	1	0
3	A	1	0	0	0	0
4	A	5	0	0	0	0
5	A	18	0	24	8	0
6	A	518	0	0	7	0
All	All	6947	0	6125	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:ASN:HB2	6:A:1045:HOH:O	1.41	1.20
1:A:131:ARG:HG3	1:A:131:ARG:HH11	1.06	1.09
1:A:220:ALA:HB1	5:A:829:GOL:H12	1.45	0.96
1:A:656:MET:HE1	1:A:671:VAL:HG11	1.51	0.90
1:A:131:ARG:HG3	1:A:131:ARG:NH1	1.85	0.87
1:A:220:ALA:HB1	5:A:829:GOL:C1	2.08	0.84
1:A:330:THR:O	1:A:331:GLU:CB	2.26	0.82
1:A:423:ILE:HD11	1:A:432:ILE:CD1	2.13	0.78
1:A:754:ARG:HH11	1:A:754:ARG:HG2	1.46	0.78
1:A:795:ILE:HD13	1:A:796:PRO:HD2	1.66	0.78
1:A:423:ILE:HD11	1:A:432:ILE:HD11	1.68	0.75
1:A:220:ALA:CB	5:A:829:GOL:H12	2.17	0.75
1:A:330:THR:O	1:A:331:GLU:HB3	1.87	0.75
2:A:826:TPP:C2	2:A:826:TPP:HN42	2.00	0.74
1:A:220:ALA:HB3	5:A:829:GOL:H32	1.71	0.73
1:A:254:GLU:OE2	1:A:262:ARG:HD3	1.89	0.72
1:A:656:MET:HE1	1:A:671:VAL:CG1	2.20	0.71
1:A:805:ASP:O	1:A:806:VAL:HB	1.90	0.69
1:A:43:ARG:HB2	1:A:131:ARG:HH12	1.59	0.68
1:A:646:SER:HB2	6:A:892:HOH:O	1.93	0.67
1:A:146:GLU:HG2	6:A:881:HOH:O	1.94	0.66
1:A:516:ALA:HB1	1:A:565:LYS:HG2	1.76	0.66
1:A:366:LYS:HD2	6:A:985:HOH:O	1.98	0.64
1:A:620:GLU:OE2	1:A:675:VAL:HG23	1.98	0.63
1:A:220:ALA:HB3	5:A:829:GOL:C3	2.27	0.63
1:A:516:ALA:CB	1:A:565:LYS:HG2	2.30	0.62
1:A:402:TYR:O	1:A:806:VAL:HA	2.00	0.61
1:A:423:ILE:HD11	1:A:432:ILE:HD12	1.84	0.60
1:A:220:ALA:CB	5:A:829:GOL:H32	2.33	0.59
1:A:170:ASN:HD21	1:A:431:ARG:HH11	1.48	0.59
1:A:131:ARG:HH11	1:A:131:ARG:CG	1.92	0.59
1:A:306:LYS:HE3	1:A:314:GLU:OE1	2.03	0.59
1:A:805:ASP:O	1:A:806:VAL:CB	2.51	0.59
1:A:571:HIS:O	1:A:597:LYS:HE3	2.03	0.57
1:A:778:ASN:HD21	5:A:831:GOL:H32	1.70	0.57
1:A:703:LYS:HE3	6:A:1004:HOH:O	2.06	0.55
1:A:583:MET:SD	1:A:675:VAL:HG21	2.45	0.55
1:A:48:MET:HE3	1:A:52:PHE:CE2	2.42	0.54
1:A:435:PRO:HA	1:A:476:GLN:O	2.08	0.54
1:A:590:LYS:HE3	6:A:1221:HOH:O	2.09	0.52
1:A:806:VAL:HG13	1:A:806:VAL:O	2.09	0.52
1:A:579:THR:HG21	1:A:628:TRP:HH2	1.76	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:VAL:HG13	1:A:610:THR:HG21	1.92	0.51
1:A:754:ARG:HD2	1:A:755:TYR:CZ	2.47	0.50
1:A:635:GLU:HB2	1:A:639:GLU:OE2	2.13	0.49
1:A:118:ASN:HD21	1:A:128:LYS:HZ2	1.60	0.49
1:A:654:GLU:OE1	1:A:754:ARG:NH1	2.46	0.48
1:A:163:HIS:HA	1:A:488:ALA:HB2	1.96	0.48
1:A:118:ASN:HD21	1:A:128:LYS:NZ	2.12	0.47
1:A:121:LYS:HE2	6:A:1217:HOH:O	2.15	0.47
1:A:214:LEU:HD22	1:A:226:ALA:HB2	1.97	0.47
1:A:131:ARG:NH1	1:A:131:ARG:CG	2.61	0.47
1:A:144:ALA:HB1	1:A:145:PRO:HD2	1.96	0.47
1:A:642:VAL:HG22	1:A:765:ILE:HD11	1.95	0.47
1:A:254:GLU:OE2	1:A:262:ARG:CD	2.61	0.46
1:A:142:HIS:O	1:A:144:ALA:N	2.45	0.46
1:A:300:LYS:NZ	1:A:320:HIS:ND1	2.64	0.46
1:A:735:LYS:NZ	1:A:750:ASN:O	2.49	0.45
1:A:795:ILE:HD13	1:A:796:PRO:CD	2.42	0.45
1:A:486:LEU:O	1:A:490:LEU:HG	2.17	0.44
1:A:230:ASP:OD1	1:A:246:GLU:OE1	2.35	0.44
1:A:646:SER:HB3	1:A:652:THR:OG1	2.17	0.43
1:A:330:THR:O	1:A:331:GLU:HB2	2.12	0.43
1:A:433:PHE:HA	1:A:474:THR:O	2.18	0.43
1:A:61:LEU:HB3	1:A:322:VAL:HG11	2.00	0.43
1:A:326:SER:HB2	1:A:333:HIS:ND1	2.34	0.43
1:A:579:THR:HG21	1:A:628:TRP:CH2	2.54	0.43
1:A:583:MET:HA	1:A:617:ALA:HB1	2.00	0.42
1:A:451:ASP:HB3	1:A:467:MET:HA	2.02	0.42
1:A:630:TRP:CD1	1:A:630:TRP:H	2.38	0.42
1:A:754:ARG:HH11	1:A:754:ARG:CG	2.22	0.42
1:A:245:TYR:OH	1:A:273:GLU:OE2	2.37	0.42
1:A:783:LYS:HE2	1:A:783:LYS:HB2	1.90	0.42
1:A:142:HIS:CD2	1:A:155:GLY:HA2	2.55	0.41
1:A:440:SER:O	1:A:605:LYS:HD3	2.21	0.41
1:A:220:ALA:CB	5:A:829:GOL:C3	2.96	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	800/845 (95%)	754 (94%)	43 (5%)	3 (0%)	30	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	331	GLU
1	A	687	ASP
1	A	543	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	677/712 (95%)	653 (96%)	24 (4%)	32	48

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	VAL
1	A	43	ARG
1	A	131	ARG
1	A	147	THR
1	A	199	LEU
1	A	214	LEU
1	A	300	LYS
1	A	381	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	396	VAL
1	A	397	THR
1	A	477	LEU
1	A	543	VAL
1	A	560	SER
1	A	605	LYS
1	A	642	VAL
1	A	646	SER
1	A	650	VAL
1	A	668	LYS
1	A	684	GLU
1	A	693	GLU
1	A	754	ARG
1	A	795	ILE
1	A	802	VAL
1	A	806	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	105	GLN
1	A	118	ASN
1	A	170	ASN
1	A	321	GLN
1	A	441	ASN
1	A	472	GLN
1	A	554	GLN
1	A	574	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	PO4	A	828	-	4,4,4	0.88	0	6,6,6	0.83	0
2	TPP	A	826	3	26,27,27	2.36	7 (26%)	38,40,40	2.53	11 (28%)
5	GOL	A	830	-	5,5,5	1.30	0	5,5,5	2.08	3 (60%)
5	GOL	A	829	-	5,5,5	1.13	0	5,5,5	1.33	1 (20%)
5	GOL	A	831	-	5,5,5	0.35	0	5,5,5	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	831	-	-	4/4/4/4	-
2	TPP	A	826	3	-	4/17/17/17	0/2/2/2
5	GOL	A	830	-	-	2/4/4/4	-
5	GOL	A	829	-	-	3/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	826	TPP	C2-N3	6.79	1.49	1.32
2	A	826	TPP	C2-S1	6.44	1.88	1.69
2	A	826	TPP	PA-O3A	-4.20	1.55	1.59
2	A	826	TPP	C7'-C5'	2.53	1.55	1.51
2	A	826	TPP	C4'-N3'	2.43	1.38	1.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	826	TPP	C5'-C4'	-2.18	1.39	1.42
2	A	826	TPP	C5-C4	2.13	1.39	1.35

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	826	TPP	S1-C2-N3	-8.34	101.73	112.30
2	A	826	TPP	C2-S1-C5	7.73	96.34	91.22
2	A	826	TPP	C6'-N1'-C2'	4.06	122.73	116.07
2	A	826	TPP	N1'-C2'-N3'	-4.02	118.84	125.53
2	A	826	TPP	C5-C4-N3	3.38	117.83	111.67
2	A	826	TPP	CM4-C4-C5	-3.04	119.97	127.75
2	A	826	TPP	CM2-C2'-N1'	2.92	120.31	117.20
5	A	830	GOL	O2-C2-C1	2.88	121.09	109.18
5	A	830	GOL	O2-C2-C3	2.55	119.72	109.18
2	A	826	TPP	C7'-C5'-C4'	2.52	126.96	122.36
2	A	826	TPP	CM2-C2'-N3'	2.46	120.81	117.13
2	A	826	TPP	O3A-PA-O1A	-2.32	103.72	110.70
2	A	826	TPP	C5'-C7'-N3	-2.25	108.11	112.97
5	A	829	GOL	O3-C3-C2	2.06	119.66	110.38
5	A	830	GOL	O1-C1-C2	2.03	119.51	110.38

There are no chirality outliers.

All (13) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	826	TPP	PA-O3A-PB-O3B
5	A	829	GOL	O1-C1-C2-C3
5	A	830	GOL	C1-C2-C3-O3
5	A	831	GOL	O1-C1-C2-C3
5	A	831	GOL	C1-C2-C3-O3
5	A	830	GOL	O1-C1-C2-O2
5	A	829	GOL	O1-C1-C2-O2
5	A	831	GOL	O1-C1-C2-O2
2	A	826	TPP	C4'-C5'-C7'-N3
2	A	826	TPP	C6'-C5'-C7'-N3
5	A	831	GOL	O2-C2-C3-O3
5	A	829	GOL	O2-C2-C3-O3
2	A	826	TPP	PA-O3A-PB-O2B

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	826	TPP	1	0
5	A	829	GOL	7	0
5	A	831	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	802/845 (94%)	-0.34	5 (0%) 85 86	15, 26, 42, 60	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	806	VAL	5.4
1	A	5	VAL	2.9
1	A	330	THR	2.5
1	A	795	ILE	2.2
1	A	638	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	A	830	6/6	0.78	0.19	29,32,34,35	0
4	PO4	A	828	5/5	0.83	0.24	83,85,86,87	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	GOL	A	831	6/6	0.84	0.21	60,62,64,66	0
5	GOL	A	829	6/6	0.90	0.22	40,44,45,46	0
2	TPP	A	826	26/26	0.98	0.05	12,21,24,29	0
3	MG	A	827	1/1	0.99	0.02	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.