



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:43 PM UTC

PDB ID : 3AHJ / pdb_00003ahj
Title : H553A mutant of Phosphoketolase from Bifidobacterium Breve
Authors : Suzuki, R.; Katayama, T.; Kim, B.-J.; Wakagi, T.; Shoun, H.; Ashida, H.; Yamamoto, K.; Fushinobu, S.
Deposited on : 2010-04-22
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	845	 83% 11% • 5%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	802	Total	C	N	O	S	0	0	0
			6374	4050	1083	1217	24			


There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP D6PAH1
A	-18	GLY	-	expression tag	UNP D6PAH1
A	-17	SER	-	expression tag	UNP D6PAH1
A	-16	SER	-	expression tag	UNP D6PAH1
A	-15	HIS	-	expression tag	UNP D6PAH1
A	-14	HIS	-	expression tag	UNP D6PAH1
A	-13	HIS	-	expression tag	UNP D6PAH1
A	-12	HIS	-	expression tag	UNP D6PAH1
A	-11	HIS	-	expression tag	UNP D6PAH1
A	-10	HIS	-	expression tag	UNP D6PAH1
A	-9	SER	-	expression tag	UNP D6PAH1
A	-8	SER	-	expression tag	UNP D6PAH1
A	-7	GLY	-	expression tag	UNP D6PAH1
A	-6	LEU	-	expression tag	UNP D6PAH1
A	-5	VAL	-	expression tag	UNP D6PAH1
A	-4	PRO	-	expression tag	UNP D6PAH1
A	-3	ARG	-	expression tag	UNP D6PAH1
A	-2	GLY	-	expression tag	UNP D6PAH1
A	-1	SER	-	expression tag	UNP D6PAH1
A	0	HIS	-	expression tag	UNP D6PAH1
A	553	ALA	HIS	engineered mutation	UNP D6PAH1

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is THIAMINE DIPHOSPHATE (CCD ID: TPP) (formula: $C_{12}H_{19}N_4O_7P_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	542	Total O 542 542	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Xylulose 5-phosphate/fructose 6-phosphate phosphoketolase

Chain A:  83% 11% • 5%



4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	175.08Å 175.08Å 164.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.61 – 2.10 38.61 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (38.61-2.10) 99.8 (38.61-2.10)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.63 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.158 , 0.191 (Not available) , 0.180	Depositor DCC
R_{free} test set	3722 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.127	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 29.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6987	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TPP, MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.58	44/6547 (0.7%)	1.21	24/8901 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (44) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	779	ALA	CA-CB	8.58	1.67	1.53
1	A	417	ALA	CA-CB	7.58	1.65	1.53
1	A	617	ALA	CA-CB	7.58	1.65	1.53
1	A	732	VAL	CA-C	6.60	1.60	1.52
1	A	180	ILE	CA-CB	-6.57	1.46	1.54
1	A	219	ILE	CA-CB	6.55	1.62	1.54
1	A	787	PHE	N-CA	-6.45	1.38	1.46
1	A	640	VAL	CA-CB	6.43	1.61	1.53
1	A	257	MET	N-CA	6.41	1.54	1.46
1	A	5	VAL	N-CA	6.34	1.58	1.46
1	A	516	ALA	CA-CB	6.23	1.63	1.53
1	A	780	PHE	CA-C	6.21	1.61	1.52
1	A	749	VAL	CA-CB	6.19	1.62	1.54
1	A	488	ALA	CA-CB	6.17	1.63	1.53
1	A	230	ASP	CG-OD1	6.11	1.36	1.25
1	A	279	ALA	CA-CB	6.08	1.62	1.53
1	A	672	VAL	CA-C	6.07	1.60	1.52
1	A	697	GLU	C-O	5.98	1.31	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	124	ALA	CA-CB	5.87	1.62	1.53
1	A	183	GLY	N-CA	5.76	1.52	1.45
1	A	291	TYR	C-O	-5.66	1.16	1.24
1	A	298	THR	C-O	-5.66	1.19	1.24
1	A	181	GLY	N-CA	5.63	1.52	1.45
1	A	185	ALA	CA-CB	5.59	1.63	1.53
1	A	757	LEU	CA-C	5.51	1.59	1.52
1	A	14	ASP	CA-C	5.50	1.60	1.53
1	A	759	ALA	C-O	5.49	1.30	1.24
1	A	140	PRO	CA-C	5.49	1.59	1.52
1	A	618	ARG	N-CA	5.36	1.52	1.46
1	A	559	THR	N-CA	5.33	1.52	1.46
1	A	516	ALA	N-CA	5.32	1.52	1.46
1	A	675	VAL	CA-CB	-5.32	1.49	1.55
1	A	61	LEU	N-CA	5.28	1.52	1.46
1	A	740	THR	CA-CB	5.28	1.59	1.53
1	A	175	PHE	C-O	-5.25	1.17	1.24
1	A	248	VAL	CA-CB	5.23	1.61	1.54
1	A	280	ALA	CA-CB	5.22	1.61	1.53
1	A	92	ILE	C-O	5.21	1.29	1.24
1	A	322	VAL	CA-CB	5.21	1.59	1.54
1	A	667	ILE	CA-CB	5.20	1.60	1.54
1	A	418	TYR	CA-C	5.14	1.59	1.52
1	A	642	VAL	CA-CB	5.13	1.61	1.54
1	A	45	ASN	CA-C	5.08	1.59	1.52
1	A	701	ALA	CA-CB	5.02	1.61	1.53

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	759	ALA	N-CA-C	6.32	117.83	111.07
1	A	218	LYS	N-CA-C	-6.27	102.73	108.75
1	A	176	VAL	CA-C-N	-6.17	113.42	119.90
1	A	176	VAL	C-N-CA	-6.17	113.42	119.90
1	A	328	ARG	NE-CZ-NH2	-6.16	113.66	119.20
1	A	176	VAL	N-CA-C	5.95	114.04	108.15
1	A	382	ILE	CB-CA-C	-5.95	103.26	111.65
1	A	561	LEU	CB-CA-C	-5.95	101.54	110.88
1	A	170	ASN	CA-C-N	-5.94	115.97	123.15
1	A	170	ASN	C-N-CA	-5.94	115.97	123.15
1	A	126	LEU	N-CA-C	-5.80	105.04	111.36
1	A	288	ARG	N-CA-C	-5.60	102.59	109.65

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	THR	N-CA-C	5.55	120.47	113.25
1	A	328	ARG	CG-CD-NE	-5.45	100.02	112.00
1	A	150	SER	N-CA-C	5.36	118.28	109.76
1	A	371	ILE	N-CA-C	-5.22	105.44	110.72
1	A	550	GLY	CA-C-N	5.22	127.27	120.28
1	A	550	GLY	C-N-CA	5.22	127.27	120.28
1	A	292	PRO	CB-CA-C	-5.20	104.58	111.23
1	A	167	ALA	N-CA-C	5.16	116.91	111.28
1	A	480	HIS	N-CA-C	5.08	116.82	111.28
1	A	49	LYS	CB-CA-C	5.06	118.08	109.53
1	A	567	PHE	N-CA-C	5.06	118.27	110.42
1	A	328	ARG	NE-CZ-NH1	5.03	126.53	121.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	597	LYS	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6374	0	6083	30	0
2	A	1	0	0	0	0
3	A	26	0	16	3	0
4	A	44	0	66	3	0
5	A	542	0	0	3	0
All	All	6987	0	6165	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:827:TPP:HN42	3:A:827:TPP:C2	1.93	0.82
1:A:635:GLU:OE1	1:A:635:GLU:HA	2.02	0.60
1:A:432:ILE:CD1	1:A:446:THR:HG21	2.33	0.58
1:A:273:GLU:OE1	5:A:1221:HOH:O	2.17	0.57
1:A:245:TYR:OH	1:A:273:GLU:OE2	2.20	0.56
1:A:328:ARG:HG2	1:A:328:ARG:O	2.05	0.55
1:A:735:LYS:HD3	5:A:1348:HOH:O	2.07	0.55
1:A:654:GLU:OE1	1:A:754:ARG:HD3	2.09	0.52
1:A:739:SER:OG	1:A:740:THR:N	2.43	0.52
1:A:797:GLU:OE1	4:A:836:EDO:O1	2.28	0.51
1:A:554:GLN:HE22	1:A:738:GLY:HA3	1.76	0.51
1:A:657:ALA:HA	1:A:801:TRP:CZ2	2.46	0.50
1:A:96:GLY:HA3	1:A:153:GLU:O	2.12	0.50
1:A:583:MET:HA	1:A:617:ALA:HB1	1.94	0.49
1:A:654:GLU:OE1	1:A:754:ARG:CD	2.60	0.49
1:A:43:ARG:HD3	5:A:1336:HOH:O	2.14	0.48
1:A:620:GLU:HG2	1:A:626:ALA:CB	2.43	0.47
1:A:131:ARG:HD2	4:A:832:EDO:O1	2.15	0.47
1:A:756:ALA:HB2	1:A:781:ARG:CZ	2.45	0.46
1:A:702:ASP:OD1	4:A:828:EDO:O1	2.33	0.46
1:A:13:LEU:HD23	1:A:13:LEU:HA	1.74	0.44
1:A:498:TRP:CE2	1:A:537:LEU:HD13	2.53	0.44
1:A:170:ASN:HD21	1:A:431:ARG:HH11	1.66	0.44
1:A:483:GLU:OE2	1:A:487:GLU:OE2	2.36	0.44
1:A:190:LEU:C	1:A:190:LEU:HD13	2.43	0.43
1:A:651:PRO:HG2	1:A:710:HIS:HA	2.00	0.43
3:A:827:TPP:HN42	3:A:827:TPP:H2	1.77	0.43
1:A:786:GLN:NE2	1:A:790:ASP:OD1	2.48	0.43
1:A:483:GLU:O	1:A:487:GLU:HG3	2.20	0.41
3:A:827:TPP:C2	3:A:827:TPP:N4'	2.73	0.41
1:A:432:ILE:HD11	1:A:446:THR:HG21	2.03	0.41
1:A:435:PRO:HA	1:A:476:GLN:O	2.20	0.40
1:A:135:TYR:CG	1:A:136:PRO:HD2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	800/845 (95%)	768 (96%)	31 (4%)	1 (0%)	48 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	302	TRP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	676/711 (95%)	670 (99%)	6 (1%)	70 78

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	ARG
1	A	147	THR
1	A	381	VAL
1	A	391	LEU
1	A	561	LEU
1	A	754	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	ASN
1	A	86	GLN
1	A	105	GLN
1	A	170	ASN
1	A	441	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	472	GLN
1	A	554	GLN
1	A	574	ASN
1	A	673	ASN
1	A	758	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	EDO	A	830	-	3,3,3	0.60	0	2,2,2	0.93	0
3	TPP	A	827	2	26,27,27	2.59	7 (26%)	38,40,40	2.58	12 (31%)
4	EDO	A	833	-	3,3,3	0.56	0	2,2,2	0.35	0
4	EDO	A	828	-	3,3,3	0.73	0	2,2,2	0.14	0
4	EDO	A	829	-	3,3,3	0.63	0	2,2,2	0.20	0
4	EDO	A	836	-	3,3,3	0.20	0	2,2,2	0.88	0
4	EDO	A	837	-	3,3,3	0.61	0	2,2,2	0.82	0
4	EDO	A	835	-	3,3,3	0.42	0	2,2,2	0.82	0
4	EDO	A	834	-	3,3,3	0.58	0	2,2,2	1.17	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	A	831	-	3,3,3	0.43	0	2,2,2	0.57	0
4	EDO	A	838	-	3,3,3	0.59	0	2,2,2	0.31	0
4	EDO	A	832	-	3,3,3	0.54	0	2,2,2	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	830	-	-	1/1/1/1	-
3	TPP	A	827	2	-	4/17/17/17	0/2/2/2
4	EDO	A	833	-	-	1/1/1/1	-
4	EDO	A	828	-	-	0/1/1/1	-
4	EDO	A	829	-	-	0/1/1/1	-
4	EDO	A	836	-	-	1/1/1/1	-
4	EDO	A	837	-	-	0/1/1/1	-
4	EDO	A	835	-	-	1/1/1/1	-
4	EDO	A	834	-	-	0/1/1/1	-
4	EDO	A	831	-	-	0/1/1/1	-
4	EDO	A	838	-	-	0/1/1/1	-
4	EDO	A	832	-	-	0/1/1/1	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	827	TPP	C2-N3	7.97	1.51	1.32
3	A	827	TPP	C2-S1	6.76	1.89	1.69
3	A	827	TPP	PA-O3A	-3.79	1.55	1.59
3	A	827	TPP	C7'-C5'	3.31	1.57	1.51
3	A	827	TPP	C4'-N3'	3.00	1.39	1.35
3	A	827	TPP	C2'-N1'	2.73	1.38	1.34
3	A	827	TPP	PA-O1A	-2.04	1.43	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	827	TPP	S1-C2-N3	-9.55	100.20	112.30
3	A	827	TPP	C2-S1-C5	6.56	95.56	91.22
3	A	827	TPP	CM2-C2'-N1'	4.30	121.78	117.20
3	A	827	TPP	CM4-C4-C5	-3.55	118.66	127.75
3	A	827	TPP	C5-C4-N3	3.33	117.73	111.67

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	827	TPP	C6'-N1'-C2'	3.22	121.36	116.07
3	A	827	TPP	C5'-C7'-N3	-2.81	106.91	112.97
3	A	827	TPP	O3A-PA-O1A	-2.66	102.69	110.70
3	A	827	TPP	C5'-C4'-N4'	-2.46	118.83	122.14
3	A	827	TPP	O2A-PA-O3A	2.39	113.73	107.27
3	A	827	TPP	O2B-PB-O3A	2.19	111.98	104.64
3	A	827	TPP	C5'-C6'-N1'	-2.18	120.28	123.83

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	827	TPP	C4'-C5'-C7'-N3
3	A	827	TPP	C6'-C5'-C7'-N3
4	A	830	EDO	O1-C1-C2-O2
4	A	836	EDO	O1-C1-C2-O2
4	A	833	EDO	O1-C1-C2-O2
4	A	835	EDO	O1-C1-C2-O2
3	A	827	TPP	PA-O3A-PB-O2B
3	A	827	TPP	PA-O3A-PB-O3B

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	827	TPP	3	0
4	A	828	EDO	1	0
4	A	836	EDO	1	0
4	A	832	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	802/845 (94%)	-0.51	3 (0%) 88 90	19, 27, 42, 59	0

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	806	VAL	2.7
1	A	230	ASP	2.6
1	A	5	VAL	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	830	4/4	0.77	0.24	64,67,68,70	0
4	EDO	A	831	4/4	0.86	0.17	58,60,61,61	0
4	EDO	A	828	4/4	0.88	0.15	48,49,52,52	0
4	EDO	A	829	4/4	0.89	0.13	54,55,55,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	EDO	A	835	4/4	0.90	0.15	54,55,55,57	0
4	EDO	A	834	4/4	0.91	0.15	43,43,47,48	0
4	EDO	A	836	4/4	0.91	0.13	42,47,49,52	0
4	EDO	A	833	4/4	0.93	0.12	40,40,42,44	0
4	EDO	A	832	4/4	0.93	0.13	51,53,54,55	0
4	EDO	A	837	4/4	0.96	0.08	27,27,27,32	0
4	EDO	A	838	4/4	0.98	0.09	35,38,40,40	0
3	TPP	A	827	26/26	0.99	0.05	20,24,32,38	0
2	MG	A	826	1/1	1.00	0.01	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.