



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 07:54 PM EDT

PDB ID : 7AHJ / pdb_00007ahj
Title : Crystal structure of PPARgamma V290M mutant ligand binding domain in complex with farglitazar
Authors : Schoenmakers, E.; Schwabe, B.T.W.; Fairall, L.; Chatterjee, K.; Schwabe, J.W.R.
Deposited on : 2020-09-24
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	
1	B	275	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8415 atoms, of which 4106 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	260	Total	C	H	N	O	S	0	1	0
			4041	1313	2009	327	381	11			
1	B	264	Total	C	H	N	O	S	0	1	0
			4145	1342	2068	338	385	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	ALA	-	expression tag	UNP P37231
A	290	MET	VAL	engineered mutation	UNP P37231
B	203	ALA	-	expression tag	UNP P37231
B	290	MET	VAL	engineered mutation	UNP P37231

- Molecule 2 is 2-(2-BENZOYL-PHENYLAMINO)-3-{4-[2-(5-METHYL-2-PHENYL-OXAZO L-4-YL)-ETHOXY]-PHENYL}-PROPIONIC ACID (CCD ID: 570) (formula: C₃₄H₃₀N₂O₅).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			70	34	29	2	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	84	Total	O	0	0
			84	84		
3	B	75	Total	O	0	0
			75	75		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 1: Peroxisome proliferator-activated receptor gamma



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.96Å 61.24Å 118.51Å 90.00° 103.17° 90.00°	Depositor
Resolution (Å)	28.92 – 2.10 28.92 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (28.92-2.10) 99.8 (28.92-2.10)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.03 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.18.2_3874, REFMAC 5.8.0238	Depositor
R, R_{free}	0.200 , 0.237 0.202 , 0.237	Depositor DCC
R_{free} test set	1903 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	29.4	Xtriage
Anisotropy	0.886	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8415	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
570

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	3/2072 (0.1%)	0.64	6/2800 (0.2%)
1	B	0.30	1/2116 (0.0%)	0.56	4/2856 (0.1%)
All	All	0.37	4/4188 (0.1%)	0.60	10/5656 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	288	ARG	CZ-NH2	-7.20	1.24	1.33
1	A	288	ARG	CG-CD	6.68	1.72	1.52
1	B	265	LYS	CE-NZ	-6.66	1.29	1.49
1	A	288	ARG	CZ-NH1	5.16	1.40	1.32

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	205	ASN	N-CA-C	-8.87	100.24	108.22
1	A	205	ASN	CB-CA-C	8.73	121.75	110.22
1	B	266	HIS	N-CA-C	-8.14	102.91	113.17
1	B	454	GLN	CA-CB-CG	-6.30	101.50	114.10
1	A	205	ASN	N-CA-CB	-6.16	104.63	111.10
1	A	427	GLU	CA-CB-CG	5.87	125.85	114.10
1	A	288	ARG	CB-CG-CD	-5.83	97.88	111.30
1	A	395	GLY	N-CA-C	5.69	120.93	114.67
1	B	266	HIS	N-CA-CB	-5.64	102.14	110.49
1	B	454	GLN	CB-CG-CD	-5.28	103.63	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2032	2009	2031	30	0
1	B	2077	2068	2082	22	0
2	A	41	29	29	6	0
3	A	84	0	0	0	0
3	B	75	0	0	0	0
All	All	4309	4106	4142	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:268:THR:HG21	1:B:469:LEU:HD11	1.37	1.04
1:B:325:ILE:HD12	1:B:388:ILE:HG23	1.45	0.94
1:A:288:ARG:HD2	1:A:288:ARG:O	1.72	0.87
1:A:286:GLN:HB3	2:A:501:570:C1C	2.14	0.78
1:A:262:ILE:HG13	1:A:263:LYS:H	1.56	0.70
1:B:358:LYS:CB	1:B:359:PRO:CD	2.74	0.66
1:A:358:LYS:HB3	1:A:359:PRO:HD3	1.79	0.63
1:A:370:PHE:HB2	1:A:445:ILE:HD11	1.81	0.62
1:A:288:ARG:HD2	1:A:288:ARG:C	2.27	0.60
1:A:286:GLN:HB3	2:A:501:570:H1C	1.82	0.60
1:A:258:GLY:O	1:A:262:ILE:HG12	2.03	0.58
1:A:282:PHE:CE1	2:A:501:570:H1D	2.39	0.58
1:B:330:LEU:HD21	1:B:364:MET:HE1	1.85	0.57
1:A:288:ARG:C	1:A:288:ARG:CD	2.79	0.56
1:B:387:PHE:CE1	1:B:439:MET:HE3	2.40	0.55
1:A:325:ILE:HD11	1:A:392:ILE:HG13	1.88	0.55
1:A:262:ILE:HG13	1:A:263:LYS:N	2.21	0.55
1:A:358:LYS:CB	1:A:359:PRO:CD	2.85	0.55
1:A:283:GLN:HA	1:A:286:GLN:HG2	1.91	0.51
1:A:370:PHE:CB	1:A:445:ILE:HD11	2.40	0.51
1:B:449:HIS:NE2	1:B:477:TYR:O	2.47	0.48
1:A:252:MET:HE1	1:A:277:VAL:HG21	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:325:ILE:HG23	1:B:388:ILE:HD12	1.96	0.47
1:B:328:THR:OG1	1:B:442:LEU:HD11	2.14	0.47
1:A:358:LYS:HB3	1:A:359:PRO:CD	2.45	0.47
1:A:359:PRO:HG2	1:A:456:ILE:HD13	1.95	0.47
1:A:443:ARG:NH1	1:B:444:GLN:OE1	2.48	0.47
1:A:290:MET:SD	1:A:466:HIS:CG	3.09	0.46
1:A:466:HIS:CG	1:A:467:PRO:HD2	2.51	0.46
1:B:266:HIS:ND1	1:B:266:HIS:C	2.74	0.46
1:A:364:MET:SD	2:A:501:570:HE1	2.56	0.46
2:A:501:570:O1G	2:A:501:570:N	2.49	0.46
1:A:226:PHE:CG	1:A:329:MET:HE1	2.51	0.45
1:B:262:ILE:O	1:B:266:HIS:HB3	2.17	0.45
1:B:387:PHE:HE1	1:B:439:MET:HE3	1.80	0.45
1:B:325:ILE:HD11	1:B:391:ILE:HB	1.99	0.44
1:B:318:LEU:O	1:B:322:VAL:HG13	2.17	0.44
1:B:357:ARG:HD2	1:B:358:LYS:CB	2.48	0.43
1:A:286:GLN:HB3	2:A:501:570:C1D	2.48	0.43
1:A:317:LEU:HD22	1:A:392:ILE:O	2.19	0.43
1:A:466:HIS:CE1	1:A:467:PRO:HD2	2.53	0.43
1:B:310:ASP:OD1	1:B:312:ASN:HB2	2.19	0.42
1:A:359:PRO:HG2	1:A:456:ILE:CD1	2.50	0.42
1:A:325:ILE:HD11	1:A:392:ILE:CG1	2.48	0.42
1:B:367:LYS:HD2	1:B:367:LYS:N	2.33	0.42
1:B:379:LEU:HD11	1:B:435:LEU:HD13	2.01	0.42
1:B:266:HIS:CE1	1:B:284:GLY:CA	3.03	0.42
1:A:384:LEU:O	1:A:388:ILE:HG23	2.20	0.41
1:A:226:PHE:CD1	1:A:329:MET:HE1	2.56	0.41
1:B:255:LEU:HD23	1:B:277:VAL:HG13	2.04	0.40
1:B:358:LYS:CB	1:B:359:PRO:HD2	2.50	0.40
1:B:338:GLY:HA3	1:B:347:PHE:CZ	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/275 (93%)	249 (98%)	5 (2%)	1 (0%)	30	28
1	B	259/275 (94%)	254 (98%)	4 (2%)	1 (0%)	30	28
All	All	514/550 (94%)	503 (98%)	9 (2%)	2 (0%)	30	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	358	LYS
1	A	358	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/247 (90%)	221 (100%)	0	100	100
1	B	224/247 (91%)	223 (100%)	1 (0%)	84	89
All	All	445/494 (90%)	444 (100%)	1 (0%)	87	93

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	266	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	345	GLN
1	A	451	GLN
1	B	323	HIS
1	B	345	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	570	A	501	-	44,45,45	1.49	7 (15%)	57,61,61	2.14	6 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	570	A	501	-	-	3/30/30/30	0/5/5/5

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	570	C3I-C3G	-3.83	1.39	1.47
2	A	501	570	C1A-N	3.79	1.45	1.37
2	A	501	570	C3G-N3H	3.59	1.35	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	570	O3F-C3D	-3.58	1.32	1.38
2	A	501	570	C1H-C1G	2.87	1.54	1.49
2	A	501	570	C3C-N3H	-2.76	1.32	1.38
2	A	501	570	O1G-C1G	-2.41	1.18	1.22

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	570	C3E-C3D-C3C	-9.84	127.17	135.22
2	A	501	570	O3F-C3D-C3E	6.81	124.79	115.88
2	A	501	570	O3F-C3G-N3H	-5.53	108.33	114.37
2	A	501	570	C1A-C1F-C1G	-5.03	118.67	121.96
2	A	501	570	O3F-C3G-C3I	3.85	125.43	118.73
2	A	501	570	O1-C-O2	-2.44	118.55	124.08

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	501	570	OH-C3A-C3B-C3C
2	A	501	570	O1-C-CA-N
2	A	501	570	O2-C-CA-N

There are no ring outliers.

1 monomer is involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	570	6	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	260/275 (94%)	0.34	20 (7%) 19 21	24, 40, 70, 91	0
1	B	264/275 (96%)	0.50	28 (10%) 11 12	18, 42, 72, 94	1 (0%)
All	All	524/550 (95%)	0.42	48 (9%) 14 15	18, 41, 70, 94	1 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	282	PHE	5.3
1	B	239	GLY	5.1
1	B	270	LEU	4.6
1	B	206	PRO	4.5
1	B	244	LYS	4.5
1	B	266	HIS	3.9
1	A	363	PHE	3.9
1	B	268	THR	3.6
1	A	239	GLY	3.4
1	B	258	GLY	3.4
1	B	275	LYS	3.3
1	B	357	ARG	3.3
1	A	463	MET	3.1
1	B	247	PHE	3.1
1	B	260	ASP	3.1
1	A	262	ILE	3.0
1	B	358	LYS	3.0
1	B	261	LYS	2.9
1	B	265	LYS	2.9
1	B	269	PRO	2.9
1	A	275	LYS	2.9
1	B	464	SER	2.9
1	A	465	LEU	2.8
1	A	279	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	203	ALA	2.7
1	B	427	GLU	2.7
1	B	262	ILE	2.7
1	A	276	GLU	2.6
1	A	252	MET	2.6
1	A	278	ALA	2.6
1	B	477	TYR	2.6
1	B	463	MET	2.6
1	A	427	GLU	2.5
1	A	358	LYS	2.5
1	A	452	LEU	2.5
1	B	259	GLU	2.4
1	A	286	GLN	2.3
1	A	362	ASP	2.3
1	B	264	PHE	2.3
1	B	267	ILE	2.3
1	B	461	THR	2.2
1	A	263	LYS	2.2
1	A	244	LYS	2.1
1	B	363	PHE	2.1
1	B	238	THR	2.1
1	B	474	LYS	2.1
1	A	464	SER	2.1
1	B	257	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	570	A	501	41/41	0.74	0.17	44,59,73,86	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.