



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:33 AM UTC

PDB ID : 3AIV / pdb_00003aiv
Title : Crystal structure of beta-glucosidase in rye complexed with an aglycone DIM-BOA
Authors : Sue, M.; Nakamura, C.; Miyamoto, T.; Yajima, S.
Deposited on : 2010-05-18
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	564	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4220 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	486	3932	2517	651	745	19	0	0	0

There are 47 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-44	MET	-	expression tag	UNP Q9FYS3
A	-43	HIS	-	expression tag	UNP Q9FYS3
A	-42	HIS	-	expression tag	UNP Q9FYS3
A	-41	HIS	-	expression tag	UNP Q9FYS3
A	-40	HIS	-	expression tag	UNP Q9FYS3
A	-39	HIS	-	expression tag	UNP Q9FYS3
A	-38	HIS	-	expression tag	UNP Q9FYS3
A	-37	SER	-	expression tag	UNP Q9FYS3
A	-36	SER	-	expression tag	UNP Q9FYS3
A	-35	GLY	-	expression tag	UNP Q9FYS3
A	-34	LEU	-	expression tag	UNP Q9FYS3
A	-33	VAL	-	expression tag	UNP Q9FYS3
A	-32	PRO	-	expression tag	UNP Q9FYS3
A	-31	ARG	-	expression tag	UNP Q9FYS3
A	-30	GLY	-	expression tag	UNP Q9FYS3
A	-29	SER	-	expression tag	UNP Q9FYS3
A	-28	GLY	-	expression tag	UNP Q9FYS3
A	-27	MET	-	expression tag	UNP Q9FYS3
A	-26	LYS	-	expression tag	UNP Q9FYS3
A	-25	GLU	-	expression tag	UNP Q9FYS3
A	-24	THR	-	expression tag	UNP Q9FYS3
A	-23	ALA	-	expression tag	UNP Q9FYS3
A	-22	ALA	-	expression tag	UNP Q9FYS3
A	-21	ALA	-	expression tag	UNP Q9FYS3
A	-20	LYS	-	expression tag	UNP Q9FYS3
A	-19	PHE	-	expression tag	UNP Q9FYS3
A	-18	GLU	-	expression tag	UNP Q9FYS3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	ARG	-	expression tag	UNP Q9FYS3
A	-16	GLN	-	expression tag	UNP Q9FYS3
A	-15	HIS	-	expression tag	UNP Q9FYS3
A	-14	MET	-	expression tag	UNP Q9FYS3
A	-13	ASP	-	expression tag	UNP Q9FYS3
A	-12	SER	-	expression tag	UNP Q9FYS3
A	-11	PRO	-	expression tag	UNP Q9FYS3
A	-10	ASP	-	expression tag	UNP Q9FYS3
A	-9	LEU	-	expression tag	UNP Q9FYS3
A	-8	GLY	-	expression tag	UNP Q9FYS3
A	-7	THR	-	expression tag	UNP Q9FYS3
A	-6	ASP	-	expression tag	UNP Q9FYS3
A	-5	ASP	-	expression tag	UNP Q9FYS3
A	-4	ASP	-	expression tag	UNP Q9FYS3
A	-3	ASP	-	expression tag	UNP Q9FYS3
A	-2	LYS	-	expression tag	UNP Q9FYS3
A	-1	ALA	-	expression tag	UNP Q9FYS3
A	0	MET	-	expression tag	UNP Q9FYS3
A	323	ALA	GLY	SEE REMARK 999	UNP Q9FYS3
A	325	SER	LEU	SEE REMARK 999	UNP Q9FYS3

- Molecule 2 is 2,4-DIHYDROXY-7-(METHYLOXY)-2H-1,4-BENZOXAZIN-3(4H)-ONE (CCD ID: HBO) (formula: C₉H₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			15	9	1	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	273	Total 273	O 273	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-glucosidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 3 2	Depositor
Cell constants a, b, c, α , β , γ	194.19Å 194.19Å 194.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.10 – 2.50 47.10 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.7 (47.10-2.50) 99.7 (47.10-2.50)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.11 (at 2.51Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.190 , 0.209 0.187 , 0.207	Depositor DCC
R_{free} test set	2194 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 35.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4220	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HBO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.76	58/4055 (1.4%)	1.19	13/5498 (0.2%)

All (58) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	456	GLY	N-CA	-10.48	1.30	1.45
1	A	147	ASP	CG-OD2	9.82	1.44	1.25
1	A	25	LYS	CD-CE	8.93	1.79	1.52
1	A	147	ASP	CA-C	8.05	1.63	1.52
1	A	251	LYS	N-CA	7.95	1.56	1.46
1	A	385	LYS	CE-NZ	7.81	1.72	1.49
1	A	346	SER	C-O	-7.46	1.14	1.24
1	A	77	VAL	CA-CB	7.37	1.64	1.54
1	A	58	TRP	CA-C	7.07	1.62	1.52
1	A	248	HIS	CA-C	6.91	1.61	1.52
1	A	186	TRP	N-CA	-6.83	1.37	1.45
1	A	147	ASP	CA-CB	6.80	1.65	1.53
1	A	458	ILE	CG1-CD1	-6.74	1.25	1.51
1	A	251	LYS	CG-CD	6.73	1.72	1.52
1	A	495	LYS	CD-CE	6.66	1.72	1.52
1	A	263	VAL	CA-CB	6.63	1.64	1.54
1	A	335	THR	CA-C	6.37	1.60	1.53
1	A	251	LYS	CA-C	6.35	1.61	1.52
1	A	251	LYS	CE-NZ	6.28	1.68	1.49
1	A	93	LYS	CD-CE	6.21	1.71	1.52
1	A	443	ILE	CG1-CD1	-6.18	1.27	1.51
1	A	248	HIS	CD2-NE2	6.15	1.44	1.37
1	A	25	LYS	CE-NZ	6.13	1.67	1.49
1	A	134	ARG	NE-CZ	6.11	1.39	1.33
1	A	44	ILE	CG1-CD1	-6.06	1.28	1.51
1	A	14	VAL	C-O	6.05	1.35	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	31	LYS	CG-CD	6.04	1.70	1.52
1	A	73	THR	CA-CB	6.01	1.62	1.53
1	A	249	TYR	CA-C	-5.84	1.45	1.53
1	A	480	ASP	N-CA	-5.78	1.38	1.46
1	A	232	HIS	CA-C	-5.71	1.45	1.52
1	A	294	VAL	CA-CB	5.64	1.60	1.54
1	A	248	HIS	CE1-NE2	5.59	1.38	1.32
1	A	259	MET	CB-CG	-5.57	1.35	1.52
1	A	434	ARG	C-O	5.57	1.31	1.24
1	A	200	TYR	CA-C	-5.48	1.45	1.52
1	A	148	THR	N-CA	5.47	1.55	1.46
1	A	125	TYR	C-O	5.42	1.30	1.24
1	A	86	GLU	N-CA	5.41	1.53	1.46
1	A	191	GLU	N-CA	5.38	1.50	1.46
1	A	162	LYS	CD-CE	5.35	1.68	1.52
1	A	248	HIS	C-O	5.32	1.31	1.24
1	A	321	LYS	CA-C	5.32	1.59	1.52
1	A	402	PRO	CA-C	5.29	1.58	1.52
1	A	93	LYS	CE-NZ	5.23	1.65	1.49
1	A	352	LEU	CA-C	-5.22	1.46	1.52
1	A	343	ASP	CG-OD1	5.22	1.35	1.25
1	A	460	ASN	C-O	5.20	1.29	1.23
1	A	62	CYS	N-CA	5.16	1.52	1.46
1	A	458	ILE	C-O	-5.16	1.18	1.24
1	A	483	LYS	CA-C	5.12	1.59	1.52
1	A	451	GLY	C-O	5.09	1.29	1.24
1	A	16	THR	CA-CB	-5.07	1.45	1.53
1	A	233	HIS	N-CA	5.07	1.52	1.46
1	A	122	ILE	C-O	5.06	1.29	1.24
1	A	259	MET	CG-SD	5.06	1.93	1.80
1	A	65	TYR	N-CA	5.04	1.52	1.46
1	A	411	ALA	CA-CB	-5.02	1.44	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	436	ILE	N-CA-C	-8.36	102.09	110.62
1	A	112	ASN	N-CA-C	6.38	120.07	112.93
1	A	473	LEU	N-CA-C	-6.10	104.55	112.23
1	A	147	ASP	N-CA-CB	-6.07	100.23	110.49
1	A	472	GLY	N-CA-C	6.01	121.11	112.82
1	A	45	GLU	N-CA-C	5.92	117.41	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	393	ILE	CB-CA-C	-5.75	104.37	112.14
1	A	412	ASP	N-CA-C	-5.55	100.19	109.24
1	A	248	HIS	CB-CA-C	-5.40	101.37	110.44
1	A	202	GLU	N-CA-C	-5.38	105.44	112.23
1	A	141	VAL	N-CA-C	5.32	115.52	107.80
1	A	27	ASP	CB-CA-C	-5.14	100.49	110.46
1	A	265	GLY	N-CA-C	-5.10	106.02	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3932	0	3693	87	0
2	A	15	0	9	0	0
3	A	273	0	0	11	0
All	All	4220	0	3702	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:25:LYS:CD	1:A:25:LYS:CE	1.79	1.59
1:A:251:LYS:CE	1:A:251:LYS:NZ	1.68	1.55
1:A:385:LYS:NZ	1:A:385:LYS:CE	1.72	1.52
1:A:25:LYS:CE	1:A:25:LYS:NZ	1.67	1.52
1:A:385:LYS:HD3	3:A:520:HOH:O	1.43	1.16
1:A:126:ASN:ND2	1:A:182:ARG:HH22	1.58	0.99
1:A:74:ASN:HD22	1:A:76:ASP:H	1.16	0.94
1:A:126:ASN:HD22	1:A:182:ARG:HH22	1.12	0.88
1:A:450:ARG:HG2	1:A:450:ARG:HH11	1.45	0.79
1:A:60:HIS:O	1:A:64:THR:HG23	1.83	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:ASN:ND2	1:A:76:ASP:H	1.81	0.77
1:A:78:ALA:HA	1:A:467:TYR:OH	1.87	0.74
1:A:432:LEU:O	1:A:436:ILE:HG12	1.94	0.68
1:A:74:ASN:HD22	1:A:76:ASP:N	1.89	0.68
1:A:181:ASP:OD2	1:A:182:ARG:HD3	1.95	0.66
1:A:57:THR:HG22	1:A:153:GLU:HG3	1.77	0.66
1:A:232:HIS:HD2	1:A:299:TYR:OH	1.79	0.65
1:A:87:GLU:HG3	3:A:661:HOH:O	1.97	0.64
1:A:251:LYS:NZ	1:A:251:LYS:CD	2.61	0.64
1:A:349:THR:HG22	3:A:790:HOH:O	1.98	0.64
1:A:281:ARG:HH22	1:A:341:HIS:HD2	1.46	0.64
1:A:232:HIS:CD2	1:A:312:PRO:HB2	2.33	0.63
1:A:198:PHE:CG	1:A:204:ILE:HD11	2.33	0.63
1:A:188:THR:HG23	1:A:257:ILE:HD11	1.82	0.62
1:A:35:PHE:HE1	1:A:436:ILE:HD12	1.67	0.60
1:A:186:TRP:HB2	1:A:257:ILE:HG12	1.84	0.60
1:A:364:THR:HG22	1:A:370:GLU:HG2	1.83	0.59
1:A:43:GLN:HG2	1:A:461:PHE:O	2.03	0.58
1:A:341:HIS:HE1	1:A:343:ASP:OD1	1.88	0.57
1:A:126:ASN:HD22	1:A:182:ARG:NH2	1.93	0.55
1:A:198:PHE:CD1	1:A:204:ILE:HD11	2.41	0.55
1:A:488:LYS:HE2	3:A:707:HOH:O	2.07	0.54
1:A:74:ASN:ND2	1:A:76:ASP:CG	2.66	0.54
1:A:232:HIS:CD2	1:A:299:TYR:OH	2.60	0.54
1:A:109:ILE:HG13	1:A:122:ILE:HD12	1.90	0.53
1:A:99:VAL:HG11	1:A:453:PHE:CZ	2.44	0.53
1:A:159:PHE:O	1:A:233:HIS:HD2	1.91	0.53
1:A:456:GLY:HA3	1:A:471:PHE:O	2.08	0.53
1:A:397:LYS:HE2	1:A:398:TYR:CZ	2.44	0.52
1:A:266:TYR:O	1:A:279:ARG:HD2	2.09	0.52
1:A:204:ILE:H	1:A:204:ILE:HD13	1.75	0.51
1:A:43:GLN:O	1:A:460:ASN:HB2	2.11	0.51
1:A:188:THR:CG2	1:A:257:ILE:HD11	2.40	0.51
1:A:458:ILE:HG22	1:A:474:VAL:HB	1.93	0.51
1:A:188:THR:HG23	1:A:257:ILE:CD1	2.41	0.50
1:A:313:MET:HE3	3:A:730:HOH:O	2.12	0.50
1:A:93:LYS:HE3	3:A:736:HOH:O	2.13	0.49
1:A:112:ASN:C	1:A:114:THR:H	2.21	0.49
1:A:245:PHE:HZ	1:A:255:SER:HB3	1.76	0.49
1:A:457:LEU:HG	1:A:458:ILE:HG23	1.95	0.49
1:A:44:ILE:HG12	1:A:81:SER:HB2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:450:ARG:HG2	1:A:450:ARG:NH1	2.20	0.48
1:A:281:ARG:HH22	1:A:341:HIS:CD2	2.29	0.48
1:A:271:ASP:HA	3:A:711:HOH:O	2.13	0.48
1:A:25:LYS:NZ	1:A:27:ASP:OD2	2.47	0.48
1:A:51:ASP:HB3	1:A:120:LYS:HG2	1.95	0.47
1:A:394:MET:HE2	1:A:403:ILE:HD11	1.96	0.47
1:A:126:ASN:ND2	1:A:182:ARG:NH2	2.43	0.47
1:A:91:ALA:HB1	1:A:474:VAL:HG21	1.98	0.46
1:A:134:ARG:HH11	1:A:134:ARG:HG3	1.81	0.46
1:A:232:HIS:HE1	3:A:602:HOH:O	1.99	0.46
1:A:251:LYS:HA	1:A:251:LYS:HD3	1.77	0.46
1:A:481:GLY:O	1:A:482:ASN:HB2	2.14	0.46
1:A:78:ALA:O	1:A:478:LYS:NZ	2.49	0.46
1:A:118:ASN:OD1	1:A:118:ASN:C	2.58	0.46
1:A:461:PHE:CE1	1:A:466:GLY:HA2	2.50	0.45
1:A:393:ILE:HD11	3:A:595:HOH:O	2.16	0.45
1:A:289:TRP:HD1	1:A:289:TRP:O	2.00	0.45
1:A:290:PHE:C	1:A:293:PRO:HD2	2.42	0.45
1:A:400:ASN:N	1:A:401:PRO:CD	2.80	0.45
1:A:19:LYS:HD3	1:A:19:LYS:HA	1.72	0.44
1:A:144:TRP:CZ2	1:A:146:TRP:HA	2.52	0.44
1:A:263:VAL:HG21	1:A:331:LEU:HG	2.00	0.43
1:A:44:ILE:HG22	1:A:460:ASN:HB3	2.01	0.43
1:A:395:LYS:HE3	1:A:446:GLY:HA3	2.01	0.43
1:A:462:GLU:O	1:A:463:TRP:C	2.60	0.43
1:A:77:VAL:HG13	3:A:608:HOH:O	2.18	0.42
1:A:115:GLY:H	1:A:170:TYR:HE1	1.67	0.42
1:A:204:ILE:HD13	1:A:204:ILE:N	2.35	0.42
1:A:289:TRP:C	1:A:289:TRP:CD1	2.98	0.42
1:A:98:LYS:HE2	3:A:771:HOH:O	2.19	0.41
1:A:218:VAL:O	1:A:218:VAL:HG22	2.20	0.41
1:A:232:HIS:CG	1:A:312:PRO:HB2	2.55	0.41
1:A:332:ASN:CG	1:A:407:GLU:HB2	2.46	0.41
1:A:416:ASP:HA	1:A:417:PRO:HD2	1.92	0.40
1:A:204:ILE:HG12	1:A:205:HIS:CD2	2.57	0.40
1:A:458:ILE:H	1:A:458:ILE:HD13	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/564 (86%)	466 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	418/483 (86%)	399 (96%)	19 (4%)	24	49

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	ILE
1	A	64	THR
1	A	77	VAL
1	A	87	GLU
1	A	109	ILE
1	A	204	ILE
1	A	212	PRO
1	A	218	VAL
1	A	257	ILE
1	A	272	SER
1	A	331	LEU
1	A	385	LYS
1	A	419	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	443	ILE
1	A	450	ARG
1	A	458	ILE
1	A	476	ILE
1	A	482	ASN
1	A	495	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	119	GLN
1	A	126	ASN
1	A	127	ASN
1	A	163	GLN
1	A	166	ASN
1	A	232	HIS
1	A	233	HIS
1	A	286	ASN
1	A	341	HIS
1	A	369	ASN
1	A	460	ASN
1	A	482	ASN
1	A	497	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HBO	A	900	-	15,16,16	5.83	12 (80%)	16,23,23	2.05	4 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HBO	A	900	-	-	0/2/18/18	0/2/2/2

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	900	HBO	C4B-N3B	10.17	1.52	1.40
2	A	900	HBO	O1B-C2B	8.86	1.60	1.44
2	A	900	HBO	C2B-C3B	7.48	1.62	1.51
2	A	900	HBO	O7B-C9B	7.21	1.63	1.42
2	A	900	HBO	C4B-C1B	7.12	1.51	1.40
2	A	900	HBO	OHB-N3B	-6.58	1.31	1.39
2	A	900	HBO	C5B-C4B	5.89	1.48	1.39
2	A	900	HBO	O1A-C2B	5.81	1.52	1.38
2	A	900	HBO	C6B-C7B	4.42	1.47	1.38
2	A	900	HBO	C6B-C5B	3.96	1.45	1.38
2	A	900	HBO	O1B-C1B	3.78	1.43	1.38
2	A	900	HBO	O3B-C3B	2.21	1.26	1.22

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	900	HBO	C9B-O7B-C7B	-5.69	105.31	117.50
2	A	900	HBO	OHB-N3B-C3B	3.68	119.97	116.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	900	HBO	O1B-C1B-C8B	2.41	120.40	116.29
2	A	900	HBO	C6B-C7B-C8B	2.30	123.60	120.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	486/564 (86%)	-0.57	13 (2%) 56 51	12, 22, 40, 59	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	14	VAL	5.9
1	A	147	ASP	4.7
1	A	499	VAL	4.2
1	A	134	ARG	3.3
1	A	162	LYS	3.1
1	A	251	LYS	3.0
1	A	31	LYS	2.7
1	A	246	LYS	2.7
1	A	418	GLU	2.6
1	A	319	GLN	2.2
1	A	252	HIS	2.1
1	A	347	ASP	2.1
1	A	495	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HBO	A	900	15/15	0.80	0.18	36,41,46,47	0

6.5 Other polymers

There are no such residues in this entry.