



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:53 AM UTC

PDB ID : 4AI7 / pdb_00004ai7
Title : Crystal structure of Laccase from *Thermus thermophilus* HB27 complexed with Hg, crystal of the apoenzyme soaked for 2 h in 5 mM HgCl₂ at 278 K.
Authors : Serrano-Posada, H.; Rudino-Pinera, E.
Deposited on : 2012-02-08
Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	439	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MPD	A	1469	-	X	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MPD	A	1470	-	X	-	-
3	MPD	A	1471	-	-	X	-
3	MPD	A	1473	-	-	X	-
3	MPD	A	1474	-	X	X	-
3	MPD	A	1476	-	-	X	-
3	MPD	A	1477	-	-	X	-
3	MPD	A	1478	-	X	-	-
3	MPD	A	1479	-	X	-	-
3	MPD	A	1481	-	X	-	-
3	MPD	A	1482	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE-LIKE PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	439	3799	2444	686	652	17	0	41	0


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	53	ILE	LEU	SEE REMARK 999	UNP Q4H436

- Molecule 2 is MERCURY (II) ION (CCD ID: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	Hg	0	0
			5	5		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0
3	A	1	Total C O 8 6 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	504	Total O 504 504	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LACCASE-LIKE PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	93.28Å 110.10Å 96.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.57 – 1.70 28.57 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (28.57-1.70) 97.4 (28.57-1.70)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.92 (at 1.70Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.152 , 0.170 0.147 , 0.166	Depositor DCC
R_{free} test set	2695 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	14.1	Xtriage
Anisotropy	0.113	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 59.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4436	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	28/3897 (0.7%)	0.77	0/5302

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	135	HIS	CD2-NE2	-10.05	1.26	1.37
1	A	97[A]	HIS	CG-ND1	-8.60	1.28	1.38
1	A	97[B]	HIS	CG-ND1	-8.60	1.28	1.38
1	A	393	HIS	CG-ND1	-7.99	1.29	1.38
1	A	396	HIS	CD2-NE2	-7.20	1.29	1.37

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	301	MET	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3799	0	3893	119	0
2	A	5	0	0	0	0
3	A	128	0	220	101	0
4	A	504	0	0	38	3
All	All	4436	0	4113	179	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ARG:CZ	3:A:1471:MPD:C5	1.79	1.51
1:A:391[B]:MET:CE	1:A:391[B]:MET:SD	2.03	1.46
3:A:1482:MPD:CM	3:A:1482:MPD:H52	1.53	1.38
3:A:1477:MPD:HM1	4:A:2455:HOH:O	1.25	1.29
3:A:1479:MPD:HM2	3:A:1479:MPD:C5	1.33	1.29

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:2283:HOH:O	4:A:2489:HOH:O[8_545]	1.94	0.26
4:A:2010:HOH:O	4:A:2246:HOH:O[8_545]	2.14	0.06
4:A:2015:HOH:O	4:A:2405:HOH:O[6_545]	2.15	0.05

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	479/439 (109%)	468 (98%)	8 (2%)	3 (1%)	21	9

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	399[A]	VAL
1	A	399[B]	VAL
1	A	301	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	409/367 (111%)	403 (98%)	6 (2%)	57	43

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	358[A]	ARG
1	A	358[B]	ARG
1	A	412	PHE
1	A	77	ARG
1	A	75	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 5 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	A	1480	-	7,7,7	0.96	1 (14%)	9,10,10	0.50	0
3	MPD	A	1467	-	7,7,7	0.59	0	9,10,10	0.47	0
3	MPD	A	1473	-	7,7,7	1.47	1 (14%)	9,10,10	1.94	3 (33%)
3	MPD	A	1478	-	7,7,7	1.16	1 (14%)	9,10,10	3.77	6 (66%)
3	MPD	A	1482	-	7,7,7	0.67	0	9,10,10	1.17	1 (11%)
3	MPD	A	1481	-	7,7,7	0.98	1 (14%)	9,10,10	4.06	6 (66%)
3	MPD	A	1472	-	7,7,7	0.67	0	9,10,10	0.60	0
3	MPD	A	1468	-	7,7,7	1.13	1 (14%)	9,10,10	1.43	3 (33%)
3	MPD	A	1471	-	7,7,7	1.53	1 (14%)	9,10,10	1.06	1 (11%)
3	MPD	A	1470	-	7,7,7	1.57	1 (14%)	9,10,10	3.66	6 (66%)
3	MPD	A	1469	-	7,7,7	0.72	0	9,10,10	3.90	6 (66%)
3	MPD	A	1474	-	7,7,7	0.73	0	9,10,10	3.86	6 (66%)
3	MPD	A	1479	-	7,7,7	1.18	1 (14%)	9,10,10	4.32	6 (66%)
3	MPD	A	1477	-	7,7,7	0.91	0	9,10,10	4.10	6 (66%)
3	MPD	A	1476	-	7,7,7	0.89	0	9,10,10	3.97	6 (66%)
3	MPD	A	1475	-	7,7,7	0.69	0	9,10,10	3.95	6 (66%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	1480	-	-	0/5/5/5	-
3	MPD	A	1467	-	-	1/5/5/5	-
3	MPD	A	1473	-	-	1/5/5/5	-
3	MPD	A	1478	-	-	2/5/5/5	-
3	MPD	A	1482	-	-	2/5/5/5	-
3	MPD	A	1481	-	-	3/5/5/5	-
3	MPD	A	1472	-	-	2/5/5/5	-
3	MPD	A	1468	-	-	1/5/5/5	-
3	MPD	A	1471	-	-	1/5/5/5	-
3	MPD	A	1470	-	-	3/5/5/5	-
3	MPD	A	1469	-	-	3/5/5/5	-
3	MPD	A	1474	-	-	3/5/5/5	-
3	MPD	A	1479	-	-	2/5/5/5	-
3	MPD	A	1477	-	-	2/5/5/5	-
3	MPD	A	1476	-	-	1/5/5/5	-
3	MPD	A	1475	-	-	2/5/5/5	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1471	MPD	O2-C2	-3.74	1.35	1.44
3	A	1470	MPD	O2-C2	-3.74	1.35	1.44
3	A	1478	MPD	O2-C2	-2.88	1.37	1.44
3	A	1473	MPD	O2-C2	-2.74	1.38	1.44
3	A	1479	MPD	O2-C2	-2.56	1.38	1.44

The worst 5 of 62 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1479	MPD	O2-C2-C3	-7.53	81.10	109.27
3	A	1477	MPD	O2-C2-C3	-7.10	82.70	109.27
3	A	1469	MPD	C1-C2-C3	-6.75	81.06	110.20
3	A	1481	MPD	CM-C2-C3	-6.10	83.88	110.20
3	A	1474	MPD	C1-C2-C3	-6.08	83.97	110.20

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1474	MPD	C2-C3-C4-C5
3	A	1475	MPD	C2-C3-C4-C5
3	A	1477	MPD	C2-C3-C4-C5
3	A	1479	MPD	C2-C3-C4-C5
3	A	1467	MPD	O2-C2-C3-C4

There are no ring outliers.

12 monomers are involved in 101 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1480	MPD	2	0
3	A	1473	MPD	15	0
3	A	1482	MPD	6	0
3	A	1481	MPD	1	0
3	A	1472	MPD	2	0
3	A	1471	MPD	29	0
3	A	1470	MPD	1	0
3	A	1474	MPD	10	0
3	A	1479	MPD	5	0
3	A	1477	MPD	19	0
3	A	1476	MPD	7	0
3	A	1475	MPD	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	439/439 (100%)	-0.51	7 (1%) 70 75	3, 13, 32, 80	41 (9%)

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	303	HIS	3.0
1	A	302	ALA	2.9
1	A	301	MET	2.9
1	A	300	GLY	2.7
1	A	338	THR	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	A	1474	8/8	0.69	0.24	31,43,56,62	8
3	MPD	A	1481	8/8	0.74	0.18	36,54,65,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MPD	A	1482	8/8	0.76	0.17	34,41,44,46	8
3	MPD	A	1475	8/8	0.77	0.21	20,36,43,46	8
3	MPD	A	1473	8/8	0.81	0.19	23,28,40,43	8
3	MPD	A	1476	8/8	0.81	0.14	21,30,33,40	8
2	HG	A	1483	1/1	0.83	0.15	58,58,58,58	1
3	MPD	A	1469	8/8	0.85	0.13	20,25,30,31	8
3	MPD	A	1471	8/8	0.85	0.16	31,45,68,71	0
2	HG	A	1466	1/1	0.86	0.14	33,33,33,33	1
3	MPD	A	1480	8/8	0.88	0.14	25,36,44,44	8
3	MPD	A	1468	8/8	0.88	0.14	28,38,39,40	8
3	MPD	A	1479	8/8	0.88	0.13	19,35,50,64	8
3	MPD	A	1477	8/8	0.89	0.14	23,33,35,50	8
3	MPD	A	1470	8/8	0.90	0.09	18,21,28,28	0
3	MPD	A	1472	8/8	0.91	0.10	17,29,49,52	8
3	MPD	A	1478	8/8	0.93	0.09	20,22,33,41	8
2	HG	A	1464	1/1	0.95	0.11	13,13,13,13	1
3	MPD	A	1467	8/8	0.97	0.05	9,12,14,19	0
2	HG	A	1463	1/1	0.97	0.07	17,17,17,17	1
2	HG	A	1465	1/1	0.98	0.09	14,14,14,14	1

6.5 Other polymers [i](#)

There are no such residues in this entry.