



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2026 – 05:19 PM UTC

PDB ID : 6AI8 / pdb_00006ai8
Title : Cab2 mutant-H337A
Authors : Zheng, P.; Zhu, Z.
Deposited on : 2018-08-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	371	76% 6% 17% 2% 75% 7% 17%
1	B	371	76% 6% 17% 2% 75% 7% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Phosphopantothenate--cysteine ligase CAB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	309	Total	C	N	O	S	0	0	0
			2520	1623	424	467	6			
1	B	307	Total	C	N	O	S	0	1	0
			2515	1623	423	463	6			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP P40506
A	-4	HIS	-	expression tag	UNP P40506
A	-3	HIS	-	expression tag	UNP P40506
A	-2	HIS	-	expression tag	UNP P40506
A	-1	HIS	-	expression tag	UNP P40506
A	0	HIS	-	expression tag	UNP P40506
A	337	ALA	HIS	engineered mutation	UNP P40506
B	-5	HIS	-	expression tag	UNP P40506
B	-4	HIS	-	expression tag	UNP P40506
B	-3	HIS	-	expression tag	UNP P40506
B	-2	HIS	-	expression tag	UNP P40506
B	-1	HIS	-	expression tag	UNP P40506
B	0	HIS	-	expression tag	UNP P40506
B	337	ALA	HIS	engineered mutation	UNP P40506

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	103	Total	O	0	0
			103	103		
4	B	89	Total	O	0	0
			89	89		

- Molecule 1: Phosphopantothenate--cysteine ligase CAB2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	53.79Å 113.23Å 57.78Å 90.00° 93.33° 90.00°	Depositor
Resolution (Å)	57.68 – 2.30 57.68 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (57.68-2.30) 99.6 (57.68-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.89 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.184 , 0.247 0.190 , 0.246	Depositor DCC
R_{free} test set	1417 reflections (4.59%)	wwPDB-VP
Wilson B-factor (Å ²)	54.8	Xtriage
Anisotropy	0.052	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 39.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5279	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.62	0/2576	0.84	1/3485 (0.0%)
1	B	0.62	0/2572	0.83	1/3479 (0.0%)
All	All	0.62	0/5148	0.84	2/6964 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	132	ARG	N-CA-C	5.88	117.69	111.28
1	B	132	ARG	N-CA-C	5.10	118.60	112.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2520	0	2537	14	0
1	B	2515	0	2532	16	0
2	A	24	0	32	2	0
2	B	18	0	24	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	103	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	89	0	0	0	0
All	All	5279	0	5125	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:248:THR:HG22	1:B:249:PRO:HD2	1.52	0.92
1:B:332:LEU:HA	1:B:342:MET:HE1	1.81	0.62
1:A:215:PHE:CE2	1:B:253:LEU:HD12	2.38	0.58
1:B:227:ILE:HG21	1:B:245:THR:HG21	1.87	0.56
1:B:264:LEU:HD23	1:B:300:TYR:CD2	2.45	0.52
1:B:87:LEU:HD21	1:B:95:ILE:CD1	2.40	0.52
1:A:271:TRP:HE1	2:A:404:GOL:H2	1.76	0.51
1:A:215:PHE:CD2	1:B:253:LEU:HD12	2.45	0.51
1:A:146:ILE:HG21	1:B:152:ILE:HD11	1.93	0.50
1:A:297:LEU:CD1	1:A:305:VAL:HG23	2.43	0.49
1:B:187:GLN:HE21	2:B:401:GOL:H2	1.79	0.48
1:B:102:THR:HG23	1:B:127:LEU:CD1	2.45	0.47
1:B:245:THR:OG1	1:B:253:LEU:HD21	2.16	0.46
1:A:217:VAL:HG12	1:A:257:LEU:HD23	1.99	0.45
1:A:253:LEU:HD12	1:B:257:LEU:HD12	1.98	0.45
1:A:108:ALA:HB3	1:A:129:PRO:HG3	1.99	0.45
1:B:311:GLN:OE1	1:B:311:GLN:N	2.50	0.45
1:A:63:PHE:O	1:A:67:GLN:HG2	2.18	0.43
1:A:57:ILE:HG12	1:A:194:SER:HB3	2.01	0.42
1:A:262:LYS:HB3	1:A:300:TYR:HE1	1.85	0.42
1:B:337:ALA:N	1:B:341:GLU:OE1	2.48	0.42
1:B:303:GLN:OE1	1:B:303:GLN:N	2.52	0.42
1:A:187:GLN:HE21	2:A:401:GOL:C3	2.34	0.40
1:A:79:THR:HA	1:A:121:LEU:O	2.21	0.40
1:A:146:ILE:HD13	1:B:152:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/371 (82%)	298 (98%)	7 (2%)	0	100	100
1	B	304/371 (82%)	294 (97%)	10 (3%)	0	100	100
All	All	609/742 (82%)	592 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	281/336 (84%)	273 (97%)	8 (3%)	38	56
1	B	280/336 (83%)	269 (96%)	11 (4%)	28	43
All	All	561/672 (84%)	542 (97%)	19 (3%)	32	49

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	GLU
1	A	46	THR
1	A	162	LYS
1	A	214	ASP
1	A	217	VAL
1	A	253	LEU
1	A	291	TYR
1	A	317	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	41	GLU
1	B	58	LYS
1	B	102	THR
1	B	152	ILE
1	B	248	THR
1	B	250	ASP
1	B	255	VAL
1	B	262	LYS
1	B	291	TYR
1	B	353	LYS
1	B	359	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	A	163	ASN
1	A	186	ASN
1	A	308	ASN
1	B	110	GLN
1	B	163	ASN
1	B	225	HIS
1	B	324	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	403	-	5,5,5	0.37	0	5,5,5	0.57	0
2	GOL	B	401	-	5,5,5	0.51	0	5,5,5	0.85	0
2	GOL	A	403	-	5,5,5	0.50	0	5,5,5	0.43	0
2	GOL	A	404	-	5,5,5	0.43	0	5,5,5	0.41	0
2	GOL	B	402	-	5,5,5	0.25	0	5,5,5	0.46	0
3	SO4	A	405	-	4,4,4	0.55	0	6,6,6	0.47	0
2	GOL	A	402	-	5,5,5	0.46	0	5,5,5	0.17	0
3	SO4	B	404	-	4,4,4	0.46	0	6,6,6	0.19	0
2	GOL	A	401	-	5,5,5	0.57	0	5,5,5	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	403	-	-	0/4/4/4	-
2	GOL	B	401	-	-	1/4/4/4	-
2	GOL	A	403	-	-	0/4/4/4	-
2	GOL	A	404	-	-	1/4/4/4	-
2	GOL	B	402	-	-	0/4/4/4	-
2	GOL	A	402	-	-	2/4/4/4	-
2	GOL	A	401	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GOL	O1-C1-C2-C3
2	A	401	GOL	C1-C2-C3-O3
2	A	402	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	402	GOL	O1-C1-C2-C3
2	A	401	GOL	O2-C2-C3-O3
2	A	401	GOL	O1-C1-C2-O2
2	A	404	GOL	O1-C1-C2-C3
2	B	401	GOL	O2-C2-C3-O3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	GOL	1	0
2	A	404	GOL	1	0
2	A	401	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	309/371 (83%)	-0.07	2 (0%) 85 86	36, 59, 102, 140	0
1	B	307/371 (82%)	0.08	9 (2%) 53 56	32, 59, 109, 133	1 (0%)
All	All	616/742 (83%)	0.00	11 (1%) 67 69	32, 59, 108, 140	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263[A]	PHE	3.4
1	B	309	LEU	3.1
1	B	244	THR	3.0
1	A	263	PHE	2.8
1	B	227	ILE	2.7
1	B	360	THR	2.3
1	B	41	GLU	2.2
1	B	329	TRP	2.2
1	B	282	LEU	2.1
1	A	243	GLY	2.1
1	B	290	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	401	6/6	0.77	0.17	65,66,68,70	0
2	GOL	B	401	6/6	0.79	0.15	55,62,65,67	0
2	GOL	A	404	6/6	0.84	0.13	72,75,80,88	0
2	GOL	A	403	6/6	0.84	0.14	66,72,74,78	0
3	SO4	A	405	5/5	0.84	0.11	65,73,74,79	0
2	GOL	B	402	6/6	0.85	0.13	65,68,71,76	0
2	GOL	B	403	6/6	0.87	0.14	78,84,85,87	0
2	GOL	A	402	6/6	0.89	0.10	68,74,75,79	0
3	SO4	B	404	5/5	0.92	0.07	74,76,79,83	0

6.5 Other polymers [i](#)

There are no such residues in this entry.