



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:20 AM UTC

PDB ID : 4AJK / pdb_00004ajk
Title : rat LDHA in complex with N-(2-(methylamino)-1,3-benzothiazol-6-yl) acetamide
Authors : Tucker, J.A.; Brassington, C.; Hassall, G.; Vogtherr, M.; Ward, R.; Tart, J.; Davies, G.; Greenwood, R.
Deposited on : 2012-02-16
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	91% 8%
1	B	331	92% 8%
1	C	331	92% 7%
1	D	331	91% 9%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 11384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-LACTATE DEHYDROGENASE A CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	1	0
			2494	1598	420	463	13			
1	B	329	Total	C	N	O	S	0	4	0
			2539	1622	431	473	13			
1	C	329	Total	C	N	O	S	0	4	0
			2525	1614	427	472	12			
1	D	331	Total	C	N	O	S	0	4	1
			2524	1613	422	476	13			

- Molecule 2 is N-(2-METHYL-1,3-BENZOTHAZOL-6-YL)-3-UREIDO-PROPANAMIDE (CCD ID: 88S) (formula: C₁₂H₁₄N₄O₂S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			19	12	4	2	1		
2	B	1	Total	C	N	O	S	0	0
			19	12	4	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	S	0	0
			19	12	4	2	1		
2	D	1	Total	C	N	O	S	0	0
			19	12	4	2	1		

- Molecule 3 is MALONATE ION (CCD ID: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		
3	A	1	Total	C	O	0	0
			7	3	4		
3	B	1	Total	C	O	0	0
			7	3	4		
3	B	1	Total	C	O	0	0
			7	3	4		
3	C	1	Total	C	O	0	0
			7	3	4		
3	C	1	Total	C	O	0	0
			7	3	4		
3	D	1	Total	C	O	0	0
			7	3	4		
3	D	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	346	Total	O	0	0
			346	346		
5	B	331	Total	O	0	0
			331	331		
5	C	257	Total	O	0	0
			257	257		
5	D	218	Total	O	0	0
			218	218		

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

Position	Amino Acid	Conservation Score (bits)
1	A1	0.25
13	K13	0.15
15	E15	0.15
16	Q16	0.15
17	V17	0.15
54	E54	0.05
80	K80	0.05
81	D81	0.05
113	V113	0.05
131	K131	0.05
139	V139	0.05
140	D140	0.05
145	V145	0.05
171	Y171	0.05
180	H180	0.05
181	P181	0.05
191	E191	0.05
194	D194	0.05
227	K227	0.05
239	E239	0.05
247	T247	0.05
253	L253	0.05
273	S273	0.05
276	I276	0.05
280	Y280	0.25
281	G281	0.15
282	I282	0.15
285	D285	0.05
286	V286	0.05
287	F287	0.05
288	L288	0.05
293	I293	0.05
301	D301	0.05
314	R314	0.05
322	L322	0.05

Chain B:

Chain C: 92% 7%

Chain D:

Node	State	Percentage
A1	Green	
I8	Green	
V9	Green	
E15	Green	
Q16	Green	
T73	Yellow	
Y82	Yellow	
R98	Green	
Q99	Green	
Q100	Green	
V109	Yellow	
V113	Yellow	
I119	Yellow	
N122	Yellow	
V123	Yellow	
L132	Yellow	
V139	Yellow	
V145	Yellow	
F152	Yellow	
L164	Yellow	
G190	Yellow	
D222	Yellow	
T247	Yellow	
A250	Yellow	
L266	Yellow	
S273	Green	
T274	Green	
K275	Green	
I276	Green	
K277	Green	
G281	Green	
L288	Yellow	
D301	Yellow	
Q326	Yellow	
I330	Green	
F331	Green	

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.17Å 81.75Å 128.97Å 90.00° 96.14° 90.00°	Depositor
Resolution (Å)	29.39 – 2.03 29.39 – 2.03	Depositor EDS
% Data completeness (in resolution range)	98.1 (29.39-2.03) 98.5 (29.39-2.03)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 2.03Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.147 , 0.190 0.154 , 0.199	Depositor DCC
R_{free} test set	4080 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	18.6	Xtriage
Anisotropy	0.402	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 64.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11384	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 88S, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/2536	1.27	8/3445 (0.2%)
1	B	0.85	0/2581	1.22	14/3500 (0.4%)
1	C	0.85	0/2567	1.27	10/3484 (0.3%)
1	D	0.82	0/2567	1.27	10/3487 (0.3%)
All	All	0.85	0/10251	1.26	42/13916 (0.3%)

There are no bond length outliers.

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	16	GLN	N-CA-C	8.64	123.51	109.85
1	C	109	VAL	N-CA-C	7.60	118.34	110.36
1	D	109	VAL	N-CA-C	7.56	118.10	110.23
1	D	152	PHE	CA-CB-CG	7.18	120.98	113.80
1	C	150	SER	N-CA-C	7.12	119.04	111.28
1	A	282	ILE	N-CA-C	6.82	118.40	108.58
1	B	119	ILE	N-CA-C	6.51	116.61	110.30
1	D	139	VAL	N-CA-C	6.40	117.08	110.36
1	A	139	VAL	N-CA-C	6.32	117.83	110.62
1	B	109	VAL	N-CA-C	6.22	116.70	110.23
1	B	89	LYS	N-CA-C	-6.21	105.53	113.23
1	A	54	GLU	N-CA-C	6.17	118.51	111.11
1	C	139	VAL	N-CA-C	6.04	116.71	110.36
1	D	119	ILE	N-CA-C	6.03	116.15	110.30
1	C	80	LYS	N-CA-C	-5.83	105.06	111.82
1	C	222	ASP	CA-CB-CG	5.81	118.41	112.60
1	A	131	LYS	N-CA-C	-5.80	100.25	109.76
1	D	100	GLN	N-CA-C	-5.79	102.19	110.59
1	B	222	ASP	CA-CB-CG	5.71	118.31	112.60
1	A	194	ASP	N-CA-C	5.68	118.25	111.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	123	VAL	N-CA-CB	5.67	117.83	110.57
1	A	239	GLU	CB-CG-CD	5.67	122.23	112.60
1	B	249	TRP	N-CA-C	5.65	117.11	111.07
1	D	9	VAL	N-CA-CB	5.65	116.88	110.72
1	A	81	ASP	CA-CB-CG	5.62	118.22	112.60
1	D	15	GLU	N-CA-C	-5.51	98.75	108.23
1	C	131	LYS	N-CA-C	-5.49	100.75	109.59
1	A	80	LYS	N-CA-C	-5.47	106.10	112.89
1	B	124	VAL	CA-C-N	5.42	127.85	120.54
1	B	124	VAL	C-N-CA	5.42	127.85	120.54
1	B	54	GLU	N-CA-C	5.39	117.57	111.11
1	D	73	THR	CB-CA-C	5.38	115.82	110.33
1	B	139	VAL	N-CA-C	5.31	116.68	110.62
1	C	137	ASN	N-CA-C	5.31	117.30	110.39
1	D	326	GLN	N-CA-C	5.17	117.66	111.71
1	B	243	LEU	N-CA-C	5.16	117.69	111.40
1	C	8	ILE	N-CA-C	5.16	115.33	108.11
1	D	222	ASP	CA-CB-CG	5.15	117.75	112.60
1	C	123	VAL	CA-C-N	5.13	127.50	120.46
1	C	123	VAL	C-N-CA	5.13	127.50	120.46
1	B	16	GLN	CA-C-N	5.12	129.66	120.59
1	B	16	GLN	C-N-CA	5.12	129.66	120.59

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2494	0	2516	11	0
1	B	2539	0	2582	8	0
1	C	2525	0	2552	8	0
1	D	2524	0	2526	11	0
2	A	19	0	14	0	0
2	B	19	0	14	0	0
2	C	19	0	14	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	19	0	14	0	0
3	A	14	0	4	0	0
3	B	14	0	4	0	0
3	C	14	0	4	0	0
3	D	14	0	4	0	0
4	B	6	0	8	0	0
4	C	6	0	8	0	0
4	D	6	0	8	0	0
5	A	346	0	0	2	0
5	B	331	0	0	0	0
5	C	257	0	0	0	0
5	D	218	0	0	1	0
All	All	11384	0	10272	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:276[B]:ILE:HG13	1:C:288:LEU:HG	1.83	0.60
1:A:191:GLU:HG3	1:A:322:LEU:HD21	1.89	0.54
1:D:164:LEU:HD11	1:D:250:ALA:HB1	1.92	0.52
1:C:276[A]:ILE:HG12	1:C:288:LEU:HG	1.91	0.51
1:A:253:LEU:HD21	1:B:40[B]:MET:SD	2.51	0.51
1:A:293:ILE:HD12	1:A:301:ASP:HB2	1.91	0.51
1:D:113:VAL:HG21	1:D:329:LEU:HD22	1.93	0.51
1:D:82:TYR:CG	1:D:122:ASN:HB3	2.48	0.48
1:C:140:ASP:HB3	1:C:287:PHE:O	2.14	0.48
1:C:312:GLU:HG2	1:C:316:LYS:HE3	1.96	0.48
1:A:180:HIS:HB2	1:D:266:LEU:O	2.15	0.46
1:B:36:ILE:O	1:B:40[B]:MET:HG2	2.16	0.46
1:A:227:LYS:HG3	5:A:2261:HOH:O	2.15	0.46
1:B:86:ALA:HA	1:B:126:TYR:HB3	1.97	0.45
1:B:293:ILE:HD12	1:B:301:ASP:HB2	1.98	0.45
1:C:293:ILE:HD12	1:C:301:ASP:HB2	1.99	0.45
1:C:276[A]:ILE:HD12	1:C:282:ILE:HG21	1.99	0.45
1:D:98:ARG:HD2	5:D:2085:HOH:O	2.17	0.45
1:A:140:ASP:HB3	1:A:287:PHE:O	2.18	0.44
1:A:171:TYR:HA	1:A:181:PRO:HG3	2.00	0.44
1:A:113:VAL:HG22	1:A:145:VAL:HG21	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:190:GLY:HA2	1:C:288:LEU:HD13	2.00	0.43
1:D:123:VAL:HG11	1:D:132:LEU:HD21	2.00	0.43
1:B:291:PRO:HG2	1:B:305:VAL:CG2	2.49	0.43
1:D:113:VAL:HG22	1:D:145:VAL:HG21	2.01	0.42
1:A:314:ARG:HD3	5:A:2258:HOH:O	2.19	0.42
1:D:275:MET:HG2	1:D:277:LYS:H	1.85	0.42
1:B:280:TYR:HB2	1:B:282:ILE:HD12	2.02	0.41
1:B:301:ASP:HB3	1:D:8:ILE:HG23	2.02	0.41
1:B:10:ASN:HA	1:D:301:ASP:OD2	2.21	0.41
1:D:190:GLY:HA2	1:D:288:LEU:HD13	2.02	0.41
1:C:292:CYS:HB3	1:C:299:ILE:HG23	2.02	0.41
1:A:276:ILE:HG12	1:A:288:LEU:HG	2.02	0.40
1:A:276:ILE:HD13	1:A:282:ILE:HD13	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	327/331 (99%)	316 (97%)	11 (3%)	0	100	100
1	B	329/331 (99%)	322 (98%)	7 (2%)	0	100	100
1	C	329/331 (99%)	322 (98%)	7 (2%)	0	100	100
1	D	333/331 (101%)	318 (96%)	15 (4%)	0	100	100
All	All	1318/1324 (100%)	1278 (97%)	40 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	269/286 (94%)	264 (98%)	5 (2%)	50	50
1	B	279/286 (98%)	276 (99%)	3 (1%)	65	68
1	C	275/286 (96%)	271 (98%)	4 (2%)	57	59
1	D	272/286 (95%)	270 (99%)	2 (1%)	76	77
All	All	1095/1144 (96%)	1081 (99%)	14 (1%)	63	63

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	VAL
1	A	247	THR
1	A	273	SER
1	A	276	ILE
1	A	285	ASP
1	B	247	THR
1	B	273	SER
1	B	276	ILE
1	C	9	VAL
1	C	247	THR
1	C	276[A]	ILE
1	C	276[B]	ILE
1	D	247	THR
1	D	273	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	10	ASN
1	A	65	GLN
1	A	100	GLN
1	A	225	GLN
1	B	10	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	107	ASN
1	B	110	GLN
1	C	114	ASN
1	C	297	ASN
1	D	10	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	88S	A	1332	-	20,20,20	0.65	1 (5%)	27,27,27	0.92	2 (7%)
3	MLI	A	1334	-	6,6,6	1.07	0	7,7,7	1.05	0
4	GOL	B	1332	-	5,5,5	0.12	0	5,5,5	0.51	0
4	GOL	C	1332	-	5,5,5	0.13	0	5,5,5	0.27	0
3	MLI	D	1335	-	6,6,6	1.25	1 (16%)	7,7,7	0.77	0
2	88S	B	1333	-	20,20,20	0.77	1 (5%)	27,27,27	0.53	0
3	MLI	B	1335	-	6,6,6	1.20	0	7,7,7	0.99	0
3	MLI	C	1334	-	6,6,6	1.06	0	7,7,7	1.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MLI	D	1334	-	6,6,6	1.05	0	7,7,7	1.09	0
4	GOL	D	1332	-	5,5,5	0.06	0	5,5,5	0.17	0
2	88S	D	1333	-	20,20,20	0.60	1 (5%)	27,27,27	0.98	2 (7%)
3	MLI	C	1335	-	6,6,6	1.22	0	7,7,7	0.92	0
3	MLI	A	1333	-	6,6,6	1.10	0	7,7,7	1.05	0
3	MLI	B	1334	-	6,6,6	1.29	0	7,7,7	0.76	0
2	88S	C	1333	-	20,20,20	0.67	1 (5%)	27,27,27	0.71	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	88S	A	1332	-	-	0/10/10/10	0/2/2/2
3	MLI	A	1334	-	-	2/4/4/4	-
4	GOL	B	1332	-	-	0/4/4/4	-
4	GOL	C	1332	-	-	0/4/4/4	-
3	MLI	D	1335	-	-	2/4/4/4	-
2	88S	B	1333	-	-	0/10/10/10	0/2/2/2
3	MLI	B	1335	-	-	2/4/4/4	-
3	MLI	C	1334	-	-	2/4/4/4	-
3	MLI	D	1334	-	-	2/4/4/4	-
4	GOL	D	1332	-	-	0/4/4/4	-
2	88S	D	1333	-	-	0/10/10/10	0/2/2/2
3	MLI	C	1335	-	-	0/4/4/4	-
3	MLI	A	1333	-	-	2/4/4/4	-
3	MLI	B	1334	-	-	0/4/4/4	-
2	88S	C	1333	-	-	0/10/10/10	0/2/2/2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	1333	88S	C2-N3	2.48	1.32	1.29
2	D	1333	88S	C2-N3	2.24	1.31	1.29
2	B	1333	88S	C2-N3	2.22	1.31	1.29
3	D	1335	MLI	O9-C3	-2.03	1.24	1.30
2	A	1332	88S	C2-N3	2.02	1.31	1.29

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1333	88S	C1-C2-S10	4.13	122.08	120.08
2	A	1332	88S	C1-C2-S10	3.75	121.90	120.08
2	C	1333	88S	C1-C2-S10	2.77	121.42	120.08
2	D	1333	88S	C1-C2-N3	-2.20	122.49	124.01
2	A	1332	88S	C1-C2-N3	-2.09	122.56	124.01

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1333	MLI	C3-C1-C2-O7
3	D	1334	MLI	C3-C1-C2-O7
3	B	1335	MLI	C2-C1-C3-O8
3	B	1335	MLI	C2-C1-C3-O9
3	A	1333	MLI	C3-C1-C2-O6
3	D	1334	MLI	C3-C1-C2-O6
3	D	1335	MLI	C2-C1-C3-O9
3	A	1334	MLI	C2-C1-C3-O9
3	C	1334	MLI	C2-C1-C3-O8
3	A	1334	MLI	C2-C1-C3-O8
3	C	1334	MLI	C2-C1-C3-O9
3	D	1335	MLI	C2-C1-C3-O8

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 88S A 1332



Ligand 88S B 1333





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/331 (99%)	-0.58	4 (1%) 76 76	7, 19, 48, 71	2 (0%)
1	B	329/331 (99%)	-0.69	3 (0%) 81 81	7, 16, 40, 66	5 (1%)
1	C	329/331 (99%)	-0.62	3 (0%) 81 81	11, 19, 40, 56	5 (1%)
1	D	331/331 (100%)	-0.43	4 (1%) 76 76	9, 23, 48, 72	6 (1%)
All	All	1319/1324 (99%)	-0.58	14 (1%) 78 78	7, 19, 44, 72	18 (1%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	331	PHE	4.3
1	B	281	GLY	3.5
1	A	331	PHE	3.4
1	C	1	ALA	2.7
1	D	15	GLU	2.5
1	A	1	ALA	2.5
1	C	281	GLY	2.5
1	A	15	GLU	2.4
1	C	331	PHE	2.2
1	B	13	LYS	2.2
1	D	16	GLN	2.2
1	A	280	TYR	2.1
1	B	15	GLU	2.1
1	D	281	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	C	1332	6/6	0.85	0.11	38,41,41,43	0
4	GOL	B	1332	6/6	0.89	0.10	31,40,40,41	0
2	88S	D	1333	19/19	0.92	0.08	27,33,36,36	0
4	GOL	D	1332	6/6	0.92	0.07	31,36,38,38	0
3	MLI	D	1334	7/7	0.95	0.07	22,25,31,35	0
2	88S	A	1332	19/19	0.95	0.07	18,23,31,32	0
2	88S	B	1333	19/19	0.96	0.06	15,18,28,28	0
2	88S	C	1333	19/19	0.96	0.06	21,24,30,33	0
3	MLI	D	1335	7/7	0.96	0.05	17,21,27,29	0
3	MLI	A	1333	7/7	0.97	0.05	18,22,32,33	0
3	MLI	A	1334	7/7	0.97	0.05	20,23,24,26	0
3	MLI	B	1334	7/7	0.97	0.05	17,18,21,26	0
3	MLI	C	1334	7/7	0.97	0.06	12,15,22,24	0
3	MLI	C	1335	7/7	0.97	0.05	18,19,25,28	0
3	MLI	B	1335	7/7	0.99	0.03	17,19,22,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 88S D 1333:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 88S A 1332:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 88S B 1333:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 88S C 1333:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.