



wwPDB EM Validation Summary Report ⓘ

Mar 13, 2026 – 08:36 AM UTC

PDB ID : 6AJ3 / pdb_00006aj3
EMDB ID : EMD-9632
Title : The structure of Enterovirus D68 procapsid
Authors : Zheng, Q.B.; Zhu, R.; Xu, L.F.; He, M.Z.; Yan, X.D.; Cheng, T.; Li, S.W.
Deposited on : 2018-08-26
Resolution : 3.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : **FAILED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Q-score	25397	10198 (3.30 - 4.30)

MolProbity failed to run properly - the sequence quality summary graphics cannot be shown.

ENTRY-COMPOSITION INFOmissingINFO

SEQUENCE-PLOTS INFOmissingINFO

2 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	13849	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING ONLY	Depositor
Microscope	FEI TECNAI F30	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON II (4k x 4k)	Depositor
Maximum map value	0.206	Depositor
Minimum map value	-0.096	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.013	Depositor
Recommended contour level	0.0613	Depositor
Map size ($\text{\AA}$)	507.6, 507.6, 507.6	wwPDB
Map dimensions	450, 450, 450	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.128, 1.128, 1.128	Depositor

3 Model quality [i](#)

3.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3 Torsion angles [i](#)

3.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

3.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

3.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

3.6 Ligand geometry [i](#)

There are no ligands in this entry.

3.7 Other polymers [i](#)

There are no such residues in this entry.

3.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


4 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-9632. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

4.1 Orthogonal projections [i](#)


4.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

4.2 Central slices [i](#)


4.2.1 Primary map



X Index: 225



Y Index: 225




Z Index: 225


The images above show central slices of the map in three orthogonal directions.

4.3 Largest variance slices [i](#)


4.3.1 Primary map



X Index: 239



Y Index: 239




Z Index: 239


The images above show the largest variance slices of the map in three orthogonal directions.

4.4 Orthogonal standard-deviation projections (False-color) [i](#)


4.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

4.5 Orthogonal surface views [i](#)

4.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0613. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


4.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

5 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

5.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


5.2 Volume estimate [i](#)



The volume at the recommended contour level is 1338 nm^3 ; this corresponds to an approximate mass of 1209 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

5.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹

6 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

6.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.263 Å⁻¹

6.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.80	-	-
Author-provided FSC curve	3.83	4.18	3.87
Unmasked-calculated*	-	-	-


*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

7 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-9632 and PDB model 6AJ3. Per-residue inclusion information can be found in section ?? on page ??.

7.1 Map-model overlays


7.1.1 Map-model overlay [i](#)



X




Y




Z


7.1.2 Map-model assembly overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0613 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

7.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

7.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0613).

7.4 Atom inclusion [i](#)



At the recommended contour level, 78% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.

7.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0613) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6460	0.4720
A	0.6320	0.4720
B	0.6670	0.4780
C	0.6420	0.4680

