



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:56 AM UTC

PDB ID : 7AJ0 / pdb_00007aj0
Title : Crystal structure of PsFucS1 sulfatase from Pseudoalteromonas sp.
Authors : Roret, T.; Mikkelsen, M.D.; Czjzek, M.; Meyer, A.S.
Deposited on : 2020-09-28
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	92%5% .
1	B	515	92%5% .
1	C	515	92%5% .
1	D	515	% 90%6% . .
1	E	515	93%. .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	515	% 93%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 48247 atoms, of which 23325 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Arylsulfatase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	501	Total	C	H	N	O	S	0	0	0
			7824	2505	3893	669	738	19			
1	B	501	Total	C	H	N	O	S	0	0	0
			7794	2499	3871	667	738	19			
1	C	501	Total	C	H	N	O	S	0	0	0
			7824	2505	3893	669	738	19			
1	D	501	Total	C	H	N	O	S	0	0	0
			7824	2505	3893	669	738	19			
1	E	501	Total	C	H	N	O	S	0	0	0
			7824	2505	3893	669	738	19			
1	F	501	Total	C	H	N	O	S	0	0	0
			7809	2502	3882	668	738	19			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		
2	D	1	Total	Ca	0	0
			1	1		
2	E	1	Total	Ca	0	0
			1	1		
2	F	1	Total	Ca	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0
3	B	2	Total Cl 2 2	0	0
3	C	2	Total Cl 2 2	0	0
3	D	2	Total Cl 2 2	0	0
3	E	2	Total Cl 2 2	0	0
3	F	2	Total Cl 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	240	Total O 240 240	0	0
4	B	220	Total O 220 220	0	0
4	C	212	Total O 212 212	0	0
4	D	246	Total O 246 246	0	0
4	E	187	Total O 187 187	0	0
4	F	225	Total O 225 225	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Arylsulfatase




• Molecule 1: Arylsulfatase



• Molecule 1: Arylsulfatase



• Molecule 1: Arylsulfatase






● Molecule 1: Arylsulfatase



● Molecule 1: Arylsulfatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	86.20Å 182.34Å 209.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.66 – 2.50 46.66 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.9 (46.66-2.50) 99.9 (46.66-2.50)	Depositor EDS
R_{merge}	0.38	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.172 , 0.210 0.172 , 0.211	Depositor DCC
R_{free} test set	5746 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.207	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 26.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	48247	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/4027	0.32	0/5460
1	B	0.15	0/4019	0.33	0/5452
1	C	0.15	0/4027	0.32	0/5460
1	D	0.15	0/4027	0.33	0/5460
1	E	0.13	0/4027	0.32	0/5460
1	F	0.14	0/4023	0.32	0/5456
All	All	0.14	0/24150	0.32	0/32748

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3931	3893	3892	20	0
1	B	3923	3871	3870	14	0
1	C	3931	3893	3892	14	0
1	D	3931	3893	3892	27	0
1	E	3931	3893	3892	13	0
1	F	3927	3882	3881	13	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
3	E	2	0	0	0	0
3	F	2	0	0	0	0
4	A	240	0	0	1	0
4	B	220	0	0	2	0
4	C	212	0	0	1	0
4	D	246	0	0	4	0
4	E	187	0	0	3	0
4	F	225	0	0	3	0
All	All	24922	23325	23319	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (91) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:21:ASN:N	4:D:701:HOH:O	2.07	0.87
1:D:373:MET:SD	4:D:936:HOH:O	2.32	0.86
1:A:94:LYS:NZ	1:A:98:GLU:OE2	2.15	0.78
1:A:170:GLY:HA3	1:A:175:MET:HE2	1.66	0.76
1:A:481:MET:HE1	1:D:301:ARG:HG3	1.68	0.74
1:A:301:ARG:HG3	1:D:481:MET:HE1	1.71	0.71
1:B:454:ASN:OD1	4:B:701:HOH:O	2.11	0.68
1:A:421:ASP:O	4:A:701:HOH:O	2.12	0.66
1:C:498:MET:SD	4:C:881:HOH:O	2.52	0.66
1:D:157:VAL:HG11	1:D:399:VAL:HG12	1.78	0.66
1:D:41:GLU:OE2	4:D:702:HOH:O	2.14	0.65
1:E:30:LYS:NZ	4:E:702:HOH:O	2.26	0.64
1:A:133:ILE:HB	1:A:175:MET:HE1	1.78	0.63
1:F:21:ASN:ND2	4:F:702:HOH:O	2.32	0.62
1:D:166:LYS:HE3	1:D:191:VAL:HG23	1.81	0.61
1:F:248:MET:HE1	1:F:288:PRO:HD2	1.81	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:37:ILE:HD11	1:E:428:GLN:CG	2.29	0.61
1:A:25:ASP:OD2	1:B:216:LYS:NZ	2.34	0.59
1:D:480:ASN:C	1:D:481:MET:HE2	2.28	0.59
1:D:37:ILE:HD11	1:D:428:GLN:HG3	1.85	0.58
1:D:122:ARG:NH2	4:D:708:HOH:O	2.37	0.57
1:F:140:THR:HG22	1:F:520:ILE:O	2.04	0.57
1:A:301:ARG:CG	1:D:481:MET:HE1	2.34	0.56
1:D:395:ARG:HB2	1:D:400:ILE:HD11	1.87	0.56
1:A:481:MET:HE1	1:D:301:ARG:CG	2.33	0.56
1:E:37:ILE:HD11	1:E:428:GLN:HG2	1.88	0.56
1:B:492:GLN:OE1	1:C:460:ARG:NH2	2.34	0.54
1:F:122:ARG:NH2	4:F:709:HOH:O	2.40	0.52
1:E:37:ILE:HD11	1:E:428:GLN:HG3	1.90	0.51
1:E:229:ARG:NH1	4:E:712:HOH:O	2.46	0.49
1:B:139:VAL:CG2	1:B:520:ILE:HB	2.43	0.48
1:D:37:ILE:HD11	1:D:428:GLN:CG	2.43	0.48
1:F:254:ARG:NH1	1:F:313:GLU:OE1	2.43	0.47
1:D:188:LEU:HD12	1:D:188:LEU:N	2.29	0.47
1:C:71:LEU:HD13	1:C:304:MET:HE1	1.98	0.46
1:B:134:LEU:O	1:B:136:GLY:N	2.49	0.46
1:F:291:LYS:NZ	4:F:706:HOH:O	2.40	0.46
1:B:474:PRO:HD2	1:B:498:MET:HE3	1.99	0.45
1:B:472:THR:O	1:C:435:ARG:NH1	2.50	0.45
1:C:432:MET:SD	1:C:447:LYS:HE3	2.57	0.44
1:F:107:SER:HB3	1:F:364:ILE:HD13	2.00	0.44
1:B:221:ARG:NE	4:B:719:HOH:O	2.51	0.44
1:B:249:ILE:HG23	1:B:250:ASP:N	2.33	0.44
1:C:188:LEU:HD12	1:C:188:LEU:N	2.32	0.44
1:A:433:TRP:CZ2	1:A:512:MET:HE1	2.53	0.43
1:A:481:MET:HE3	1:D:296:ALA:HB1	2.00	0.43
1:F:188:LEU:HD12	1:F:188:LEU:N	2.33	0.43
1:B:188:LEU:HD23	1:B:188:LEU:H	1.83	0.43
1:C:403:LEU:HD22	1:C:409:PHE:CE2	2.54	0.43
1:C:249:ILE:HG23	1:C:250:ASP:N	2.33	0.42
1:E:139:VAL:HG23	1:E:520:ILE:HB	2.01	0.42
1:A:98:GLU:HA	1:A:376:ALA:HB1	2.00	0.42
1:A:105:MET:CE	1:A:389:ILE:HD11	2.50	0.42
1:A:188:LEU:HD12	1:A:188:LEU:N	2.34	0.42
1:B:432:MET:SD	1:B:447:LYS:HE3	2.59	0.42
1:E:441:LEU:HD12	1:E:441:LEU:C	2.45	0.42
1:E:494:PHE:HD2	1:E:498:MET:HE2	1.85	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:403:LEU:HD22	1:A:409:PHE:CZ	2.55	0.41
1:C:441:LEU:C	1:C:441:LEU:HD12	2.45	0.41
1:F:248:MET:HE1	1:F:288:PRO:CD	2.48	0.41
1:E:31:PRO:O	4:E:701:HOH:O	2.21	0.41
1:E:357:THR:HG21	1:E:439:THR:OG1	2.20	0.41
1:C:129:LEU:HD21	1:C:143:PRO:HG2	2.01	0.41
1:D:265:VAL:HG22	1:D:323:GLN:HG3	2.01	0.41
1:D:157:VAL:HG11	1:D:399:VAL:CG1	2.49	0.41
1:D:342:TRP:CG	1:D:343:PRO:HA	2.56	0.41
1:A:296:ALA:HB1	1:D:481:MET:HE3	2.02	0.41
1:C:433:TRP:CZ2	1:C:512:MET:HE1	2.56	0.41
1:D:105:MET:SD	1:D:384:MET:HG2	2.60	0.41
1:D:338:ASN:O	1:D:345:SER:HB3	2.21	0.41
1:A:176:LEU:HD12	1:A:234:ARG:NE	2.35	0.41
1:F:249:ILE:HG23	1:F:250:ASP:N	2.36	0.41
1:A:342:TRP:CG	1:A:343:PRO:HA	2.55	0.41
1:A:487:ALA:HB3	1:D:459:ARG:NH1	2.36	0.41
1:C:364:ILE:HB	1:C:365:PRO:HD2	2.03	0.41
1:D:191:VAL:HG22	1:D:192:GLN:N	2.36	0.41
1:D:432:MET:SD	1:D:447:LYS:HE3	2.61	0.41
1:D:498:MET:N	1:D:499:PRO:HD2	2.36	0.41
1:E:188:LEU:H	1:E:188:LEU:HD12	1.86	0.41
1:F:441:LEU:C	1:F:441:LEU:HD12	2.45	0.41
1:B:342:TRP:CG	1:B:343:PRO:HA	2.56	0.41
1:B:350:TRP:CZ3	1:B:483:PRO:HB2	2.55	0.40
1:C:74:GLY:O	1:C:81:GLY:HA3	2.21	0.40
1:F:432:MET:SD	1:F:447:LYS:HE3	2.61	0.40
1:C:98:GLU:HA	1:C:376:ALA:HB1	2.02	0.40
1:D:357:THR:HG21	1:D:439:THR:OG1	2.21	0.40
1:E:249:ILE:HG23	1:E:250:ASP:N	2.36	0.40
1:B:441:LEU:C	1:B:441:LEU:HD12	2.46	0.40
1:F:139:VAL:CG2	1:F:520:ILE:HB	2.52	0.40
1:A:390:TYR:CE1	1:A:394:LEU:HD11	2.57	0.40
1:E:133:ILE:HG21	1:E:175:MET:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	499/515 (97%)	483 (97%)	16 (3%)	0	100	100
1	B	499/515 (97%)	481 (96%)	15 (3%)	3 (1%)	21	38
1	C	499/515 (97%)	485 (97%)	14 (3%)	0	100	100
1	D	499/515 (97%)	475 (95%)	24 (5%)	0	100	100
1	E	499/515 (97%)	479 (96%)	19 (4%)	1 (0%)	43	63
1	F	499/515 (97%)	478 (96%)	20 (4%)	1 (0%)	43	63
All	All	2994/3090 (97%)	2881 (96%)	108 (4%)	5 (0%)	43	63

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	135	THR
1	B	138	LYS
1	F	138	LYS
1	B	136	GLY
1	E	136	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	421/433 (97%)	420 (100%)	1 (0%)	87	95
1	B	419/433 (97%)	415 (99%)	4 (1%)	68	86
1	C	421/433 (97%)	419 (100%)	2 (0%)	81	92

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	421/433 (97%)	415 (99%)	6 (1%)	59	81
1	E	421/433 (97%)	416 (99%)	5 (1%)	63	83
1	F	420/433 (97%)	414 (99%)	6 (1%)	59	81
All	All	2523/2598 (97%)	2499 (99%)	24 (1%)	68	86

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	364	ILE
1	B	188	LEU
1	B	219	GLN
1	B	318	LEU
1	B	491	HIS
1	C	123	LEU
1	C	335	ASN
1	D	22	VAL
1	D	138	LYS
1	D	188	LEU
1	D	345	SER
1	D	364	ILE
1	D	428	GLN
1	E	95	LEU
1	E	156	LYS
1	E	262	ASN
1	E	282	ASN
1	E	325	GLU
1	F	135	THR
1	F	188	LEU
1	F	213	LEU
1	F	325	GLU
1	F	335	ASN
1	F	345	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	282	ASN
1	A	326	ASN
1	B	91	ASN
1	B	454	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	91	ASN
1	C	168	HIS
1	C	263	GLN
1	C	335	ASN
1	C	378	GLN
1	D	168	HIS
1	E	91	ASN
1	E	108	GLN
1	E	168	HIS
1	E	219	GLN
1	F	168	HIS
1	F	491	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 18 ligands modelled in this entry, 18 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/515 (97%)	-0.34	2 (0%) 88 86	23, 29, 42, 57	0
1	B	501/515 (97%)	-0.28	2 (0%) 88 86	24, 32, 44, 57	0
1	C	501/515 (97%)	-0.26	2 (0%) 88 86	24, 32, 45, 62	0
1	D	501/515 (97%)	-0.34	4 (0%) 82 80	22, 29, 42, 52	0
1	E	501/515 (97%)	-0.27	1 (0%) 91 89	24, 32, 44, 57	0
1	F	501/515 (97%)	-0.26	5 (0%) 79 76	25, 31, 43, 57	0
All	All	3006/3090 (97%)	-0.29	16 (0%) 87 85	22, 31, 44, 62	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	521	GLY	3.6
1	F	139	VAL	3.1
1	D	400	ILE	2.9
1	D	304	MET	2.7
1	A	138	LYS	2.7
1	A	521	GLY	2.6
1	B	135	THR	2.5
1	F	520	ILE	2.5
1	D	406	ASN	2.5
1	F	140	THR	2.4
1	F	211	LYS	2.4
1	E	521	GLY	2.1
1	D	105	MET	2.1
1	C	141	GLN	2.1
1	B	138	LYS	2.1
1	C	521	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CL	C	602	1/1	0.91	0.13	53,53,53,53	0
3	CL	C	601	1/1	0.96	0.09	26,26,26,26	0
3	CL	B	602	1/1	0.97	0.07	44,44,44,44	0
2	CA	B	600	1/1	0.97	0.06	22,22,22,22	0
3	CL	A	602	1/1	0.97	0.10	35,35,35,35	0
3	CL	E	602	1/1	0.97	0.12	43,43,43,43	0
2	CA	C	600	1/1	0.98	0.04	31,31,31,31	0
3	CL	B	601	1/1	0.98	0.07	25,25,25,25	0
3	CL	D	603	1/1	0.98	0.10	24,24,24,24	0
3	CL	E	601	1/1	0.98	0.11	28,28,28,28	0
2	CA	D	601	1/1	0.98	0.03	22,22,22,22	0
3	CL	F	601	1/1	0.98	0.11	24,24,24,24	0
3	CL	F	602	1/1	0.98	0.06	44,44,44,44	0
2	CA	A	600	1/1	0.99	0.03	22,22,22,22	0
2	CA	E	600	1/1	0.99	0.07	22,22,22,22	0
3	CL	D	602	1/1	0.99	0.08	42,42,42,42	0
2	CA	F	600	1/1	0.99	0.05	22,22,22,22	0
3	CL	A	601	1/1	1.00	0.08	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CA B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around CA C 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA D 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CA A 600:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA E 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.