



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:06 PM UTC

PDB ID : 5AKJ / pdb_00005akj
Title : ligand complex structure of soluble epoxide hydrolase
Authors : Oster, L.; Tapani, S.; Xue, Y.; Kack, H.
Deposited on : 2015-03-03
Resolution : 2.03 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	13299 (2.04-2.00)
Clashscore	190562	1022 (2.02-2.02)
Ramachandran outliers	187476	1014 (2.02-2.02)
Sidechain outliers	187428	1014 (2.02-2.02)
RSRZ outliers	180081	13314 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	549	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BIFUNCTIONAL EPOXIDE HYDROLASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	546	4359	2792	735	794	38	0	4	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP P34913

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		
3	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 4 is 2-(1H-BENZIMIDAZOL-2-YLSULFANYL)ETHANOL (CCD ID: 6N4) (formula: C₉H₁₀N₂OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	13	9	2	1	1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	484	Total	O	0	0
			484	484		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BIFUNCTIONAL EPOXIDE HYDROLASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.57Å 92.57Å 244.22Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.41 – 2.03 29.41 – 2.03	Depositor EDS
% Data completeness (in resolution range)	97.9 (29.41-2.03) 97.8 (29.41-2.03)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.03Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.184 , 0.223 (Not available) , 0.188	Depositor DCC
R_{free} test set	2033 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	24.5	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 54.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4886	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 6N4, SO4, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.87	0/4464	1.32	21/6046 (0.3%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	330	VAL	N-CA-CB	7.76	118.99	110.53
1	A	160	LYS	N-CA-C	-7.27	100.92	109.93
1	A	292	ASP	N-CA-C	-7.07	96.12	108.20
1	A	305	ILE	CA-C-N	6.95	131.82	120.63
1	A	305	ILE	C-N-CA	6.95	131.82	120.63
1	A	545	ASP	N-CA-C	6.55	121.94	113.88
1	A	480	LEU	N-CA-C	6.42	119.09	111.71
1	A	144	LYS	N-CA-C	6.11	118.73	111.71
1	A	312	VAL	N-CA-CB	5.89	119.42	110.58
1	A	381	PHE	CA-CB-CG	-5.79	108.01	113.80
1	A	505	GLN	N-CA-C	5.79	119.16	111.75
1	A	444	GLU	CA-C-N	5.74	127.90	120.44
1	A	444	GLU	C-N-CA	5.74	127.90	120.44
1	A	388	GLN	N-CA-C	5.62	117.21	111.14
1	A	500	VAL	N-CA-CB	5.62	115.78	109.99
1	A	460	ARG	N-CA-C	5.46	116.93	110.97
1	A	412	SER	N-CA-C	5.42	118.68	111.75
1	A	516	ARG	CA-C-N	5.28	126.47	121.46
1	A	516	ARG	C-N-CA	5.28	126.47	121.46
1	A	184	ASP	CA-CB-CG	5.16	117.75	112.60
1	A	180	VAL	N-CA-C	5.07	115.27	108.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4359	0	4362	53	0
2	A	10	0	0	0	0
3	A	20	0	30	4	0
4	A	13	0	10	1	0
5	A	484	0	0	3	0
All	All	4886	0	4402	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:322:ASP:OD1	1:A:351[A]:ARG:NH2	2.00	0.92
1:A:547:ARG:HH11	1:A:547:ARG:HG3	1.37	0.89
1:A:376:LYS:HA	1:A:376:LYS:HE2	1.59	0.84
1:A:34:ARG:HH11	1:A:34:ARG:HG2	1.41	0.83
1:A:425:ALA:HB1	1:A:430:VAL:HG21	1.59	0.82
1:A:85:ASN:O	3:A:1554:DMS:H23	1.83	0.78
1:A:440:ARG:HH11	1:A:440:ARG:HG3	1.51	0.76
1:A:339:MET:HE2	1:A:343:TYR:HE2	1.49	0.76
1:A:547:ARG:HH11	1:A:547:ARG:CG	2.02	0.72
1:A:339:MET:HE2	1:A:343:TYR:CE2	2.28	0.68
1:A:309:CYS:SG	1:A:312:VAL:HG13	2.34	0.67
1:A:452:GLN:HE22	1:A:455:LYS:NZ	1.94	0.64
1:A:34:ARG:HG2	1:A:34:ARG:NH1	2.12	0.62
1:A:547:ARG:HG3	1:A:547:ARG:NH1	2.11	0.62
1:A:375:ILE:HG22	1:A:376:LYS:HE3	1.82	0.61
1:A:376:LYS:HE2	1:A:376:LYS:CA	2.31	0.60
1:A:452:GLN:HE22	1:A:455:LYS:HZ2	1.53	0.56
1:A:387:PHE:HE1	1:A:428:LEU:HD21	1.70	0.55
1:A:266:GLY:HA3	1:A:335:ASP:HB3	1.89	0.54
1:A:440:ARG:HH11	1:A:440:ARG:CG	2.19	0.54
1:A:142:GLU:HA	1:A:145[A]:MET:HE3	1.91	0.53
1:A:546:ALA:O	1:A:547:ARG:HB2	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:184:ASP:HB3	1:A:190:LEU:HD13	1.92	0.52
1:A:387:PHE:CE1	1:A:428:LEU:HD21	2.47	0.50
1:A:315:LYS:HD3	5:A:2303:HOH:O	2.10	0.49
1:A:339:MET:HE3	1:A:339:MET:HA	1.95	0.49
1:A:452:GLN:NE2	1:A:455:LYS:NZ	2.60	0.49
1:A:410:ARG:NH2	1:A:437:SER:O	2.46	0.49
1:A:199:VAL:HG11	1:A:216:VAL:HG21	1.96	0.47
1:A:369:MET:HG2	1:A:373:GLU:HB3	1.97	0.47
1:A:315:LYS:HD2	1:A:348:TYR:OH	2.15	0.47
1:A:44:GLY:HA2	3:A:1553:DMS:H22	1.97	0.46
1:A:442:VAL:HG21	1:A:447:ILE:HD11	1.98	0.46
1:A:410:ARG:HB2	1:A:416:VAL:HG12	1.97	0.46
1:A:22:VAL:HG21	1:A:95:ALA:CB	2.45	0.46
1:A:245:LYS:HB2	1:A:246:PRO:HD2	1.99	0.45
1:A:460:ARG:HG3	1:A:461:GLY:N	2.32	0.44
1:A:44:GLY:H	3:A:1553:DMS:H22	1.82	0.43
1:A:168:PHE:O	1:A:172:THR:HG23	2.19	0.43
1:A:233:ASN:HD22	1:A:234:PRO:HD2	1.83	0.43
1:A:519:ILE:HG22	1:A:522:CYS:HB2	2.00	0.43
1:A:417:LEU:HD11	1:A:429:PHE:HE1	1.84	0.43
1:A:164:GLN:HG2	5:A:2170:HOH:O	2.20	0.42
1:A:335:ASP:OD1	4:A:1555:6N4:N7	2.52	0.42
1:A:20:PHE:CZ	1:A:41:PHE:CE2	3.08	0.42
1:A:85:ASN:O	3:A:1554:DMS:C2	2.63	0.41
1:A:304:GLU:HB3	1:A:306:GLU:OE2	2.21	0.41
1:A:419:MET:HE1	1:A:428:LEU:HD12	2.03	0.41
1:A:11:ASP:CG	1:A:19:VAL:HG12	2.45	0.40
1:A:55:LYS:O	1:A:157:GLY:HA2	2.21	0.40
1:A:60:LEU:HD22	1:A:126:TRP:NE1	2.36	0.40
1:A:421:LYS:HA	1:A:421:LYS:HD3	1.64	0.40
1:A:245:LYS:NZ	5:A:2249:HOH:O	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	548/549 (100%)	535 (98%)	12 (2%)	1 (0%)	43	40

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	268	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	476/474 (100%)	440 (92%)	36 (8%)	12	6

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	20	PHE
1	A	25	ARG
1	A	72	ARG
1	A	84	LYS
1	A	97	SER
1	A	107	GLN
1	A	115	LYS
1	A	132	GLU
1	A	172	THR
1	A	190	LEU
1	A	196	LEU
1	A	207	ASP
1	A	222	LEU
1	A	224	THR
1	A	249	ARG
1	A	253	VAL
1	A	306	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	310	MET
1	A	312	VAL
1	A	315	LYS
1	A	330	VAL
1	A	353	ARG
1	A	370	SER
1	A	375	ILE
1	A	376	LYS
1	A	382	ASP
1	A	421	LYS
1	A	424	GLU
1	A	428	LEU
1	A	434	GLU
1	A	440	ARG
1	A	460	ARG
1	A	471	ARG
1	A	502	GLN
1	A	547	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	GLN
1	A	233	ASN
1	A	452	GLN
1	A	513	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	DMS	A	1553	-	3,3,3	0.84	0	3,3,3	0.80	0
3	DMS	A	1551	-	3,3,3	0.77	0	3,3,3	0.41	0
2	SO4	A	1548	-	4,4,4	0.63	0	6,6,6	0.15	0
2	SO4	A	1549	-	4,4,4	0.36	0	6,6,6	0.06	0
3	DMS	A	1550	-	3,3,3	0.73	0	3,3,3	1.29	0
3	DMS	A	1552	-	3,3,3	0.78	0	3,3,3	0.85	0
4	6N4	A	1555	-	14,14,14	2.82	8 (57%)	18,18,18	1.56	2 (11%)
3	DMS	A	1554	-	3,3,3	0.82	0	3,3,3	1.45	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	6N4	A	1555	-	-	2/4/4/4	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1555	6N4	C1-C6	5.07	1.47	1.38
4	A	1555	6N4	C2-C3	4.03	1.45	1.38
4	A	1555	6N4	C8-S10	-4.01	1.70	1.75
4	A	1555	6N4	C3-C4	3.33	1.45	1.39
4	A	1555	6N4	C2-C1	3.31	1.45	1.38
4	A	1555	6N4	C8-N7	2.90	1.36	1.30
4	A	1555	6N4	C6-C5	2.40	1.43	1.40
4	A	1555	6N4	C11-S10	2.28	1.91	1.81

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1555	6N4	C4-C5-N7	-3.11	105.17	109.42
4	A	1555	6N4	N9-C8-N7	-3.08	109.69	113.25
3	A	1554	DMS	C2-S-C1	-2.08	87.75	98.42

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1555	6N4	C12-C11-S10-C8
4	A	1555	6N4	N7-C8-S10-C11

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1553	DMS	2	0
4	A	1555	6N4	1	0
3	A	1554	DMS	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	546/549 (99%)	-0.19	7 (1%) 75 75	9, 25, 49, 95	4 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	20	PHE	3.6
1	A	547	ARG	3.6
1	A	19	VAL	3.1
1	A	376	LYS	3.0
1	A	378	ASN	2.2
1	A	18	ALA	2.0
1	A	420	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	DMS	A	1553	4/4	0.69	0.21	55,59,60,66	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DMS	A	1552	4/4	0.80	0.19	65,65,68,75	0
3	DMS	A	1554	4/4	0.81	0.15	60,66,67,71	0
3	DMS	A	1550	4/4	0.82	0.25	53,66,67,69	0
2	SO4	A	1549	5/5	0.89	0.15	74,78,79,80	0
2	SO4	A	1548	5/5	0.94	0.17	49,53,54,54	0
4	6N4	A	1555	13/13	0.94	0.08	26,27,29,30	0
3	DMS	A	1551	4/4	0.97	0.09	33,36,42,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.