



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:12 PM UTC

PDB ID : 7AKH / pdb_00007akh
Title : Structure of DYRK2 in complex with compound 58
Authors : Dokurno, P.; Surgenor, A.E.; Kotschy, A.
Deposited on : 2020-10-01
Resolution : 2.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	429	 65% 25% 6%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3237 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Dual specificity tyrosine-phosphorylation-regulated kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	402	3185	2017	559	583	3	23	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	MET	-	initiating methionine	UNP Q92630
A	125	HIS	-	expression tag	UNP Q92630
A	126	HIS	-	expression tag	UNP Q92630
A	127	HIS	-	expression tag	UNP Q92630
A	128	HIS	-	expression tag	UNP Q92630
A	129	HIS	-	expression tag	UNP Q92630
A	130	HIS	-	expression tag	UNP Q92630
A	131	SER	-	expression tag	UNP Q92630
A	132	SER	-	expression tag	UNP Q92630
A	133	GLY	-	expression tag	UNP Q92630
A	134	VAL	-	expression tag	UNP Q92630
A	135	ASP	-	expression tag	UNP Q92630
A	136	LEU	-	expression tag	UNP Q92630
A	137	GLY	-	expression tag	UNP Q92630
A	138	THR	-	expression tag	UNP Q92630
A	139	GLU	-	expression tag	UNP Q92630
A	140	ASN	-	expression tag	UNP Q92630
A	141	LEU	-	expression tag	UNP Q92630
A	142	TYR	-	expression tag	UNP Q92630
A	143	PHE	-	expression tag	UNP Q92630
A	144	GLN	-	expression tag	UNP Q92630
A	145	SER	-	expression tag	UNP Q92630

- Molecule 2 is 4-[3-[(2 {S})-2-(6-bromanylpyridin-2-yl)oxypropyl]-2-methyl-imidazo[4,5-b]pyridin-5-yl]pyridine-2,6-diamine (CCD ID: RK8) (formula: C₂₀H₂₀BrN₇O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	Br	C	N	O	0	0
			29	1	20	7	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	22	Total	O	0	0
			22	22		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity tyrosine-phosphorylation-regulated kinase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 42 21 2	Depositor
Cell constants a, b, c, α , β , γ	84.24Å 84.24Å 150.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.85 20.00 – 2.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-2.85) 99.5 (20.00-2.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.52 (at 2.83Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.185 , 0.245 0.188 , 0.243	Depositor DCC
R_{free} test set	655 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	73.7	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 31.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3237	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PTR, SEP, CL, RK8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	0/3221	1.43	3/4350 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	292	THR	CA-CB-OG1	-6.55	99.78	109.60
1	A	491	PRO	N-CA-C	5.53	120.28	111.26
1	A	244	VAL	N-CA-C	-5.01	108.41	113.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3185	0	3084	76	1
2	A	29	0	0	3	0
3	A	1	0	0	0	0
4	A	22	0	0	0	0
All	All	3237	0	3084	76	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:481:ARG:CZ	1:A:487:LEU:HD23	1.98	0.92
1:A:481:ARG:HG2	1:A:487:LEU:HD22	1.53	0.89
1:A:514:GLN:HB3	1:A:524:MET:HE2	1.68	0.73
1:A:401:MET:N	1:A:402:PRO:HD2	2.07	0.69
1:A:307:ASN:HA	1:A:355:LEU:HD12	1.76	0.68
1:A:490:PRO:O	1:A:493:SER:HB3	1.97	0.65
1:A:509:LEU:HD12	1:A:509:LEU:O	1.96	0.65
1:A:212:GLN:NE2	1:A:212:GLN:HA	2.11	0.65
1:A:232:SEP:O2P	1:A:232:SEP:N	2.31	0.64
1:A:315:ASN:O	1:A:315:ASN:ND2	2.32	0.63
1:A:315:ASN:HD21	1:A:318:GLN:HB2	1.67	0.59
1:A:364:ILE:O	1:A:364:ILE:HD12	2.04	0.58
1:A:514:GLN:CB	1:A:524:MET:HE2	2.35	0.57
1:A:440:MET:SD	1:A:462:PRO:HG3	2.45	0.57
1:A:256:GLU:HB2	1:A:259:PHE:HD2	1.70	0.56
1:A:509:LEU:HD11	1:A:513:LYS:CE	2.35	0.56
1:A:437:LEU:HD13	1:A:494:ARG:HD3	1.88	0.55
1:A:481:ARG:CZ	1:A:487:LEU:CD2	2.81	0.55
1:A:233:PHE:CZ	2:A:601:RK8:BR1	3.15	0.55
1:A:233:PHE:HZ	2:A:601:RK8:BR1	2.46	0.54
1:A:407:SER:O	1:A:411:ILE:HG13	2.05	0.54
1:A:155:THR:OG1	1:A:158:GLN:HG3	2.08	0.54
1:A:204:ASP:CG	1:A:296:HIS:HE2	2.15	0.54
1:A:265:GLU:OE1	1:A:268:ARG:NH2	2.39	0.53
1:A:481:ARG:NH1	1:A:487:LEU:HD23	2.22	0.53
1:A:364:ILE:HD12	1:A:364:ILE:C	2.34	0.52
1:A:509:LEU:HD11	1:A:513:LYS:HE3	1.92	0.51
1:A:315:ASN:HD22	1:A:318:GLN:H	1.58	0.51
1:A:228:ILE:HG21	1:A:238:LYS:HB2	1.92	0.51
1:A:139:GLU:O	1:A:139:GLU:HG3	2.12	0.50
1:A:284:VAL:O	1:A:365:LYS:HE2	2.12	0.50
1:A:156:PRO:HG3	1:A:178:TYR:O	2.12	0.50
1:A:136:LEU:HD22	1:A:154:MET:CE	2.42	0.50
1:A:455:PHE:C	1:A:463:ARG:HG3	2.36	0.49
1:A:438:LEU:HD13	1:A:516:LEU:O	2.12	0.49
1:A:315:ASN:ND2	1:A:318:GLN:HB2	2.28	0.49
1:A:436:GLU:HG2	1:A:494:ARG:HB2	1.94	0.49
1:A:469:THR:OG1	1:A:475:VAL:HG23	2.13	0.49
1:A:322:LEU:N	1:A:323:PRO:CD	2.76	0.48
1:A:465:CYS:HB3	1:A:479:GLY:HA2	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:GLU:H	1:A:157:GLU:CD	2.21	0.47
1:A:322:LEU:HD12	1:A:322:LEU:O	2.14	0.47
1:A:307:ASN:CA	1:A:355:LEU:HD12	2.42	0.47
1:A:345:ILE:HD11	1:A:375:GLU:HA	1.96	0.47
1:A:212:GLN:HA	1:A:212:GLN:HE21	1.79	0.47
1:A:438:LEU:HD23	1:A:438:LEU:N	2.30	0.47
1:A:509:LEU:HD11	1:A:513:LYS:HE2	1.96	0.47
1:A:438:LEU:CD1	1:A:516:LEU:O	2.63	0.47
1:A:320:PHE:CE2	1:A:415:LEU:HD22	2.50	0.46
1:A:509:LEU:HD12	1:A:509:LEU:C	2.40	0.45
1:A:182:TYR:CZ	1:A:243:LYS:HD2	2.52	0.45
1:A:160:MET:HE1	1:A:172:HIS:CG	2.53	0.44
1:A:160:MET:HE1	1:A:172:HIS:CD2	2.52	0.44
1:A:400:GLY:O	1:A:403:ILE:HG12	2.17	0.44
1:A:458:SEP:O2P	1:A:458:SEP:N	2.50	0.44
1:A:496:TRP:O	1:A:499:ALA:HB3	2.17	0.44
1:A:153:PRO:HG2	1:A:180:GLU:HB3	2.00	0.43
1:A:532:HIS:CG	1:A:533:PRO:HD2	2.54	0.43
1:A:394:ILE:O	1:A:451:ARG:NH1	2.52	0.43
1:A:355:LEU:HD22	1:A:367:ILE:HG21	2.01	0.43
1:A:438:LEU:HD11	1:A:516:LEU:HB3	2.01	0.42
1:A:387:PHE:HB2	1:A:388:TYR:CE1	2.54	0.42
1:A:199:ASN:HB3	1:A:204:ASP:O	2.19	0.42
1:A:355:LEU:HD21	2:A:601:RK8:C3	2.49	0.42
1:A:136:LEU:HD22	1:A:154:MET:HE3	2.02	0.42
1:A:529:ALA:O	1:A:535:LEU:HD12	2.20	0.42
1:A:421:LEU:O	1:A:422:LEU:HD23	2.20	0.42
1:A:445:LEU:HD12	1:A:445:LEU:HA	1.88	0.42
1:A:252:MET:HE3	1:A:252:MET:HB3	1.93	0.41
1:A:435:ILE:HA	1:A:439:GLY:O	2.20	0.41
1:A:456:VAL:HG13	1:A:460:GLY:HA2	2.02	0.41
1:A:186:LEU:H	1:A:186:LEU:HG	1.66	0.41
1:A:345:ILE:O	1:A:372:SER:HA	2.20	0.41
1:A:415:LEU:HD23	1:A:415:LEU:HA	1.85	0.41
1:A:240:TYR:OH	1:A:245:HIS:HA	2.20	0.40
1:A:449:SER:HB2	1:A:452:ALA:HB2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:139:GLU:OE2	1:A:361:ARG:NH1[6_555]	2.08	0.12

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/429 (92%)	369 (93%)	28 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	332/372 (89%)	292 (88%)	40 (12%)	5	9

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	VAL
1	A	164	MET
1	A	177	SER
1	A	187	ASN
1	A	191	ARG
1	A	195	THR
1	A	209	SER
1	A	226	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	230	LYS
1	A	235	GLN
1	A	237	VAL
1	A	252	MET
1	A	256	GLU
1	A	265	GLU
1	A	269	ILE
1	A	279	ASP
1	A	289	GLU
1	A	322	LEU
1	A	355	LEU
1	A	359	GLN
1	A	361	ARG
1	A	365	LYS
1	A	371	SER
1	A	375	GLU
1	A	383	ILE
1	A	427	GLU
1	A	435	ILE
1	A	444	LYS
1	A	445	LEU
1	A	447	ASP
1	A	467	VAL
1	A	469	THR
1	A	475	VAL
1	A	483	ARG
1	A	487	LEU
1	A	493	SER
1	A	495	GLU
1	A	507	LEU
1	A	509	LEU
1	A	512	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	140	ASN
1	A	158	GLN
1	A	192	GLN
1	A	212	GLN
1	A	262	GLN
1	A	272	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	276	GLN
1	A	315	ASN
1	A	330	HIS
1	A	342	ASN
1	A	478	ASN
1	A	528	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	PTR	A	382	1	15,16,17	0.91	1 (6%)	17,22,24	0.93	0
1	SEP	A	458	1	8,9,10	0.71	0	7,12,14	1.47	1 (14%)
1	SEP	A	232	1	8,9,10	0.80	0	7,12,14	0.78	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	382	1	-	0/10/11/13	0/1/1/1
1	SEP	A	458	1	-	1/6/8/10	-
1	SEP	A	232	1	-	3/6/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	382	PTR	P-OH	-2.84	1.54	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	458	SEP	O3P-P-OG	2.40	112.93	106.67

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	232	SEP	C-CA-CB-OG
1	A	232	SEP	CA-CB-OG-P
1	A	232	SEP	CB-OG-P-O1P
1	A	458	SEP	CA-CB-OG-P

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	458	SEP	1	0
1	A	232	SEP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RK8	A	601	-	32,32,32	0.56	0	43,46,46	2.92	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RK8	A	601	-	-	3/12/12/12	0/4/4/4

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	RK8	O1-C16-N7	-12.85	110.24	119.49
2	A	601	RK8	O1-C16-C17	10.97	128.05	115.85
2	A	601	RK8	O1-C2-C3	-4.31	100.85	105.97
2	A	601	RK8	BR1-C20-C19	3.15	121.89	118.69
2	A	601	RK8	C1-C2-C3	2.77	117.84	111.69
2	A	601	RK8	C11-C15-C14	2.64	120.04	117.35
2	A	601	RK8	N3-C10-N1	2.38	129.66	125.67
2	A	601	RK8	BR1-C20-N7	-2.18	113.74	115.68

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	RK8	N7-C16-O1-C2
2	A	601	RK8	C17-C16-O1-C2
2	A	601	RK8	C1-C2-C3-N1

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	RK8	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	399/429 (93%)	-0.58	0 100 100	47, 74, 112, 176	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	A	458	10/11	0.84	0.09	73,104,113,128	0
1	SEP	A	232	10/11	0.89	0.10	69,94,109,120	4
1	PTR	A	382	16/17	0.98	0.04	54,60,67,70	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	RK8	A	601	29/29	0.92	0.09	43,63,131,137	1
3	CL	A	602	1/1	0.96	0.08	70,70,70,70	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.