



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 01:42 AM UTC

PDB ID : 6ALA / pdb_00006ala
Title : Crystal structure of H108A peptidylglycine alpha-hydroxylating monooxygenase (PHM) in complex with citrate
Authors : Maheshwari, S.; Rudzka, K.; Gabelli, S.B.; Amzel, L.M.
Deposited on : 2017-08-07
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	C	312	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5003 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peptidyl-glycine alpha-amidating monooxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	297	Total	C	N	O	S	0	0	0
			2324	1482	393	424	25			
1	C	304	Total	C	N	O	S	0	1	0
			2383	1520	403	435	25			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	108	ALA	HIS	engineered mutation	UNP P14925
C	108	ALA	HIS	engineered mutation	UNP P14925

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: C₆H₅O₇) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	C	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0
4	C	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	95	Total O 95 95	0	0
5	C	95	Total O 95 95	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptidyl-glycine alpha-amidating monooxygenase



- Molecule 1: Peptidyl-glycine alpha-amidating monooxygenase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	171.37Å 52.46Å 116.46Å 90.00° 128.74° 90.00°	Depositor
Resolution (Å)	29.05 – 2.59 29.05 – 2.59	Depositor EDS
% Data completeness (in resolution range)	98.4 (29.05-2.59) 98.4 (29.05-2.59)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.21 (at 2.57Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.178 , 0.238 0.188 , 0.245	Depositor DCC
R_{free} test set	1234 reflections (4.71%)	wwPDB-VP
Wilson B-factor (Å ²)	29.1	Xtriage
Anisotropy	0.028	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5003	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.52% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CU, FLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	0/2390	1.07	6/3249 (0.2%)
1	C	1.00	1/2452 (0.0%)	1.04	6/3336 (0.2%)
All	All	1.00	1/4842 (0.0%)	1.06	12/6585 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	72	THR	CA-CB	5.53	1.58	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ALA	CA-C-N	-7.22	113.24	123.20
1	A	142	ALA	C-N-CA	-7.22	113.24	123.20
1	C	133	ASP	N-CA-CB	-6.49	99.52	110.49
1	A	338	VAL	CB-CA-C	-5.88	104.11	112.22
1	A	103	MET	N-CA-C	5.57	118.07	111.33
1	C	220	VAL	N-CA-C	5.54	116.03	107.78
1	A	215	PRO	O-C-N	5.39	123.79	121.31
1	C	338	VAL	CB-CA-C	-5.26	104.96	112.22
1	C	214	ILE	CA-C-N	-5.19	115.92	119.66
1	C	214	ILE	C-N-CA	-5.19	115.92	119.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	PHE	N-CA-C	5.04	118.74	112.59
1	C	307	GLY	N-CA-C	5.01	118.77	110.55

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	132	THR	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2324	0	2251	30	0
1	C	2383	0	2308	26	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	13	0	5	1	0
3	C	13	0	5	0	0
4	A	42	0	56	7	0
4	C	36	0	48	0	0
5	A	95	0	0	3	0
5	C	95	0	0	2	0
All	All	5003	0	4673	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ARG:HH21	1:A:100:ARG:NE	1.62	0.96
1:A:67:ARG:HH21	1:A:100:ARG:HE	0.94	0.91
1:C:114:CYS:HB2	1:C:133:ASP:OD2	1.82	0.80
1:A:67:ARG:NH2	1:A:100:ARG:HE	1.78	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:ARG:HH11	1:A:351:ASN:HD21	1.39	0.70
1:A:127:ASP:OD1	1:A:128:GLU:N	2.28	0.66
1:C:105:THR:HG22	1:C:178:ALA:HB1	1.77	0.65
1:C:235[A]:HIS:ND1	1:C:282:ASP:OD1	2.31	0.63
1:C:208:MET:HE3	1:C:314:MET:CE	2.28	0.63
1:A:205:TYR:OH	4:A:1003:GOL:H12	1.98	0.62
1:C:58:ASP:O	1:C:60:SER:N	2.33	0.62
1:A:138:LEU:HD11	4:A:1007:GOL:C3	2.29	0.62
1:C:236:VAL:HG12	1:C:277:VAL:HG21	1.82	0.61
1:C:226:SER:O	1:C:338:VAL:HG22	2.02	0.60
1:A:138:LEU:HD11	4:A:1007:GOL:H31	1.82	0.59
1:A:252:GLY:HA3	1:A:317:LEU:HD21	1.86	0.56
1:C:133:ASP:OD1	5:C:1101:HOH:O	2.17	0.56
1:C:119:SER:HB3	1:C:124:TRP:CD2	2.42	0.55
1:A:105:THR:HG22	1:A:178:ALA:HB1	1.87	0.55
1:C:201:ILE:HG12	1:C:325:ALA:HA	1.88	0.53
1:C:288:ILE:HG21	1:C:342:MET:HE1	1.92	0.52
1:A:67:ARG:NH2	1:A:100:ARG:NE	2.45	0.51
1:A:252:GLY:HA3	1:A:317:LEU:CD2	2.40	0.51
1:A:93:PHE:CE1	1:A:197:PRO:HA	2.47	0.50
1:A:210:VAL:O	4:A:1008:GOL:O2	2.29	0.48
1:C:133:ASP:OD1	1:C:134:LYS:N	2.46	0.48
1:C:103:MET:HG3	1:C:143:ARG:HA	1.94	0.48
1:A:254:ARG:NH1	1:A:281:VAL:HG22	2.29	0.47
1:A:235:HIS:HB2	5:A:1104:HOH:O	2.13	0.47
1:C:260:TRP:CZ3	1:C:346:ILE:HG13	2.49	0.47
1:C:115:ASN:ND2	1:C:165:LYS:HG2	2.30	0.47
4:A:1006:GOL:C3	5:A:1166:HOH:O	2.62	0.46
1:A:154:VAL:HG22	1:A:200:LEU:HB2	1.97	0.46
1:A:90:GLU:OE2	1:A:195:ARG:NH2	2.49	0.46
4:A:1006:GOL:H32	5:A:1166:HOH:O	2.16	0.45
1:A:236:VAL:HG12	1:A:277:VAL:HG21	1.99	0.45
3:A:1002:FLC:OG1	3:A:1002:FLC:OHB	2.21	0.44
1:C:86:LEU:HD22	1:C:87:PRO:HD2	1.98	0.44
1:C:96:ASP:OD2	1:C:192:HIS:ND1	2.46	0.44
1:C:175:ASP:OD1	1:C:177:SER:OG	2.35	0.44
1:A:201:ILE:HG12	1:A:325:ALA:HA	1.98	0.44
1:C:93:PHE:CE1	1:C:197:PRO:HA	2.52	0.44
1:C:154:VAL:HG22	1:C:200:LEU:HB2	1.99	0.44
1:A:336:LYS:HD2	1:A:338:VAL:HG13	2.00	0.43
1:A:191:VAL:HG23	1:A:193:LEU:HD11	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:115:ASN:N	1:C:133:ASP:OD2	2.41	0.43
1:C:133:ASP:CG	5:C:1101:HOH:O	2.61	0.42
1:A:65:ASP:OD2	1:A:67:ARG:CZ	2.68	0.42
1:A:122:SER:O	1:A:123:TYR:HB3	2.20	0.41
1:C:232:TYR:HB3	1:C:323:MET:CE	2.50	0.41
1:C:283:VAL:HG21	1:C:289:LEU:HD21	2.02	0.41
1:C:323:MET:HE1	1:C:329:LEU:HD22	2.00	0.41
1:A:154:VAL:HA	1:A:200:LEU:O	2.19	0.41
1:A:211:ASP:C	1:A:211:ASP:OD1	2.63	0.41
1:A:265:ARG:HG2	1:A:354:ILE:HD11	2.02	0.41
1:A:288:ILE:HD12	4:A:1004:GOL:H31	2.02	0.40
1:A:342:MET:HE2	1:A:342:MET:HB3	1.95	0.40
1:C:226:SER:HA	1:C:289:LEU:O	2.22	0.40
1:A:270:LEU:HB3	1:A:271:PRO:HD2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/312 (95%)	284 (96%)	10 (3%)	1 (0%)	36	58
1	C	303/312 (97%)	293 (97%)	7 (2%)	3 (1%)	12	28
All	All	598/624 (96%)	577 (96%)	17 (3%)	4 (1%)	18	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	181	ASP
1	C	133	ASP
1	C	59	ALA
1	C	217	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/266 (96%)	242 (95%)	12 (5%)	23	48
1	C	261/266 (98%)	251 (96%)	10 (4%)	29	56
All	All	515/532 (97%)	493 (96%)	22 (4%)	26	51

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	138	LEU
1	A	150	LEU
1	A	162	THR
1	A	177	SER
1	A	219	LYS
1	A	257	ASN
1	A	259	GLN
1	A	278	GLU
1	A	284	THR
1	A	301	THR
1	A	333	THR
1	A	356	VAL
1	C	55	THR
1	C	86	LEU
1	C	103	MET
1	C	177	SER
1	C	228	GLN
1	C	270	LEU
1	C	284	THR
1	C	301	THR
1	C	329	LEU
1	C	333	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	HIS
1	A	198	GLN
1	A	222	ASN
1	A	228	GLN
1	A	269	GLN
1	A	279	HIS
1	A	316	ASN
1	A	351	ASN
1	C	115	ASN
1	C	183	HIS
1	C	222	ASN
1	C	305	HIS
1	C	351	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1008	-	5,5,5	0.45	0	5,5,5	0.38	0
4	GOL	A	1009	-	5,5,5	0.32	0	5,5,5	0.26	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	C	1003	-	5,5,5	0.67	0	5,5,5	1.27	1 (20%)
4	GOL	A	1003	-	5,5,5	0.39	0	5,5,5	1.33	0
4	GOL	C	1004	-	5,5,5	0.60	0	5,5,5	0.54	0
3	FLC	C	1002	2	12,12,12	1.63	2 (16%)	17,17,17	2.66	7 (41%)
4	GOL	A	1006	-	5,5,5	0.73	0	5,5,5	1.61	2 (40%)
4	GOL	C	1005	-	5,5,5	0.30	0	5,5,5	0.46	0
4	GOL	A	1005	-	5,5,5	0.35	0	5,5,5	0.99	0
4	GOL	C	1008	-	5,5,5	0.81	0	5,5,5	1.16	0
4	GOL	A	1004	-	5,5,5	0.70	0	5,5,5	0.54	0
4	GOL	C	1006	-	5,5,5	0.80	0	5,5,5	0.77	0
4	GOL	C	1007	-	5,5,5	0.37	0	5,5,5	0.40	0
4	GOL	A	1007	-	5,5,5	0.87	0	5,5,5	1.40	1 (20%)
3	FLC	A	1002	2	12,12,12	0.96	0	17,17,17	1.35	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1008	-	-	2/4/4/4	-
4	GOL	A	1009	-	-	2/4/4/4	-
4	GOL	C	1003	-	-	1/4/4/4	-
4	GOL	A	1003	-	-	2/4/4/4	-
4	GOL	C	1004	-	-	2/4/4/4	-
3	FLC	C	1002	2	-	8/16/16/16	-
4	GOL	A	1006	-	-	2/4/4/4	-
4	GOL	C	1005	-	-	2/4/4/4	-
4	GOL	A	1005	-	-	0/4/4/4	-
4	GOL	C	1008	-	-	4/4/4/4	-
4	GOL	A	1004	-	-	4/4/4/4	-
4	GOL	C	1006	-	-	3/4/4/4	-
4	GOL	C	1007	-	-	1/4/4/4	-
4	GOL	A	1007	-	-	2/4/4/4	-
3	FLC	A	1002	2	-	3/16/16/16	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1002	FLC	CB-CBC	3.05	1.56	1.53
3	C	1002	FLC	CG-CGC	2.09	1.57	1.50

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1002	FLC	CG-CB-CBC	5.21	121.55	110.03
3	C	1002	FLC	OHB-CB-CBC	-4.96	101.92	108.96
3	C	1002	FLC	OB2-CBC-CB	4.63	122.03	113.14
3	C	1002	FLC	CB-CG-CGC	3.33	123.02	113.92
3	C	1002	FLC	OG2-CGC-OG1	-3.01	115.59	123.33
3	C	1002	FLC	OG1-CGC-CG	2.99	131.43	122.95
3	C	1002	FLC	OB2-CBC-OB1	-2.77	115.00	123.86
4	A	1007	GOL	O2-C2-C1	2.63	120.05	109.18
4	A	1006	GOL	O3-C3-C2	-2.60	98.68	110.38
3	A	1002	FLC	OG1-CGC-CG	-2.57	115.68	122.95
3	A	1002	FLC	CA-CB-CBC	2.26	115.03	110.03
4	A	1006	GOL	O2-C2-C3	-2.17	100.18	109.18
4	C	1003	GOL	O2-C2-C1	-2.15	100.26	109.18
3	A	1002	FLC	OB2-CBC-CB	2.15	117.27	113.14

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1002	FLC	CA-CB-CBC-OB1
3	C	1002	FLC	CA-CB-CBC-OB2
3	C	1002	FLC	OHB-CB-CBC-OB1
3	C	1002	FLC	OHB-CB-CBC-OB2
3	C	1002	FLC	CA-CB-CG-CGC
3	C	1002	FLC	CBC-CB-CG-CGC
3	C	1002	FLC	OHB-CB-CG-CGC
4	A	1003	GOL	C1-C2-C3-O3
4	A	1007	GOL	C1-C2-C3-O3
4	A	1008	GOL	O1-C1-C2-C3
4	C	1005	GOL	C1-C2-C3-O3
4	C	1008	GOL	O1-C1-C2-O2
4	C	1008	GOL	O2-C2-C3-O3
3	A	1002	FLC	CBC-CB-CG-CGC
3	A	1002	FLC	CA-CB-CG-CGC
4	A	1004	GOL	O1-C1-C2-C3
4	A	1004	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	1006	GOL	C1-C2-C3-O3
4	A	1009	GOL	C1-C2-C3-O3
4	C	1004	GOL	C1-C2-C3-O3
4	C	1006	GOL	O1-C1-C2-C3
4	C	1008	GOL	O1-C1-C2-C3
4	C	1008	GOL	C1-C2-C3-O3
4	A	1004	GOL	O2-C2-C3-O3
4	A	1007	GOL	O2-C2-C3-O3
4	A	1008	GOL	O1-C1-C2-O2
4	C	1005	GOL	O2-C2-C3-O3
4	A	1003	GOL	O2-C2-C3-O3
4	C	1003	GOL	O2-C2-C3-O3
3	A	1002	FLC	OHB-CB-CG-CGC
4	A	1004	GOL	O1-C1-C2-O2
4	A	1009	GOL	O2-C2-C3-O3
3	C	1002	FLC	CAC-CA-CB-OHB
4	A	1006	GOL	O2-C2-C3-O3
4	C	1006	GOL	C1-C2-C3-O3
4	C	1006	GOL	O1-C1-C2-O2
4	C	1007	GOL	O2-C2-C3-O3
4	C	1004	GOL	O2-C2-C3-O3


There are no ring outliers.


6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1008	GOL	1	0
4	A	1003	GOL	1	0
4	A	1006	GOL	2	0
4	A	1004	GOL	1	0
4	A	1007	GOL	2	0
3	A	1002	FLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	297/312 (95%)	-0.10	9 (3%)	52 47	12, 28, 56, 86	6 (2%)
1	C	304/312 (97%)	-0.12	7 (2%)	61 55	11, 28, 57, 87	7 (2%)
All	All	601/624 (96%)	-0.11	16 (2%)	56 50	11, 28, 56, 87	13 (2%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	356	VAL	5.6
1	C	356	VAL	4.8
1	A	127	ASP	4.3
1	A	193	LEU	3.3
1	A	103	MET	3.3
1	C	133	ASP	3.3
1	A	355	PRO	3.2
1	C	59	ALA	3.1
1	A	67	ARG	3.0
1	A	60	SER	3.0
1	C	193	LEU	2.9
1	C	53	PRO	2.8
1	C	56	PRO	2.5
1	C	355	PRO	2.2
1	A	266	GLN	2.0
1	A	142	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	C	1007	6/6	0.67	0.20	66,70,73,74	0
4	GOL	A	1007	6/6	0.81	0.17	35,40,41,46	0
4	GOL	A	1004	6/6	0.81	0.17	42,52,55,59	0
4	GOL	C	1008	6/6	0.82	0.26	52,57,60,63	0
4	GOL	A	1003	6/6	0.85	0.16	29,33,41,52	0
4	GOL	C	1006	6/6	0.86	0.16	45,54,55,55	0
3	FLC	C	1002	13/13	0.87	0.15	27,35,53,62	0
4	GOL	C	1003	6/6	0.87	0.16	25,35,39,40	0
4	GOL	C	1004	6/6	0.87	0.15	55,60,63,67	0
4	GOL	A	1009	6/6	0.89	0.20	59,61,64,74	0
4	GOL	A	1008	6/6	0.89	0.20	44,53,54,58	0
4	GOL	A	1005	6/6	0.92	0.13	40,47,50,54	0
4	GOL	A	1006	6/6	0.92	0.11	29,33,38,38	0
4	GOL	C	1005	6/6	0.93	0.11	35,36,37,39	0
3	FLC	A	1002	13/13	0.97	0.07	27,30,35,36	0
2	CU	C	1001	1/1	0.99	0.02	35,35,35,35	0
2	CU	A	1001	1/1	1.00	0.02	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FLC C 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FLC A 1002:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CU C 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.