



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 02:52 AM UTC

PDB ID : 4AMI / pdb_00004ami
Title : Turkey beta1 adrenergic receptor with stabilising mutations and bound biased agonist bucindolol
Authors : Warne, T.; Edwards, P.C.; Leslie, A.G.; Tate, C.G.
Deposited on : 2012-03-11
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	
1	B	315	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4673 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-1 ADRENERGIC RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2237	1479	365	374	19			
1	B	288	Total	C	N	O	S	0	0	0
			2292	1513	378	381	20			

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	31	MET	-	expression tag	UNP P07700
A	32	GLY	-	expression tag	UNP P07700
A	369	HIS	-	expression tag	UNP P07700
A	370	HIS	-	expression tag	UNP P07700
A	371	HIS	-	expression tag	UNP P07700
A	372	HIS	-	expression tag	UNP P07700
A	373	HIS	-	expression tag	UNP P07700
A	68	SER	ARG	engineered mutation	UNP P07700
A	90	VAL	MET	engineered mutation	UNP P07700
A	116	LEU	CYS	engineered mutation	UNP P07700
A	227	ALA	TYR	engineered mutation	UNP P07700
A	282	LEU	ALA	engineered mutation	UNP P07700
A	327	ALA	PHE	engineered mutation	UNP P07700
A	338	MET	PHE	engineered mutation	UNP P07700
A	358	ALA	CYS	engineered mutation	UNP P07700
B	31	MET	-	expression tag	UNP P07700
B	32	GLY	-	expression tag	UNP P07700
B	369	HIS	-	expression tag	UNP P07700
B	370	HIS	-	expression tag	UNP P07700
B	371	HIS	-	expression tag	UNP P07700
B	372	HIS	-	expression tag	UNP P07700
B	373	HIS	-	expression tag	UNP P07700
B	68	SER	ARG	engineered mutation	UNP P07700
B	90	VAL	MET	engineered mutation	UNP P07700
B	116	LEU	CYS	engineered mutation	UNP P07700

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	227	ALA	TYR	engineered mutation	UNP P07700
B	282	LEU	ALA	engineered mutation	UNP P07700
B	327	ALA	PHE	engineered mutation	UNP P07700
B	338	MET	PHE	engineered mutation	UNP P07700
B	358	ALA	CYS	engineered mutation	UNP P07700

- Molecule 2 is 2-[(2S)-3-[[1-(1H-indol-3-yl)-2-methyl-propan-2-yl]amino]-2-oxidanyl-propoxy]benzenecarbonitrile (CCD ID: G90) (formula: $C_{22}H_{25}N_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			27	22	3	2		
2	B	1	Total	C	N	O	0	0
			27	22	3	2		

- Molecule 3 is HEGA-10 (CCD ID: 2CV) (formula: $C_{18}H_{37}NO_7$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	14	1	2		
3	B	1	Total	C	N	O	0	0
			15	12	1	2		
3	B	1	Total	C	N	O	0	0
			14	12	1	1		
3	B	1	Total	C	N	O	0	0
			23	17	1	5		
3	B	1	Total	C			0	0
			8	8				


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	5	Total	O	0	0
			5	5		
4	B	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: BETA-1 ADRENERGIC RECEPTOR



• Molecule 1: BETA-1 ADRENERGIC RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	89.84Å 60.68Å 107.82Å 90.00° 110.78° 90.00°	Depositor
Resolution (Å)	44.82 – 3.20 44.82 – 3.20	Depositor EDS
% Data completeness (in resolution range)	95.7 (44.82-3.20) 95.7 (44.82-3.20)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.7.0019	Depositor
R, R_{free}	0.242 , 0.279 0.241 , 0.272	Depositor DCC
R_{free} test set	898 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	100.6	Xtriage
Anisotropy	0.238	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4673	wwPDB-VP
Average B, all atoms (Å ²)	93.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 2CV, G90

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/2291	0.81	0/3125
1	B	0.40	0/2346	0.80	0/3196
All	All	0.41	0/4637	0.81	0/6321

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2237	0	2313	21	0
1	B	2292	0	2381	19	0
2	A	27	0	25	4	0
2	B	27	0	25	2	0
3	A	17	0	26	1	0
3	B	60	0	87	2	0
4	A	5	0	0	0	0
4	B	8	0	0	0	0
All	All	4673	0	4857	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:172:VAL:HG11	1:B:214:ILE:HG21	1.67	0.75
1:B:172:VAL:HG11	1:B:214:ILE:CG2	2.26	0.66
1:A:125:VAL:CG1	2:A:1359:G90:H11	2.32	0.60
1:A:164:THR:HG23	3:B:1361:2CV:H151	1.83	0.60
1:B:290:LYS:O	1:B:294:ILE:HG12	2.03	0.59
1:A:175:LEU:HD13	1:B:175:LEU:HD21	1.85	0.58
1:A:121:ASP:OD2	2:A:1359:G90:H32C	2.05	0.56
1:A:118:THR:HG21	1:A:201:PHE:CE1	2.42	0.54
1:A:285:GLU:O	1:A:289:LEU:HD23	2.08	0.53
1:A:125:VAL:HG11	2:A:1359:G90:H11	1.89	0.52
1:A:306:PHE:CE2	2:A:1359:G90:H52C	2.44	0.52
1:A:320:VAL:HG13	1:A:324:LEU:HD23	1.91	0.52
1:B:321:PRO:HB2	1:B:323:TRP:CD1	2.45	0.52
1:B:101:LEU:HD12	1:B:107:TRP:HE3	1.75	0.51
1:B:75:LEU:HD13	1:B:153:MET:HG3	1.93	0.51
1:B:125:VAL:CG1	2:B:1360:G90:H11	2.41	0.51
1:A:33:ALA:HA	1:A:36:LEU:HD12	1.94	0.50
1:B:143:ILE:HD11	1:B:289:LEU:HD11	1.94	0.49
1:A:139:ARG:HA	1:A:139:ARG:NE	2.27	0.48
1:B:139:ARG:NE	1:B:139:ARG:HA	2.28	0.48
1:B:123:LEU:HD22	1:B:174:PHE:HE1	1.79	0.48
1:A:75:LEU:HD13	1:A:153:MET:HG3	1.95	0.47
1:B:177:ILE:HA	1:B:182:TRP:CD1	2.50	0.47
1:A:213:ILE:HA	1:A:217:TYR:HB2	1.98	0.46
1:B:106:THR:HG22	1:B:197:GLY:O	2.16	0.46
3:A:1360:2CV:H151	1:B:164:THR:HG23	1.98	0.45
1:A:177:ILE:HA	1:A:182:TRP:CD1	2.52	0.45
1:A:118:THR:HG21	1:A:201:PHE:CD1	2.52	0.44
1:B:168:ILE:O	1:B:172:VAL:HG23	2.17	0.44
1:B:306:PHE:CE2	2:B:1360:G90:H52C	2.53	0.44
1:B:209:ILE:HD11	1:B:314:VAL:HG11	2.00	0.44
1:B:213:ILE:HA	1:B:217:TYR:HB2	2.00	0.44
1:A:356:LEU:C	1:A:358:ALA:H	2.26	0.43
1:B:290:LYS:HB3	1:B:345:ARG:HH22	1.83	0.43
1:A:214:ILE:HG12	3:B:1358:2CV:H271	1.98	0.43
1:A:209:ILE:HD11	1:A:314:VAL:HG11	2.01	0.42
1:A:109:TRP:HB3	1:A:113:LEU:HD12	2.01	0.42
1:A:115:GLU:OE2	1:A:180:HIS:NE2	2.53	0.42
1:A:205:ARG:NH1	1:A:314:VAL:O	2.52	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	278/315 (88%)	268 (96%)	9 (3%)	1 (0%)	30	62
1	B	284/315 (90%)	277 (98%)	7 (2%)	0	100	100
All	All	562/630 (89%)	545 (97%)	16 (3%)	1 (0%)	43	73

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	357	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/273 (89%)	242 (100%)	1 (0%)	84	86
1	B	250/273 (92%)	244 (98%)	6 (2%)	43	70
All	All	493/546 (90%)	486 (99%)	7 (1%)	59	77

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	329	ASN
1	B	106	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	171	LEU
1	B	172	VAL
1	B	175	LEU
1	B	190	LEU
1	B	329	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	ASN
1	B	59	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	G90	A	1359	-	28,29,29	2.71	5 (17%)	38,40,40	1.88	9 (23%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	2CV	A	1360	-	16,16,25	0.64	0	17,17,30	0.73	1 (5%)
3	2CV	B	1358	-	14,14,25	0.70	0	14,14,30	0.78	0
3	2CV	B	1361	-	22,22,25	0.80	0	25,26,30	0.70	0
3	2CV	B	1362	-	7,7,25	0.44	0	6,6,30	0.43	0
2	G90	B	1360	-	28,29,29	2.73	5 (17%)	38,40,40	1.60	7 (18%)
3	2CV	B	1359	-	13,13,25	0.66	0	14,14,30	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G90	A	1359	-	-	10/18/18/18	0/3/3/3
3	2CV	A	1360	-	-	5/18/18/34	-
3	2CV	B	1358	-	-	7/13/13/34	-
3	2CV	B	1361	-	-	12/28/28/34	-
3	2CV	B	1362	-	-	4/5/5/34	-
2	G90	B	1360	-	-	6/18/18/18	0/3/3/3
3	2CV	B	1359	-	-	4/13/13/34	-

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1360	G90	C7-C12	-11.45	1.27	1.44
2	A	1359	G90	C7-C12	-10.71	1.29	1.44
2	A	1359	G90	CD2-CE2	5.62	1.48	1.41
2	B	1360	G90	CD2-CE2	5.23	1.47	1.41
2	A	1359	G90	C7-C6	4.63	1.49	1.40
2	A	1359	G90	CA-N	-4.26	1.45	1.49
2	B	1360	G90	C7-C6	4.26	1.49	1.40
2	B	1360	G90	CA-N	-3.59	1.45	1.49
2	A	1359	G90	CD2-CG	3.00	1.48	1.44
2	B	1360	G90	CD2-CG	2.84	1.48	1.44

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1359	G90	C3-N-CA	7.16	124.04	116.35
2	B	1360	G90	C3-N-CA	5.46	122.21	116.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1359	G90	CZ2-CE2-CD2	-4.07	118.25	122.19
2	B	1360	G90	CZ2-CE2-CD2	-3.79	118.52	122.19
2	A	1359	G90	C1-CA-CB	2.95	113.86	109.91
2	A	1359	G90	CE3-CD2-CE2	2.77	121.71	118.84
2	B	1360	G90	C5-O2-C6	2.64	124.08	118.21
2	B	1360	G90	CE2-NE1-CD1	2.55	111.35	109.08
2	A	1359	G90	CZ2-CE2-NE1	2.49	134.73	130.74
2	B	1360	G90	CE3-CD2-CE2	2.44	121.37	118.84
2	B	1360	G90	CZ2-CE2-NE1	2.42	134.62	130.74
2	B	1360	G90	O2-C6-C7	2.40	121.43	115.80
2	A	1359	G90	O2-C6-C7	2.32	121.25	115.80
2	A	1359	G90	CE2-NE1-CD1	2.27	111.09	109.08
3	A	1360	2CV	C35-N33-C36	2.16	117.37	115.23
2	A	1359	G90	C1-CA-N	-2.09	105.02	109.91
2	A	1359	G90	O2-C6-C11	-2.04	119.51	123.95

There are no chirality outliers.

All (48) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1359	G90	N-C3-C4-C5
2	A	1359	G90	C1-CA-N-C3
2	A	1359	G90	C2-CA-N-C3
2	A	1359	G90	CB-CA-N-C3
2	A	1359	G90	N-CA-CB-CG
2	B	1360	G90	C3-C4-C5-O2
3	A	1360	2CV	N33-C35-C60-O63
3	B	1358	2CV	N33-C35-C60-O63
3	B	1361	2CV	N33-C35-C60-O63
3	B	1361	2CV	N33-C36-C37-C40
3	B	1361	2CV	N33-C36-C37-O47
3	B	1361	2CV	O49-C40-C41-O51
3	B	1361	2CV	C21-C24-C27-C30
3	B	1358	2CV	C27-C30-N33-C35
2	A	1359	G90	O1-C4-C5-O2
3	B	1358	2CV	O34-C30-N33-C35
3	B	1361	2CV	O49-C40-C41-C42
2	A	1359	G90	C3-C4-C5-O2
2	A	1359	G90	N-C3-C4-O1
3	A	1360	2CV	C21-C24-C27-C30
3	B	1359	2CV	C12-C15-C18-C21
3	B	1358	2CV	C15-C18-C21-C24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	B	1358	2CV	C9-C12-C15-C18
3	B	1362	2CV	C12-C15-C18-C21
3	B	1359	2CV	C18-C21-C24-C27
3	B	1361	2CV	C37-C40-C41-C42
2	B	1360	G90	C11-C6-O2-C5
3	A	1360	2CV	C9-C12-C15-C18
3	B	1359	2CV	C21-C24-C27-C30
2	B	1360	G90	C7-C6-O2-C5
2	A	1359	G90	C11-C6-O2-C5
3	B	1361	2CV	C12-C15-C18-C21
3	B	1362	2CV	C18-C21-C24-C27
3	A	1360	2CV	C18-C21-C24-C27
3	A	1360	2CV	C12-C15-C18-C21
2	A	1359	G90	C7-C6-O2-C5
3	B	1361	2CV	C18-C21-C24-C27
2	B	1360	G90	O1-C4-C5-O2
3	B	1358	2CV	C21-C24-C27-C30
3	B	1361	2CV	C37-C40-C41-O51
3	B	1361	2CV	C15-C18-C21-C24
3	B	1362	2CV	C21-C24-C27-C30
3	B	1362	2CV	C9-C12-C15-C18
3	B	1361	2CV	C9-C12-C15-C18
3	B	1359	2CV	C9-C12-C15-C18
3	B	1358	2CV	C15-C12-C9-C1
2	B	1360	G90	N1-C12-C7-C6
2	B	1360	G90	N-CA-CB-CG



There are no ring outliers.



5 monomers are involved in 9 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1359	G90	4	0
3	A	1360	2CV	1	0
3	B	1358	2CV	1	0
3	B	1361	2CV	1	0
2	B	1360	G90	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/315 (89%)	0.55	25 (8%) 15 10	55, 92, 136, 155	0
1	B	288/315 (91%)	0.57	31 (10%) 11 7	56, 89, 132, 150	0
All	All	570/630 (90%)	0.56	56 (9%) 13 9	55, 90, 135, 155	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	278	SER	8.2
1	B	280	VAL	7.8
1	A	318	ASP	6.7
1	A	147	PHE	6.2
1	B	238	ILE	6.2
1	A	284	ARG	5.4
1	B	237	GLN	5.0
1	B	282	LEU	4.9
1	B	71	ARG	4.1
1	B	46	LEU	3.9
1	A	36	LEU	3.8
1	B	236	GLU	3.7
1	A	92	LEU	3.6
1	B	86	ALA	3.5
1	B	187	PRO	3.3
1	A	71	ARG	3.2
1	B	357	LEU	3.2
1	B	61	LEU	3.1
1	A	124	CYS	3.1
1	A	358	ALA	3.1
1	B	279	ARG	2.9
1	A	317	ARG	2.8
1	B	57	ALA	2.8
1	A	35	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	209	ILE	2.7
1	B	323	TRP	2.7
1	B	200	ASP	2.7
1	B	239	ARG	2.6
1	B	111	SER	2.6
1	A	283	MET	2.6
1	B	88	LEU	2.5
1	A	282	LEU	2.5
1	A	322	ASP	2.5
1	B	47	LEU	2.4
1	A	86	ALA	2.4
1	A	39	GLN	2.4
1	B	72	LEU	2.4
1	B	42	ALA	2.3
1	B	284	ARG	2.3
1	B	35	LEU	2.2
1	A	178	MET	2.2
1	A	316	ASN	2.2
1	B	240	LYS	2.2
1	B	179	MET	2.2
1	A	152	LEU	2.2
1	B	33	ALA	2.2
1	A	323	TRP	2.2
1	A	125	VAL	2.1
1	B	106	THR	2.1
1	A	349	PHE	2.1
1	A	310	ASN	2.1
1	B	356	LEU	2.1
1	A	348	ASP	2.0
1	B	43	GLY	2.0
1	B	281	MET	2.0
1	A	148	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	2CV	B	1362	8/26	0.82	0.44	71,73,78,80	0
3	2CV	B	1359	14/26	0.86	0.22	79,98,120,123	0
3	2CV	A	1360	17/26	0.88	0.17	62,103,123,125	0
3	2CV	B	1361	23/26	0.92	0.17	67,125,143,148	0
3	2CV	B	1358	15/26	0.92	0.21	78,87,96,101	0
2	G90	A	1359	27/27	0.93	0.10	66,72,74,74	0
2	G90	B	1360	27/27	0.95	0.10	63,69,75,75	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around 2CV B 1359:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV A 1360:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2CV B 1361:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 2CV B 1358:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around G90 A 1359:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around G90 B 1360:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.