



## wwPDB EM Validation Summary Report ⓘ

Mar 14, 2026 – 01:13 AM UTC

PDB ID : 7AML / pdb\_00007aml  
EMDB ID : EMD-11822  
Title : RET/GDNF/GFRa1 extracellular complex Cryo-EM structure  
Authors : Adams, S.E.; Earl, C.P.; Purkiss, A.G.; McDonald, N.Q.  
Deposited on : 2020-10-09  
Resolution : 3.50 Å (reported)  
Based on initial model : 7AB8

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	13950 ( 3.00 - 4.00 )

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	615	
1	D	615	
2	B	353	
2	E	353	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	C	101	 71% 26%
3	F	101	 71% 26%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 31456 atoms, of which 15438 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase receptor Ret.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	A	596	9267	2963	4546	807	923	28	0	0
1	D	596	9267	2963	4546	807	923	28	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	505	ARG	LYS	conflict	UNP A8E7C6
A	627	GLY	-	expression tag	UNP A8E7C6
A	628	SER	-	expression tag	UNP A8E7C6
A	629	GLU	-	expression tag	UNP A8E7C6
A	630	PHE	-	expression tag	UNP A8E7C6
A	631	GLU	-	expression tag	UNP A8E7C6
A	632	ASN	-	expression tag	UNP A8E7C6
A	633	LEU	-	expression tag	UNP A8E7C6
A	634	TYR	-	expression tag	UNP A8E7C6
A	635	PHE	-	expression tag	UNP A8E7C6
A	636	GLN	-	expression tag	UNP A8E7C6
D	505	ARG	LYS	conflict	UNP A8E7C6
D	627	GLY	-	expression tag	UNP A8E7C6
D	628	SER	-	expression tag	UNP A8E7C6
D	629	GLU	-	expression tag	UNP A8E7C6
D	630	PHE	-	expression tag	UNP A8E7C6
D	631	GLU	-	expression tag	UNP A8E7C6
D	632	ASN	-	expression tag	UNP A8E7C6
D	633	LEU	-	expression tag	UNP A8E7C6
D	634	TYR	-	expression tag	UNP A8E7C6
D	635	PHE	-	expression tag	UNP A8E7C6
D	636	GLN	-	expression tag	UNP A8E7C6

- Molecule 2 is a protein called GDNF family receptor alpha.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
2	B	304	4719	1478	2334	427	448	32	0	0
2	E	304	4719	1478	2334	427	448	32	0	0

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	352	GLY	-	expression tag	UNP Q98TT9
B	353	SER	-	expression tag	UNP Q98TT9
B	354	GLU	-	expression tag	UNP Q98TT9
B	355	PHE	-	expression tag	UNP Q98TT9
B	356	LEU	-	expression tag	UNP Q98TT9
B	357	GLU	-	expression tag	UNP Q98TT9
B	358	VAL	-	expression tag	UNP Q98TT9
B	359	LEU	-	expression tag	UNP Q98TT9
B	360	PHE	-	expression tag	UNP Q98TT9
B	361	GLN	-	expression tag	UNP Q98TT9
B	362	GLY	-	expression tag	UNP Q98TT9
B	363	PRO	-	expression tag	UNP Q98TT9
B	364	GLY	-	expression tag	UNP Q98TT9
B	365	GLY	-	expression tag	UNP Q98TT9
B	366	GLY	-	expression tag	UNP Q98TT9
B	367	GLU	-	expression tag	UNP Q98TT9
B	368	ASN	-	expression tag	UNP Q98TT9
B	369	LEU	-	expression tag	UNP Q98TT9
B	370	TYR	-	expression tag	UNP Q98TT9
B	371	PHE	-	expression tag	UNP Q98TT9
B	372	GLN	-	expression tag	UNP Q98TT9
E	352	GLY	-	expression tag	UNP Q98TT9
E	353	SER	-	expression tag	UNP Q98TT9
E	354	GLU	-	expression tag	UNP Q98TT9
E	355	PHE	-	expression tag	UNP Q98TT9
E	356	LEU	-	expression tag	UNP Q98TT9
E	357	GLU	-	expression tag	UNP Q98TT9
E	358	VAL	-	expression tag	UNP Q98TT9
E	359	LEU	-	expression tag	UNP Q98TT9
E	360	PHE	-	expression tag	UNP Q98TT9
E	361	GLN	-	expression tag	UNP Q98TT9
E	362	GLY	-	expression tag	UNP Q98TT9
E	363	PRO	-	expression tag	UNP Q98TT9
E	364	GLY	-	expression tag	UNP Q98TT9
E	365	GLY	-	expression tag	UNP Q98TT9

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	366	GLY	-	expression tag	UNP Q98TT9
E	367	GLU	-	expression tag	UNP Q98TT9
E	368	ASN	-	expression tag	UNP Q98TT9
E	369	LEU	-	expression tag	UNP Q98TT9
E	370	TYR	-	expression tag	UNP Q98TT9
E	371	PHE	-	expression tag	UNP Q98TT9
E	372	GLN	-	expression tag	UNP Q98TT9


- Molecule 3 is a protein called Glial cell line-derived neurotrophic factor.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	C	98	Total	C	H	N	O	S	0	0
			1550	488	763	139	153	7		
3	F	98	Total	C	H	N	O	S	0	0
			1550	488	763	139	153	7		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
			Total	Ca	
4	A	4	Total	Ca	0
			4	4	
4	D	4	Total	Ca	0
			4	4	

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).




Mol	Chain	Residues	Atoms					AltConf
5	A	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	A	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	A	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	A	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	A	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	A	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	A	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	D	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	D	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	D	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	D	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	D	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	D	1	Total	C	H	N	O	0
			23	8	9	1	5	
5	C	1	Total	C	H	N	O	0
			24	8	10	1	5	
5	F	1	Total	C	H	N	O	0
			24	8	10	1	5	


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase receptor Ret




- Molecule 1: Proto-oncogene tyrosine-protein kinase receptor Ret







• Molecule 2: GDNF family receptor alpha



• Molecule 2: GDNF family receptor alpha





• Molecule 3: Glial cell line-derived neurotrophic factor



• Molecule 3: Glial cell line-derived neurotrophic factor



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	382547	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	48.6	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	46296	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	3.552	Depositor
Minimum map value	-2.321	Depositor
Average map value	0.003	Depositor
Map value standard deviation	0.049	Depositor
Recommended contour level	0.269	Depositor
Map size ( $\text{\AA}$ )	345.6, 345.6, 345.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.08, 1.08, 1.08	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.23	1/4823 (0.0%)	0.45	1/6550 (0.0%)
1	D	0.26	1/4823 (0.0%)	0.46	2/6550 (0.0%)
2	B	0.24	0/2429	0.45	1/3271 (0.0%)
2	E	0.24	0/2429	0.46	1/3271 (0.0%)
3	C	0.35	0/801	0.79	0/1077
3	F	0.50	1/801 (0.1%)	0.89	0/1077
All	All	0.27	3/16106 (0.0%)	0.51	5/21796 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	206	ILE	N-CA	-5.50	1.39	1.46
1	D	502	ASP	N-CA	-5.07	1.39	1.46
1	A	502	ASP	N-CA	-5.02	1.39	1.46

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	222	SER	N-CA-C	-6.66	98.35	109.07
2	E	222	SER	N-CA-C	-6.30	98.92	109.07
1	D	562	ASP	N-CA-C	5.73	118.01	111.02
1	A	142	SER	N-CA-C	-5.42	101.15	109.76
1	D	142	SER	N-CA-C	-5.20	102.58	110.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4721	4546	4555	102	0
1	D	4721	4546	4555	96	0
2	B	2385	2334	2334	54	0
2	E	2385	2334	2334	52	0
3	C	787	763	763	28	0
3	F	787	763	763	26	0
4	A	4	0	0	0	0
4	D	4	0	0	1	0
5	A	98	66	91	6	0
5	C	14	10	13	0	0
5	D	98	66	91	6	0
5	F	14	10	13	0	0
All	All	16018	15438	15512	354	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 354 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:274:GLU:N	1:A:346:ASN:OD1	2.04	0.91
1:D:274:GLU:N	1:D:346:ASN:OD1	2.04	0.90
2:E:269:ASN:OD1	2:E:291:SER:OG	1.94	0.86
2:B:269:ASN:OD1	2:B:291:SER:OG	1.94	0.84
2:B:151:ASN:O	2:B:155:ASN:ND2	2.13	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	594/615 (97%)	559 (94%)	35 (6%)	0	100	100
1	D	594/615 (97%)	559 (94%)	35 (6%)	0	100	100
2	B	298/353 (84%)	279 (94%)	19 (6%)	0	100	100
2	E	298/353 (84%)	278 (93%)	20 (7%)	0	100	100
3	C	96/101 (95%)	96 (100%)	0	0	100	100
3	F	96/101 (95%)	90 (94%)	6 (6%)	0	100	100
All	All	1976/2138 (92%)	1861 (94%)	115 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	535/553 (97%)	529 (99%)	6 (1%)	65	74
1	D	535/553 (97%)	528 (99%)	7 (1%)	61	72
2	B	272/312 (87%)	272 (100%)	0	100	100
2	E	272/312 (87%)	272 (100%)	0	100	100
3	C	90/92 (98%)	90 (100%)	0	100	100
3	F	90/92 (98%)	88 (98%)	2 (2%)	45	65
All	All	1794/1914 (94%)	1779 (99%)	15 (1%)	70	77

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	185	ASN
3	F	142	CYS
1	D	259	ASN
3	F	169	CYS
1	D	572	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	296	GLN
1	D	295	ASN
1	D	296	GLN
2	B	245	ASN
2	E	245	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 8 are monoatomic - leaving 16 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	NAG	D	711	1	14,14,15	0.25	0	17,19,21	0.40	0
5	NAG	A	707	1	14,14,15	0.38	0	17,19,21	0.53	0
5	NAG	A	711	1	14,14,15	0.26	0	17,19,21	0.39	0
5	NAG	D	706	1	14,14,15	0.24	0	17,19,21	0.56	0
5	NAG	D	707	1	14,14,15	0.39	0	17,19,21	0.53	0
5	NAG	D	709	1	14,14,15	0.29	0	17,19,21	0.36	0
5	NAG	D	710	1	14,14,15	0.25	0	17,19,21	0.39	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	NAG	A	708	1	14,14,15	0.27	0	17,19,21	0.35	0
5	NAG	D	708	1	14,14,15	0.27	0	17,19,21	0.35	0
5	NAG	A	705	1	14,14,15	0.25	0	17,19,21	0.40	0
5	NAG	C	301	3	14,14,15	0.23	0	17,19,21	0.44	0
5	NAG	F	301	3	14,14,15	0.23	0	17,19,21	0.45	0
5	NAG	A	706	1	14,14,15	0.24	0	17,19,21	0.56	0
5	NAG	D	705	1	14,14,15	0.25	0	17,19,21	0.40	0
5	NAG	A	710	1	14,14,15	0.25	0	17,19,21	0.39	0
5	NAG	A	709	1	14,14,15	0.29	0	17,19,21	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	D	711	1	-	4/6/23/26	0/1/1/1
5	NAG	A	707	1	-	2/6/23/26	0/1/1/1
5	NAG	A	711	1	-	4/6/23/26	0/1/1/1
5	NAG	D	706	1	-	1/6/23/26	0/1/1/1
5	NAG	D	707	1	-	2/6/23/26	0/1/1/1
5	NAG	D	709	1	-	2/6/23/26	0/1/1/1
5	NAG	D	710	1	-	1/6/23/26	0/1/1/1
5	NAG	A	708	1	-	1/6/23/26	0/1/1/1
5	NAG	D	708	1	-	1/6/23/26	0/1/1/1
5	NAG	A	705	1	-	3/6/23/26	0/1/1/1
5	NAG	C	301	3	-	2/6/23/26	0/1/1/1
5	NAG	F	301	3	-	2/6/23/26	0/1/1/1
5	NAG	A	706	1	-	1/6/23/26	0/1/1/1
5	NAG	D	705	1	-	3/6/23/26	0/1/1/1
5	NAG	A	710	1	-	1/6/23/26	0/1/1/1
5	NAG	A	709	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	301	NAG	C4-C5-C6-O6
5	F	301	NAG	C4-C5-C6-O6
5	C	301	NAG	O5-C5-C6-O6
5	F	301	NAG	O5-C5-C6-O6
5	A	705	NAG	C8-C7-N2-C2

There are no ring outliers.

10 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	D	711	NAG	1	0
5	A	711	NAG	1	0
5	D	706	NAG	1	0
5	D	709	NAG	1	0
5	D	710	NAG	2	0
5	A	708	NAG	1	0
5	D	708	NAG	1	0
5	A	706	NAG	1	0
5	A	710	NAG	1	0
5	A	709	NAG	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11822. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


### 6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 131




Y Index: 138




Z Index: 145


### 6.3.2 Raw map



X Index: 131



Y Index: 138




Z Index: 144


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.269. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_11822\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 165 nm<sup>3</sup>; this corresponds to an approximate mass of 149 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.286 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.286 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	4.12	4.90	4.18
Unmasked-calculated*	4.09	4.88	4.15


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from author-provided FSC intersecting FSC 0.143 CUT-OFF 4.12 differs from the reported value 3.5 by more than 10 %

The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.09 differs from the reported value 3.5 by more than 10 %

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-11822 and PDB model 7AML. Per-residue inclusion information can be found in section [3](#) on page [8](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.269 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.269).















## 9.4 Atom inclusion [i](#)



At the recommended contour level, 74% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.269) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7410	 0.3300
A	 0.7150	 0.3240
B	 0.7540	 0.3370
C	 0.8780	 0.3760
D	 0.7130	 0.3230
E	 0.7420	 0.3230
F	 0.8620	 0.3570

