



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:17 AM UTC

PDB ID : 8AM0 / pdb_00008am0
Title : Crystal structure of human T1061E PI3Kalpha in complex with its regulatory subunit and the inhibitor GDC-0077 (Inavolisib)
Authors : Goncalves, M.; Johnson, J.L.; Roewer, K.M.
Deposited on : 2022-08-02
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4591 (2.84-2.80)
Clashscore	190562	5010 (2.84-2.80)
Ramachandran outliers	187476	4916 (2.84-2.80)
Sidechain outliers	187428	4918 (2.84-2.80)
RSRZ outliers	180081	4594 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1068	10% 95% • •
2	B	293	12% 86% • 13%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 10845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1031	Total	C	N	O	S	0	0	0
			8443	5401	1447	1526	69			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1061	GLU	THR	engineered mutation	UNP P42336


- Molecule 2 is a protein called Isoform 3 of Phosphatidylinositol 3-kinase regulatory subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	255	Total	C	N	O	S	0	0	0
			2161	1353	385	418	5			

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	301	MET	-	initiating methionine	UNP P27986-3
B	302	HIS	-	expression tag	UNP P27986-3
B	303	HIS	-	expression tag	UNP P27986-3
B	304	HIS	-	expression tag	UNP P27986-3
B	305	HIS	-	expression tag	UNP P27986-3
B	306	HIS	-	expression tag	UNP P27986-3
B	307	HIS	-	expression tag	UNP P27986-3

- Molecule 3 is (2R)-2-[[2-[(4S)-4-[bis(fluoranyl)methyl]-2-oxidanylidene-1,3-oxazolidin-3-yl]-5,6-dihydroimidazo[1,2-d][1,4]benzoxazepin-9-yl]amino]propanamide (CCD ID: MWF) (formula: C₁₈H₁₉F₂N₅O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			29	18	2	5	4		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	169	Total	O	0	0
			169	169		
7	B	20	Total	O	0	0
			20	20		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphatidylinositol 4,5-bisphosphate 3-kinase catalytic subunit alpha isoform



- Molecule 2: Isoform 3 of Phosphatidylinositol 3-kinase regulatory subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	99.81Å 107.68Å 136.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.39 – 2.82 47.39 – 2.82	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.39-2.82) 100.0 (47.39-2.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.33 (at 2.81Å)	Xtriage
Refinement program	BUSTER 2.11.8 (8-JUN-2022)	Depositor
R, R_{free}	0.285 , 0.300 0.272 , 0.291	Depositor DCC
R_{free} test set	1768 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	59.8	Xtriage
Anisotropy	0.308	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10845	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, MWF, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	1/8635 (0.0%)	0.88	0/11667
2	B	0.70	0/2195	0.91	0/2939
All	All	0.72	1/10830 (0.0%)	0.88	0/14606

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	393	ILE	CA-CB	8.63	1.58	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8443	0	8462	7	0
2	B	2161	0	2129	1	0
3	A	29	0	0	0	0
4	A	8	0	12	0	0
4	B	4	0	6	0	0
5	A	10	0	14	0	0
6	A	1	0	0	0	0
7	A	169	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	20	0	0	0	0
All	All	10845	0	10623	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:PRO:N	1:A:127:GLU:OE1	2.34	0.60
1:A:917:HIS:NE2	1:A:920:ASN:ND2	2.62	0.48
1:A:901:CYS:HG	1:A:954:PHE:HE2	1.61	0.46
2:B:419:LYS:HD2	2:B:419:LYS:O	2.17	0.44
1:A:142:ASN:O	1:A:145:ASN:OD1	2.38	0.42
1:A:875:HIS:ND1	1:A:962:ILE:HG21	2.36	0.41
1:A:140:ARG:NH2	1:A:307:TYR:CE2	2.89	0.40
1:A:375:ARG:NH1	1:A:419:HIS:HE1	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1023/1068 (96%)	980 (96%)	43 (4%)	0	100	100
2	B	249/293 (85%)	246 (99%)	3 (1%)	0	100	100
All	All	1272/1361 (94%)	1226 (96%)	46 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	946/974 (97%)	942 (100%)	4 (0%)	84	94
2	B	235/272 (86%)	231 (98%)	4 (2%)	53	81
All	All	1181/1246 (95%)	1173 (99%)	8 (1%)	76	91

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	196	VAL
1	A	209	LEU
1	A	251	ILE
1	A	1062	ILE
2	B	419	LYS
2	B	422	VAL
2	B	431	TYR
2	B	498	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	75	GLN
1	A	107	ASN
1	A	284	ASN
1	A	759	HIS
1	A	822	ASN
2	B	417	ASN
2	B	552	GLN
2	B	572	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	PGE	A	2004	-	9,9,9	0.12	0	8,8,8	0.10	0
4	EDO	B	601	-	3,3,3	0.24	0	2,2,2	0.31	0
4	EDO	A	2003	-	3,3,3	0.24	0	2,2,2	0.29	0
4	EDO	A	2002	-	3,3,3	0.24	0	2,2,2	0.30	0
3	MWF	A	2001	-	30,32,32	0.96	3 (10%)	36,47,47	1.27	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PGE	A	2004	-	-	4/7/7/7	-
4	EDO	B	601	-	-	1/1/1/1	-
4	EDO	A	2003	-	-	1/1/1/1	-
4	EDO	A	2002	-	-	1/1/1/1	-
3	MWF	A	2001	-	-	5/13/39/39	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2001	MWF	F1-C18	-2.13	1.33	1.36
3	A	2001	MWF	C13-C14	-2.09	1.48	1.51
3	A	2001	MWF	F2-C18	-2.08	1.33	1.36

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2001	MWF	C4-N2-C2	3.85	129.22	122.60
3	A	2001	MWF	F2-C18-C15	2.74	112.52	109.29
3	A	2001	MWF	C12-C11-N3	-2.66	111.02	112.77
3	A	2001	MWF	F1-C18-C15	2.64	112.40	109.29

There are no chirality outliers.

All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	2001	MWF	C1-C2-C3-N1
3	A	2001	MWF	C1-C2-C3-O1
5	A	2004	PGE	O2-C3-C4-O3
5	A	2004	PGE	O1-C1-C2-O2
4	B	601	EDO	O1-C1-C2-O2
3	A	2001	MWF	C3-C2-N2-C4
4	A	2002	EDO	O1-C1-C2-O2
5	A	2004	PGE	C4-C3-O2-C2
3	A	2001	MWF	N3-C11-N5-C15
3	A	2001	MWF	N2-C2-C3-O1
4	A	2003	EDO	O1-C1-C2-O2
5	A	2004	PGE	O3-C5-C6-O4

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1031/1068 (96%)	0.89	105 (10%) 12 9	37, 61, 116, 141	0
2	B	255/293 (87%)	1.15	36 (14%) 6 5	45, 83, 125, 138	0
All	All	1286/1361 (94%)	0.94	141 (10%) 10 8	37, 64, 120, 141	0

All (141) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	411	GLY	4.8
2	B	418	PRO	4.8
2	B	589	VAL	4.5
1	A	3	PRO	4.4
1	A	2	PRO	4.0
2	B	360	ALA	3.9
2	B	413	LEU	3.8
2	B	392	PHE	3.8
1	A	313	THR	3.7
1	A	872	PHE	3.6
1	A	200	PRO	3.6
1	A	213	HIS	3.6
1	A	487	PRO	3.5
1	A	939	ASP	3.5
2	B	366	GLY	3.4
1	A	225	ILE	3.4
1	A	197	ILE	3.3
2	B	401	VAL	3.3
1	A	291	GLU	3.3
2	B	356	LEU	3.3
1	A	217	PRO	3.2
1	A	314	ALA	3.2
2	B	417	ASN	3.2
1	A	947	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	504	ALA	3.2
1	A	207	TYR	3.2
2	B	405	ILE	3.2
1	A	222	ALA	3.1
1	A	221	ILE	3.1
1	A	220	VAL	3.1
1	A	344	VAL	3.1
1	A	328	TRP	3.1
1	A	216	VAL	3.0
2	B	396	LEU	3.0
1	A	198	VAL	2.9
2	B	419	LYS	2.9
1	A	193	VAL	2.9
1	A	967	GLY	2.9
1	A	239	LEU	2.9
2	B	372	LEU	2.9
2	B	333	TRP	2.8
1	A	209	LEU	2.8
1	A	1059	PHE	2.8
1	A	773	SER	2.8
1	A	215	CYS	2.7
1	A	195	TRP	2.7
1	A	208	THR	2.7
1	A	727	THR	2.7
1	A	959	ASP	2.7
1	A	495	HIS	2.7
1	A	312	SER	2.7
1	A	192	VAL	2.7
1	A	940	HIS	2.7
1	A	232	MET	2.6
1	A	336	ILE	2.6
1	A	219	GLN	2.6
1	A	478	ASP	2.6
1	A	492	ILE	2.6
1	A	956	LEU	2.6
1	A	214	ASP	2.6
1	A	224	ALA	2.5
2	B	414	ALA	2.5
1	A	211	ILE	2.5
2	B	381	ILE	2.5
2	B	393	SER	2.5
2	B	395	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	402	VAL	2.5
1	A	474	GLU	2.5
2	B	420	LEU	2.5
1	A	7	SER	2.5
1	A	196	VAL	2.5
1	A	486	PHE	2.5
1	A	152	ASP	2.4
1	A	302	PHE	2.4
1	A	365	GLU	2.4
1	A	278	MET	2.4
1	A	255	CYS	2.4
2	B	518	GLU	2.4
1	A	297	LEU	2.4
1	A	190	ILE	2.4
1	A	864	GLY	2.4
2	B	343	VAL	2.4
1	A	194	ILE	2.4
1	A	945	PHE	2.3
2	B	421	ASP	2.3
1	A	287	LEU	2.3
2	B	331	ALA	2.3
1	A	130	MET	2.3
1	A	881	LEU	2.3
1	A	303	THR	2.3
1	A	5	PRO	2.3
2	B	408	TYR	2.3
1	A	204	LYS	2.2
1	A	1062	ILE	2.2
1	A	498	TRP	2.2
1	A	557	TYR	2.2
2	B	505	SER	2.2
2	B	581	LEU	2.2
1	A	427	ILE	2.2
1	A	500	VAL	2.2
2	B	397	THR	2.2
1	A	343	TYR	2.2
2	B	582	MET	2.2
1	A	227	LYS	2.2
1	A	340	CYS	2.2
2	B	431	TYR	2.2
1	A	938	LEU	2.2
1	A	496	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	443	GLU	2.2
1	A	593	ILE	2.2
2	B	509	ILE	2.2
1	A	559	VAL	2.2
1	A	229	THR	2.2
1	A	185	LEU	2.2
1	A	236	SER	2.2
1	A	522	GLU	2.2
1	A	943	LYS	2.1
1	A	153	LEU	2.1
2	B	355	PHE	2.1
1	A	951	ARG	2.1
1	A	937	PHE	2.1
1	A	1057	TRP	2.1
1	A	333	ALA	2.1
1	A	412	ARG	2.1
1	A	121	ILE	2.1
1	A	304	MET	2.1
1	A	1022	ILE	2.1
1	A	59	HIS	2.1
1	A	228	LYS	2.1
2	B	428	VAL	2.1
2	B	404	LEU	2.1
1	A	4	ARG	2.1
2	B	335	TRP	2.1
1	A	61	LEU	2.0
1	A	299	MET	2.0
1	A	199	SER	2.0
1	A	585	CYS	2.0
1	A	226	ARG	2.0
1	A	309	ARG	2.0
1	A	347	ASN	2.0
1	A	765	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	2003	4/4	0.70	0.19	74,74,74,74	0
5	PGE	A	2004	10/10	0.70	0.19	69,69,69,69	0
4	EDO	B	601	4/4	0.71	0.18	73,73,73,73	0
4	EDO	A	2002	4/4	0.76	0.17	51,51,51,51	0
3	MWF	A	2001	29/29	0.93	0.09	45,45,45,45	0
6	CL	A	2005	1/1	0.95	0.07	64,64,64,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.