



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:19 AM UTC

PDB ID : 2ANR / pdb_00002anr
Title : Crystal structure (II) of Nova-1 KH1/KH2 domain tandem with 25nt RNA hairpin
Authors : Malinina, L.; Teplova, M.; Musunuru, K.; Teplov, A.; Darnell, J.C.; Burley, S.K.; Darnell, R.B.; Patel, D.J.
Deposited on : 2005-08-11
Resolution : 1.94 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)
RNA backbone	3983	1010 (2.30-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	25	 64% 28% • •
2	A	178	 3% 58% 24% • • 13%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 1823 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 5'-R(*CP*(5BU)P*CP*GP*CP*GP*GP*AP*UP*CP*A P*GP*UP*CP*AP*CP*CP*CP*AP*AP*GP*CP*GP*AP*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	25	Total	Br	C	N	O	P	0	0	0
			531	1	238	98	170	24			

- Molecule 2 is a protein called RNA-binding protein Nova-1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	A	155	Total	C	N	O	S	Se	0	2	0
			1183	747	207	226	1	2			

There are 29 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	PRO	-	expression tag	UNP Q9JKN6
A	2	LEU	-	expression tag	UNP Q9JKN6
A	3	GLY	-	expression tag	UNP Q9JKN6
A	4	SER	-	expression tag	UNP Q9JKN6
A	?	-	ARG	deletion	UNP Q9JKN6
A	?	-	ILE	deletion	UNP Q9JKN6
A	?	-	LYS	deletion	UNP Q9JKN6
A	?	-	GLN	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	LEU	deletion	UNP Q9JKN6
A	?	-	PRO	deletion	UNP Q9JKN6
A	?	-	SER	deletion	UNP Q9JKN6
A	?	-	SER	deletion	UNP Q9JKN6
A	?	-	PRO	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	LYS	deletion	UNP Q9JKN6
A	?	-	SER	deletion	UNP Q9JKN6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	SER	deletion	UNP Q9JKN6
A	?	-	PRO	deletion	UNP Q9JKN6
A	?	-	SER	deletion	UNP Q9JKN6
A	?	-	ASP	deletion	UNP Q9JKN6
A	?	-	PRO	deletion	UNP Q9JKN6
A	?	-	MET	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	THR	deletion	UNP Q9JKN6
A	?	-	SER	deletion	UNP Q9JKN6
A	150	ASN	GLU	engineered mutation	UNP Q9JKN6

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total K 1 1	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	2	Total Mg 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	50	Total O 50 50	0	0
5	A	56	Total O 56 56	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-R(*CP*(5BU)P*CP*GP*CP*GP*GP*AP*UP*CP*AP*GP*UP*CP*AP*CP*CP*CP*AP*AP*GP*CP*GP*AP*G)-3'

Chain B: 



- Molecule 2: RNA-binding protein Nova-1

Chain A: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.45Å 37.88Å 38.52Å 90.00° 96.64° 90.00°	Depositor
Resolution (Å)	20.00 – 1.94 20.00 – 1.94	Depositor EDS
% Data completeness (in resolution range)	94.2 (20.00-1.94) 94.1 (20.00-1.94)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.38 (at 1.91Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.227 , 0.268 0.243 , 0.300	Depositor DCC
R_{free} test set	858 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	38.8	Xtriage
Anisotropy	0.306	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1823	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 5BU, MG, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	0.80	0/570	0.94	1/887 (0.1%)
2	A	0.77	0/1205	1.04	0/1620
All	All	0.78	0/1775	1.00	1/2507 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	212	G	C3'-C2'-C1'	5.73	107.03	101.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	A	79	MSE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	531	0	273	5	0
2	A	1183	0	1233	33	0
3	B	1	0	0	0	0
4	B	2	0	0	0	0
5	A	56	0	0	0	0
5	B	50	0	0	0	0
All	All	1823	0	1506	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:119:ILE:HA	2:A:126:VAL:HG23	1.44	0.99
2:A:130:MSE:HG3	2:A:137:VAL:HG12	1.54	0.87
2:A:119:ILE:HA	2:A:126:VAL:CG2	2.07	0.83
2:A:30:GLN:O	2:A:34:GLU:HG3	1.81	0.80
1:B:216:C:O2	2:A:40:LYS:HD2	1.91	0.71
2:A:5:GLN:HG2	2:A:59:GLN:HG2	1.83	0.59
2:A:79:MSE:H	2:A:80:PRO:HD3	1.70	0.57
2:A:118:ILE:O	2:A:126:VAL:HG22	2.06	0.55
2:A:134:GLY:HA3	2:A:161:GLN:HG2	1.89	0.55
2:A:5:GLN:CG	2:A:59:GLN:HG2	2.37	0.54
2:A:62:ILE:HD11	2:A:174:GLN:CD	2.33	0.53
2:A:134:GLY:CA	2:A:161:GLN:HG2	2.38	0.53
2:A:126:VAL:CG1	2:A:137:VAL:HG13	2.39	0.53
2:A:130:MSE:CG	2:A:137:VAL:HG12	2.34	0.52
2:A:114:THR:HG22	2:A:176:ASP:OD2	2.10	0.52
2:A:44:SER:HA	2:A:54:ARG:HD2	1.94	0.50
2:A:126:VAL:HG12	2:A:130:MSE:HE2	1.94	0.49
2:A:150:ASN:N	2:A:150:ASN:OD1	2.45	0.49
2:A:79:MSE:H	2:A:80:PRO:CD	2.26	0.49
2:A:43:LYS:O	2:A:46:ASP:HB2	2.14	0.48
2:A:158:GLU:HB2	2:A:161:GLN:OE1	2.11	0.48
2:A:11:LEU:HB2	2:A:107:LYS:HB2	1.95	0.47
1:B:209:U:H2'	1:B:210:C:O4'	2.14	0.47
2:A:158:GLU:H	2:A:161:GLN:NE2	2.11	0.47
2:A:42:SER:HB3	2:A:46:ASP:HB3	1.97	0.45
1:B:212:G:H4'	1:B:214:C:C5	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:113:SER:OG	2:A:178:GLN:OE1	2.33	0.44
2:A:5:GLN:HG2	2:A:59:GLN:CG	2.47	0.43
2:A:79:MSE:N	2:A:80:PRO:CD	2.82	0.42
1:B:214:C:H4'	1:B:215:A:H5'	2.00	0.42
2:A:160:GLU:H	2:A:160:GLU:CD	2.27	0.42
1:B:202:5BU:H2'	1:B:203:C:C6	2.55	0.41
2:A:118:ILE:O	2:A:126:VAL:CG2	2.69	0.41
2:A:130:MSE:HG2	2:A:135:ALA:O	2.20	0.41
2:A:12:ILE:HA	2:A:13:PRO:HD2	1.96	0.40
2:A:130:MSE:O	2:A:133:SER:O	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	151/178 (85%)	142 (94%)	6 (4%)	3 (2%)	6 1

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	79	MSE
2	A	141	GLN
2	A	80	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	130/147 (88%)	107 (82%)	23 (18%)	2 0

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	5	GLN
2	A	10	VAL
2	A	14	SER
2	A	30	GLN
2	A	46	ASP
2	A	52	THR
2	A	63[A]	GLU
2	A	63[B]	GLU
2	A	77	ARG
2	A	78	GLU
2	A	101	ASP
2	A	102	ARG
2	A	112	ASN
2	A	114	THR
2	A	121	LYS
2	A	131	GLU
2	A	138	GLN
2	A	149	GLN
2	A	150	ASN
2	A	161	GLN
2	A	163	ARG
2	A	164	LYS
2	A	168	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	59	GLN
2	A	112	ASN
2	A	132	GLN
2	A	138	GLN
2	A	149	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	B	24/25 (96%)	1 (4%)	1 (4%)

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	B	213	U

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	B	212	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	5BU	B	202	1	19,22,23	1.82	3 (15%)	27,32,35	2.14	10 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	5BU	B	202	1	-	0/7/25/26	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	202	5BU	C4-C5	-5.71	1.33	1.45
1	B	202	5BU	C6-N1	-4.13	1.31	1.38
1	B	202	5BU	C2-N1	-2.51	1.34	1.38

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	202	5BU	O4-C4-C5	-4.87	119.59	125.80
1	B	202	5BU	BR-C5-C4	4.30	122.97	118.02
1	B	202	5BU	C1'-N1-C2	4.00	124.78	117.59
1	B	202	5BU	C6-N1-C2	-3.46	117.86	121.30
1	B	202	5BU	O4'-C1'-N1	2.65	114.37	108.36
1	B	202	5BU	C2'-C1'-N1	-2.52	106.22	113.25
1	B	202	5BU	O4-C4-N3	2.45	124.72	120.11
1	B	202	5BU	C1'-N1-C6	-2.30	117.36	121.15
1	B	202	5BU	C5-C6-N1	2.14	125.91	120.91
1	B	202	5BU	C5-C4-N3	2.04	115.69	113.34

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	202	5BU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	B	24/25 (96%)	-0.20	0	100	100	16, 22, 29, 45	0
2	A	153/178 (85%)	0.46	5 (3%)	49	55	13, 24, 39, 46	4 (2%)
All	All	177/203 (87%)	0.37	5 (2%)	55	62	13, 24, 39, 46	4 (2%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	80	PRO	3.9
2	A	143	PRO	2.6
2	A	148	LEU	2.6
2	A	104	ASN	2.3
2	A	136	TRP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	5BU	B	202	21/22	0.80	0.10	38,40,46,47	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MG	B	302	1/1	0.84	0.09	15,15,15,15	1
4	MG	B	306	1/1	0.84	0.07	4,4,4,4	1
3	K	B	301	1/1	0.96	0.04	19,19,19,19	1

6.5 Other polymers [i](#)

There are no such residues in this entry.