



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:26 PM UTC

PDB ID : 2ANW / pdb_00002anw
Title : Expression, crystallization and three-dimensional structure of the catalytic domain of human plasma kallikrein: Implications for structure-based design of protease inhibitors
Authors : Tang, J.; Yu, C.L.; Williams, S.R.; Springman, E.; Jeffery, D.; Sprengeler, P.A.; Estevez, A.; Sampang, J.; Shrader, W.; Spencer, J.R.; Young, W.B.; McGrath, M.E.; Katz, B.A.
Deposited on : 2005-08-11
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	241	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2981 atoms, of which 867 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called plasma kallikrein, light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	238	2321	1203	430	325	352	11	0	3	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	SER	CYS	engineered mutation	UNP P03952

- Molecule 2 is BENZAMIDINE (CCD ID: BEN) (formula: C₇H₈N₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	N		
2	A	1	18	7	9	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	210	Total	H	O	0	4
			642	428	214		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: plasma kallikrein, light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	79.89Å 63.19Å 50.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	7.00 – 1.85 7.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.4 (7.00-1.85) 94.6 (7.00-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.80 (at 1.86Å)	Xtriage
Refinement program	XTALVIEW, X-PLOR 3.851	Depositor
R, R_{free}	0.221 , 0.283 0.213 , 0.213	Depositor DCC
R_{free} test set	2123 reflections (10.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.2	Xtriage
Anisotropy	0.654	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 89.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2981	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.39	10/1951 (0.5%)	1.84	41/2646 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	40	HIS	C-N	20.43	1.61	1.33
1	A	39	ARG	C-N	20.37	1.63	1.33
1	A	134	THR	C-N	8.62	1.54	1.34
1	A	57	HIS	CD2-NE2	-7.05	1.30	1.37
1	A	101	HIS	CD2-NE2	-6.92	1.30	1.37
1	A	133	TYR	C-N	-5.74	1.25	1.33
1	A	91	HIS	CD2-NE2	-5.71	1.31	1.37
1	A	132	ILE	CA-CB	5.57	1.60	1.53
1	A	40	HIS	CD2-NE2	-5.33	1.31	1.37
1	A	203	HIS	CD2-NE2	-5.21	1.32	1.37

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	133	TYR	O-C-N	-10.14	110.81	122.68
1	A	189	ASP	CA-CB-CG	9.54	122.14	112.60
1	A	165	ASN	OD1-CG-ND2	-9.31	113.28	122.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	86	LYS	N-CA-C	-8.93	101.95	113.12
1	A	243	GLN	OE1-CD-NE2	-8.33	114.27	122.60
1	A	40	HIS	O-C-N	-8.32	111.95	123.01
1	A	101	HIS	CB-CG-CD2	-8.15	120.60	131.20
1	A	57	HIS	CB-CG-CD2	-7.68	121.22	131.20
1	A	134	THR	O-C-N	7.40	132.43	122.59
1	A	72	ASN	OD1-CG-ND2	-6.93	115.67	122.60
1	A	212	ILE	O-C-N	-6.54	115.44	123.10
1	A	153	ASN	OD1-CG-ND2	-6.37	116.23	122.60
1	A	199	LEU	N-CA-C	-6.26	97.07	108.02
1	A	57	HIS	CA-CB-CG	6.24	120.04	113.80
1	A	49	HIS	CB-CG-CD2	-6.07	123.31	131.20
1	A	49	HIS	CB-CG-ND1	6.07	131.80	122.70
1	A	91	HIS	CB-CG-CD2	-6.01	123.38	131.20
1	A	163	VAL	O-C-N	-5.98	116.75	123.03
1	A	38(B)	GLN	OE1-CD-NE2	-5.93	116.67	122.60
1	A	179	ARG	CB-CG-CD	5.90	124.88	111.30
1	A	141	TRP	CG-CD1-NE1	-5.89	102.54	110.20
1	A	165	ASN	CB-CG-ND2	5.89	125.24	116.40
1	A	159	ASN	CB-CG-ND2	5.76	125.04	116.40
1	A	37	LEU	O-C-N	-5.66	117.90	122.03
1	A	39	ARG	O-C-N	5.65	129.92	123.31
1	A	237	TRP	CG-CD1-NE1	-5.60	102.93	110.20
1	A	159	ASN	OD1-CG-ND2	-5.47	117.13	122.60
1	A	115	THR	O-C-N	-5.36	115.34	122.74
1	A	57	HIS	CB-CG-ND1	5.34	130.72	122.70
1	A	243	GLN	CG-CD-NE2	5.25	124.27	116.40
1	A	27	TRP	CB-CG-CD1	-5.21	119.08	126.90
1	A	124	PRO	CB-CA-C	5.20	118.44	110.75
1	A	88	ILE	O-C-N	-5.15	117.78	123.18
1	A	123	LEU	CA-C-O	-5.12	115.99	120.60
1	A	91	HIS	CA-C-N	5.10	128.46	120.82
1	A	91	HIS	C-N-CA	5.10	128.46	120.82
1	A	185	LYS	CA-CB-CG	5.09	124.29	114.10
1	A	97	SER	CA-CB-OG	5.09	121.28	111.10
1	A	50	GLN	N-CA-C	5.09	118.58	111.56
1	A	119	LYS	CA-C-N	5.07	125.05	120.03
1	A	119	LYS	C-N-CA	5.07	125.05	120.03

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	133	TYR	Peptide,Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1891	430	1855	22	0
2	A	9	9	8	0	0
3	A	214	428	0	4	0
All	All	2114	867	1863	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:240:GLU:O	1:A:244:SER:HB2	2.04	0.57
1:A:38[B]:THR:HG22	3:A:725:HOH:O	2.06	0.56
1:A:138:VAL:HG21	1:A:160:ILE:HD12	1.92	0.50
1:A:113:ASN:HB3	3:A:397:HOH:O	2.11	0.50
1:A:61:LEU:O	1:A:65(C):VAL:HG22	2.12	0.49
1:A:139:THR:HG22	1:A:157:LYS:HA	1.96	0.47
1:A:134:THR:O	1:A:134:THR:HG23	2.14	0.46
1:A:36:LYS:HB3	3:A:911:HOH:O	2.16	0.46
1:A:142:GLY:HA2	1:A:151:ILE:HD13	1.99	0.45
1:A:29:TRP:CG	1:A:121:ILE:HB	2.52	0.44
1:A:236:ASP:O	1:A:240:GLU:HG2	2.18	0.44
1:A:211:GLY:HA2	1:A:229:THR:O	2.18	0.43
1:A:93:ASN:HB2	1:A:101:HIS:CD2	2.54	0.43
1:A:72:ASN:HA	1:A:153:ASN:O	2.19	0.43
1:A:33:LEU:HD22	1:A:66:ILE:HG12	2.01	0.42
1:A:115:THR:O	1:A:119:LYS:HG3	2.19	0.42
1:A:185:LYS:HG3	1:A:223:GLU:CD	2.45	0.42
1:A:86:LYS:HE3	3:A:922:HOH:O	2.20	0.42
1:A:213:THR:HA	1:A:228:TYR:CD2	2.56	0.41
1:A:217:GLU:OE1	1:A:217:GLU:HA	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:PHE:HA	1:A:150:GLU:O	2.20	0.41
1:A:35:VAL:HG22	1:A:41:LEU:HG	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	239/241 (99%)	232 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	208/207 (100%)	196 (94%)	12 (6%)	18	6

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	73	LEU
1	A	134	THR
1	A	145	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	163	VAL
1	A	167	GLU
1	A	173	GLN
1	A	173(A)	ASP
1	A	186	GLU
1	A	186(C)	LYS
1	A	192	LYS
1	A	244	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	21	ASN
1	A	38(B)	GLN
1	A	93	ASN
1	A	118	GLN
1	A	153	ASN
1	A	156	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BEN	A	249	-	9,9,9	1.50	3 (33%)	7,11,11	1.03	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BEN	A	249	-	-	0/4/4/4	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	249	BEN	C1-C	-2.99	1.41	1.47
2	A	249	BEN	C6-C1	2.08	1.42	1.39
2	A	249	BEN	C-N1	2.06	1.36	1.27

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	39:ARG	C	40:HIS	N	1.63
1	A	40:HIS	C	41:LEU	N	1.61

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	238/241 (98%)	-0.44	3 (1%) 75 79	12, 28, 52, 70	9 (3%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	61	LEU	4.2
1	A	60(C)	PRO	4.1
1	A	244	SER	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BEN	A	249	9/9	0.91	0.06	24,26,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.