



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:03 PM UTC

PDB ID : 3ANS / pdb_00003ans
Title : Human soluble epoxide hydrolase in complex with a synthetic inhibitor
Authors : Chiyo, N.; Ishii, T.; Hourai, S.; Yanagi, K.
Deposited on : 2010-09-08
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	336	16% 82% 10% 7%
1	B	336	11% 81% 11% 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5348 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Epoxide hydrolase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	314	Total	C	N	O	S	0	0	0
			2533	1629	424	457	23			
1	B	314	Total	C	N	O	S	0	0	0
			2533	1629	424	457	23			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	220	MET	-	expression tag	UNP P34913
A	221	LYS	-	expression tag	UNP P34913
A	222	LYS	-	expression tag	UNP P34913
A	223	GLY	-	expression tag	UNP P34913
A	224	HIS	-	expression tag	UNP P34913
A	225	HIS	-	expression tag	UNP P34913
A	226	HIS	-	expression tag	UNP P34913
A	227	HIS	-	expression tag	UNP P34913
A	228	HIS	-	expression tag	UNP P34913
A	229	HIS	-	expression tag	UNP P34913
B	220	MET	-	expression tag	UNP P34913
B	221	LYS	-	expression tag	UNP P34913
B	222	LYS	-	expression tag	UNP P34913
B	223	GLY	-	expression tag	UNP P34913
B	224	HIS	-	expression tag	UNP P34913
B	225	HIS	-	expression tag	UNP P34913
B	226	HIS	-	expression tag	UNP P34913
B	227	HIS	-	expression tag	UNP P34913
B	228	HIS	-	expression tag	UNP P34913
B	229	HIS	-	expression tag	UNP P34913

- Molecule 2 is 4-cyano-N-[(1S,2R)-2-phenylcyclopropyl]benzamide (CCD ID: S38) (formula: C₁₇H₁₄N₂O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			20	17	2	1		
2	B	1	Total	C	N	O	0	0
			20	17	2	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	107	Total	O	0	0
			107	107		
3	B	135	Total	O	0	0
			135	135		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Epoxide hydrolase 2



• Molecule 1: Epoxide hydrolase 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	129.63Å 80.35Å 88.69Å 90.00° 126.20° 90.00°	Depositor
Resolution (Å)	31.21 – 1.98 31.21 – 1.98	Depositor EDS
% Data completeness (in resolution range)	98.7 (31.21-1.98) 98.7 (31.21-1.98)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.76 (at 1.98Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.219 , 0.260 0.215 , 0.254	Depositor DCC
R_{free} test set	2566 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.501	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 37.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5348	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: S38

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.60	0/2609	0.82	0/3539
1	B	0.61	0/2609	0.85	1/3539 (0.0%)
All	All	0.61	0/5218	0.83	1/7078 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	505	GLN	N-CA-C	5.46	117.99	111.71

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	291	MET	Peptide
1	B	291	MET	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2533	0	2461	19	0
1	B	2533	0	2461	29	0
2	A	20	0	13	3	0
2	B	20	0	12	4	0
3	A	107	0	0	2	0
3	B	135	0	0	2	0
All	All	5348	0	4947	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:339:MET:HE3	2:B:602:S38:N20	1.69	1.07
1:A:339:MET:HE3	2:A:601:S38:N20	1.81	0.95
1:B:460:ARG:HH21	1:B:460:ARG:HG3	1.30	0.94
1:B:513:HIS:H	1:B:513:HIS:CD2	1.90	0.89
1:A:247:ARG:HH11	1:A:247:ARG:HG2	1.47	0.79
1:B:339:MET:CE	2:B:602:S38:N20	2.45	0.79
1:B:486:ILE:O	1:B:513:HIS:CE1	2.39	0.75
1:B:460:ARG:HH21	1:B:460:ARG:CG	2.05	0.68
1:B:524:HIS:NE2	2:B:602:S38:H11	2.09	0.67
1:A:524:HIS:NE2	2:A:601:S38:H11	2.15	0.62
1:B:266:GLY:HA3	1:B:335:ASP:HB3	1.81	0.62
1:A:247:ARG:HH11	1:A:247:ARG:CG	2.13	0.62
1:A:266:GLY:HA3	1:A:335:ASP:HB3	1.83	0.61
1:A:428:LEU:HD21	2:A:601:S38:H15	1.84	0.60
1:B:306:GLU:HG3	1:B:471:ARG:NH1	2.17	0.59
1:B:277:GLN:HG3	1:B:535:ASN:ND2	2.22	0.55
1:A:347:PHE:CZ	1:A:479:SER:HB3	2.41	0.55
1:A:366:ASN:HB3	1:A:369:MET:HE2	1.90	0.54
1:A:479:SER:O	1:A:481:GLY:N	2.41	0.53
1:A:413:ASP:OD1	1:A:440:ARG:NH2	2.39	0.53
1:B:500:VAL:HG12	3:B:748:HOH:O	2.10	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:355:VAL:O	1:B:488:ALA:HA	2.13	0.48
1:B:486:ILE:O	1:B:513:HIS:HE1	1.89	0.48
1:A:245:LYS:HB2	1:A:246:PRO:HD2	1.96	0.48
1:B:468:ASN:OD1	1:B:471:ARG:NH1	2.47	0.47
1:A:277:GLN:HG3	1:A:535:ASN:ND2	2.29	0.47
1:B:500:VAL:CG1	3:B:748:HOH:O	2.61	0.47
1:B:386:TYR:OH	1:B:396:GLU:OE1	2.29	0.47
1:B:305:ILE:HG22	1:B:471:ARG:HH12	1.79	0.46
1:A:339:MET:HE2	1:A:360:THR:CG2	2.46	0.46
1:A:507:MET:HA	1:A:510:TRP:CE2	2.51	0.46
1:B:428:LEU:HD21	2:B:602:S38:H15	1.99	0.45
1:B:378:ASN:HD22	1:B:380:VAL:H	1.64	0.45
1:A:247:ARG:CG	1:A:247:ARG:NH1	2.76	0.45
1:B:420:HIS:HB2	1:B:421:LYS:HD2	2.00	0.44
1:B:460:ARG:HG3	1:B:460:ARG:NH2	2.10	0.44
1:B:306:GLU:HA	1:B:471:ARG:HH11	1.83	0.43
1:B:306:GLU:HG3	1:B:471:ARG:HH11	1.80	0.43
1:B:342:TRP:CG	1:B:362:PHE:HB2	2.54	0.42
1:B:332:ILE:HG12	1:B:356:ALA:HB3	2.02	0.42
1:B:254:GLU:HA	1:B:288:VAL:O	2.19	0.42
1:A:389:GLU:HA	1:A:390:PRO:HD3	1.92	0.42
1:A:315:LYS:HG3	3:A:755:HOH:O	2.19	0.41
1:A:311:GLU:HG2	1:A:315:LYS:HE2	2.01	0.41
1:B:345:ALA:HB1	1:B:486:ILE:HD11	2.02	0.41
1:B:237:MET:HB3	1:B:253:VAL:HG13	2.02	0.41
1:A:469:MET:SD	3:A:942:HOH:O	2.63	0.41
1:B:507:MET:HA	1:B:510:TRP:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	312/336 (93%)	300 (96%)	10 (3%)	2 (1%)	21	12
1	B	312/336 (93%)	303 (97%)	8 (3%)	1 (0%)	36	27
All	All	624/672 (93%)	603 (97%)	18 (3%)	3 (0%)	24	14

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	480	LEU
1	B	268	PRO
1	A	268	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/296 (93%)	264 (96%)	12 (4%)	26	14
1	B	276/296 (93%)	264 (96%)	12 (4%)	26	14
All	All	552/592 (93%)	528 (96%)	24 (4%)	26	14

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	ARG
1	A	340	LEU
1	A	346	LEU
1	A	369	MET
1	A	385	LEU
1	A	421	LYS
1	A	423	CYS
1	A	431	ASN
1	A	435	GLU
1	A	480	LEU
1	A	489	LEU
1	A	508	GLU
1	B	340	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	346	LEU
1	B	378	ASN
1	B	380	VAL
1	B	418	SER
1	B	421	LYS
1	B	423	CYS
1	B	460	ARG
1	B	471	ARG
1	B	498	VAL
1	B	508	GLU
1	B	513	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	378	ASN
1	A	384	GLN
1	A	453	GLN
1	A	535	ASN
1	B	328	GLN
1	B	366	ASN
1	B	378	ASN
1	B	384	GLN
1	B	452	GLN
1	B	513	HIS
1	B	535	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	S38	A	601	-	21,22,22	1.06	2 (9%)	24,30,30	1.44	4 (16%)
2	S38	B	602	-	21,22,22	1.12	3 (14%)	24,30,30	1.32	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	S38	A	601	-	-	2/14/19/19	0/3/3/3
2	S38	B	602	-	-	2/14/19/19	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	602	S38	C12-C10	-2.82	1.47	1.51
2	A	601	S38	C12-C10	-2.57	1.48	1.51
2	A	601	S38	C9-N8	-2.19	1.42	1.46
2	B	602	S38	C9-N8	-2.15	1.42	1.46
2	B	602	S38	C5-C7	-2.02	1.45	1.50

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	602	S38	C9-N8-C7	-3.63	116.34	122.94
2	A	601	S38	C9-N8-C7	-3.28	116.96	122.94
2	A	601	S38	C6-C5-C4	3.07	122.47	118.57
2	B	602	S38	C6-C5-C4	2.88	122.23	118.57
2	A	601	S38	C3-C4-C5	-2.42	118.21	120.80
2	A	601	S38	O18-C7-C5	-2.25	116.44	120.90

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	602	S38	C11-C10-C12-C13
2	A	601	S38	C11-C10-C12-C17
2	B	602	S38	C11-C10-C12-C17
2	A	601	S38	C11-C10-C12-C13


There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	S38	3	0
2	B	602	S38	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/336 (93%)	0.79	53 (16%) 4 5	18, 31, 71, 79	0
1	B	314/336 (93%)	0.61	38 (12%) 8 12	16, 28, 70, 76	0
All	All	628/672 (93%)	0.70	91 (14%) 6 8	16, 29, 71, 79	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	377	ALA	7.0
1	A	380	VAL	6.1
1	A	377	ALA	5.9
1	B	510	TRP	5.6
1	B	232	CYS	5.4
1	B	372	LEU	5.3
1	B	380	VAL	5.2
1	A	375	ILE	4.9
1	A	372	LEU	4.7
1	A	381	PHE	4.7
1	A	425	ALA	4.7
1	A	416	VAL	4.6
1	B	512	PRO	4.4
1	A	417	LEU	4.4
1	B	420	HIS	4.3
1	B	425	ALA	4.3
1	A	510	TRP	4.2
1	A	423	CYS	4.1
1	A	379	PRO	4.1
1	B	430	VAL	4.0
1	A	433	PRO	3.9
1	A	422	VAL	3.9
1	B	375	ILE	3.9
1	A	480	LEU	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	371	PRO	3.9
1	B	473	TRP	3.8
1	B	381	PHE	3.8
1	A	418	SER	3.7
1	B	378	ASN	3.7
1	B	423	CYS	3.5
1	B	369	MET	3.5
1	B	480	LEU	3.5
1	B	513	HIS	3.5
1	A	430	VAL	3.5
1	B	422	VAL	3.5
1	A	369	MET	3.4
1	A	419	MET	3.4
1	B	421	LYS	3.4
1	B	366	ASN	3.4
1	A	506	HIS	3.3
1	B	376	LYS	3.2
1	A	374	SER	3.2
1	B	371	PRO	3.1
1	A	367	PRO	3.1
1	B	419	MET	3.0
1	A	421	LYS	3.0
1	A	363	ILE	3.0
1	A	420	HIS	3.0
1	A	473	TRP	3.0
1	A	427	GLY	3.0
1	A	544	SER	2.9
1	A	232	CYS	2.9
1	B	424	GLU	2.9
1	B	426	GLY	2.8
1	A	479	SER	2.8
1	A	431	ASN	2.7
1	B	428	LEU	2.7
1	A	481	GLY	2.7
1	A	500	VAL	2.7
1	B	382	ASP	2.6
1	A	412	SER	2.6
1	B	367	PRO	2.6
1	B	379	PRO	2.6
1	A	366	ASN	2.6
1	B	374	SER	2.6
1	B	432	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	509	ASP	2.5
1	A	505	GLN	2.5
1	A	236	ASP	2.4
1	B	373	GLU	2.4
1	A	426	GLY	2.4
1	A	383	TYR	2.3
1	A	364	PRO	2.2
1	A	365	ALA	2.2
1	A	343	TYR	2.2
1	A	362	PHE	2.2
1	A	385	LEU	2.2
1	A	545	ASP	2.2
1	A	499	LEU	2.2
1	A	368	ASN	2.1
1	B	545	ASP	2.1
1	B	481	GLY	2.1
1	B	385	LEU	2.1
1	A	445	GLU	2.1
1	A	378	ASN	2.1
1	A	429	PHE	2.1
1	A	382	ASP	2.1
1	B	506	HIS	2.0
1	A	432	SER	2.0
1	B	544	SER	2.0
1	A	233	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ


There are no oligosaccharides in this entry.


6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	S38	B	602	20/20	0.84	0.12	36,40,44,44	0
2	S38	A	601	20/20	0.86	0.12	42,45,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.