



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 01:21 AM UTC

PDB ID : 4ANB / pdb_00004anb
Title : Crystal structures of human MEK1 with carboxamide-based allosteric inhibitor XL518 (GDC-0973), or related analogs.
Authors : Rice, K.D.; Aay, N.; Anand, N.K.; Blazey, C.M.; Bowles, O.J.; Bussenius, J.; Costanzo, S.; Curtis, J.K.; Defina, S.C.; Dubenko, L.; Engst, S.; Joshi, A.A.; Kennedy, A.R.; Kim, A.I.; Koltun, E.S.; Loughheed, J.C.; Manalo, J.C.L.; Martini, J.F.; Nuss, J.M.; Peto, C.J.; Tsang, T.H.; Yu, P.; Johnston, S.
Deposited on : 2012-03-16
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	9% 72% 17% 6%

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DUAL SPECIFICITY MITOGEN-ACTIVATED PROTEIN KINASE KINASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	0	0
			2209	1412	376	405	16			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	58	MET	-	expression tag	UNP Q02750
A	59	ALA	-	expression tag	UNP Q02750
A	60	LEU	-	expression tag	UNP Q02750
A	393	GLU	-	expression tag	UNP Q02750
A	394	GLY	-	expression tag	UNP Q02750
A	395	HIS	-	expression tag	UNP Q02750
A	396	HIS	-	expression tag	UNP Q02750
A	397	HIS	-	expression tag	UNP Q02750
A	398	HIS	-	expression tag	UNP Q02750
A	399	HIS	-	expression tag	UNP Q02750
A	400	HIS	-	expression tag	UNP Q02750
A	218	GLU	SER	engineered mutation	UNP Q02750
A	222	GLU	SER	engineered mutation	UNP Q02750
A	328	ALA	GLY	engineered mutation	UNP Q02750

- Molecule 2 is [3-(AMINOMETHYL)-3-OXIDANYL-AZETIDIN-1-YL]-[3,4-BIS(FLUORANYL)-2-[(2-FLUORANYL-4-IODANYL-PHENYL)AMINO]PHENYL]METHANONE (CCD ID: YQY) (formula: C₁₇H₁₅F₃IN₃O₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	F	I	N	O	0	0
			26	17	3	1	3	2		

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		


- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	57	Total	O	0	0
			57	57		

● Molecule 1: DUAL SPECIFICITY MITOGEN-ACTIVATED PROTEIN KINASE KINASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	109.00Å 109.00Å 48.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	94.49 – 2.20 94.40 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (94.49-2.20) 99.4 (94.40-2.20)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.04 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.226 , 0.296 0.223 , 0.292	Depositor DCC
R_{free} test set	858 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	56.8	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.039 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2324	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACP, YQY, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.25	2/2254 (0.1%)	1.28	13/3038 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	262	PRO	C-N	26.09	1.65	1.32
1	A	208	ASP	CA-C	-5.44	1.46	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	262	PRO	O-C-N	-7.45	112.59	122.64
1	A	211	VAL	CB-CA-C	-7.44	102.27	112.02
1	A	262	PRO	CA-C-N	-6.05	115.25	123.96
1	A	262	PRO	C-N-CA	-6.05	115.25	123.96
1	A	174	ILE	CB-CA-C	-5.62	104.68	112.04
1	A	215	LEU	N-CA-C	-5.59	105.27	111.36
1	A	334	PHE	N-CA-C	-5.53	104.65	111.40
1	A	88	LYS	CA-C-N	5.41	125.05	119.05
1	A	88	LYS	C-N-CA	5.41	125.05	119.05
1	A	161	ILE	N-CA-C	-5.32	101.77	108.05
1	A	211	VAL	N-CA-C	5.29	115.50	110.53
1	A	205	LYS	N-CA-C	5.21	117.93	109.24
1	A	310	ILE	N-CA-C	5.15	115.77	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2209	0	2237	41	0
2	A	26	0	15	3	0
3	A	31	0	14	0	0
4	A	1	0	0	0	0
5	A	57	0	0	5	2
All	All	2324	0	2266	44	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:67:ASP:O	1:A:88:LYS:HE2	1.70	0.89
1:A:170:SER:O	1:A:174:ILE:HG12	1.79	0.81
1:A:311:PHE:CD2	5:A:612:HOH:O	2.35	0.80
1:A:187:MET:HE3	1:A:240:TYR:CD1	2.18	0.78
1:A:187:MET:HE3	1:A:240:TYR:HD1	1.50	0.74
1:A:183:LYS:HB3	1:A:184:HIS:CD2	2.23	0.72
1:A:229:TYR:OH	1:A:255:GLU:OE1	2.09	0.70
1:A:316:TYR:CE1	1:A:320:GLU:HG3	2.29	0.67
1:A:183:LYS:HD3	1:A:184:HIS:NE2	2.12	0.64
1:A:108:ARG:O	1:A:112:ILE:HG13	1.97	0.64
1:A:320:GLU:O	1:A:344:LYS:NZ	2.28	0.62
1:A:186:ILE:HD13	1:A:211:VAL:HG13	1.81	0.62
1:A:215:LEU:O	1:A:219:MET:HB2	2.01	0.61
1:A:378:THR:HG22	1:A:379:ILE:HG23	1.84	0.60
1:A:160:ARG:HG2	1:A:257:ALA:O	2.03	0.57
1:A:187:MET:CE	1:A:240:TYR:HD1	2.18	0.57
1:A:339:ASN:O	1:A:343:ILE:HD13	2.05	0.57
1:A:363:ARG:NH2	5:A:602:HOH:O	2.30	0.54
1:A:189:ARG:HA	1:A:240:TYR:CZ	2.44	0.52
1:A:111:ILE:HG21	1:A:139:ILE:HD13	1.91	0.52
1:A:307:PRO:O	1:A:308:MET:HB3	2.09	0.52
1:A:255:GLU:HB2	1:A:262:PRO:HD3	1.93	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:343:ILE:HG22	1:A:345:ASN:H	1.76	0.50
1:A:254:VAL:HG12	1:A:262:PRO:HB3	1.94	0.49
2:A:501:YQY:H211	2:A:501:YQY:H12	1.95	0.49
1:A:67:ASP:C	1:A:88:LYS:HE2	2.37	0.49
1:A:73:GLU:HG2	1:A:75:GLY:H	1.78	0.49
2:A:501:YQY:H12	2:A:501:YQY:C21	2.42	0.48
1:A:222:GLU:HG2	1:A:223:PHE:CZ	2.49	0.48
1:A:241:SER:HB2	5:A:646:HOH:O	2.13	0.47
1:A:205:LYS:NZ	5:A:605:HOH:O	2.38	0.46
1:A:213:GLY:HA2	1:A:216:ILE:HD12	1.96	0.46
1:A:112:ILE:HG22	1:A:116:GLN:NE2	2.32	0.44
1:A:118:LEU:HB2	1:A:129:PHE:CD2	2.52	0.43
1:A:66:ASP:O	1:A:88:LYS:NZ	2.23	0.43
1:A:132:ALA:HA	1:A:140:SER:O	2.19	0.43
1:A:343:ILE:HD13	1:A:343:ILE:N	2.34	0.43
1:A:229:TYR:CD1	1:A:251:LEU:HD23	2.54	0.42
2:A:501:YQY:C21	2:A:501:YQY:C12	2.97	0.42
1:A:74:LEU:HD12	1:A:82:VAL:HG12	2.01	0.42
1:A:320:GLU:HB3	1:A:321:PRO:CD	2.50	0.41
1:A:88:LYS:HE3	1:A:88:LYS:HB2	1.86	0.41
1:A:167:GLY:HA2	1:A:360:PHE:CE2	2.56	0.41
1:A:186:ILE:HA	5:A:611:HOH:O	2.20	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:655:HOH:O	5:A:657:HOH:O[5_554]	2.11	0.09
5:A:633:HOH:O	5:A:657:HOH:O[5_554]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	280/301 (93%)	266 (95%)	13 (5%)	1 (0%)	30	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	308	MET

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	243/258 (94%)	228 (94%)	15 (6%)	16	20

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	66	ASP
1	A	94	MET
1	A	98	LEU
1	A	112	ILE
1	A	135	SER
1	A	164	GLN
1	A	183	LYS
1	A	187	MET
1	A	211	VAL
1	A	219	MET
1	A	222	GLU
1	A	226	THR
1	A	329	VAL
1	A	338	VAL
1	A	343	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	78	ASN
1	A	122	ASN
1	A	164	GLN
1	A	221	ASN
1	A	335	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	YQY	A	501	-	26,28,28	1.49	4 (15%)	28,42,42	2.06	10 (35%)
3	ACP	A	502	4	31,33,33	1.73	7 (22%)	47,52,52	2.15	13 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	YQY	A	501	-	-	2/12/25/25	0/3/3/3
3	ACP	A	502	4	-	1/19/38/38	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	ACP	PB-O3A	4.46	1.63	1.58
2	A	501	YQY	C24-C20	-4.17	1.52	1.56
3	A	502	ACP	C5-C6	3.82	1.51	1.41
3	A	502	ACP	C5-C4	3.38	1.45	1.39
3	A	502	ACP	PA-O3A	3.32	1.63	1.59
3	A	502	ACP	C5-N7	-2.89	1.33	1.39
2	A	501	YQY	C2-I14	2.74	2.17	2.10
2	A	501	YQY	C3-C4	2.67	1.43	1.38
2	A	501	YQY	F26-C9	-2.49	1.31	1.34
3	A	502	ACP	PG-O3G	-2.35	1.49	1.55
3	A	502	ACP	PB-O1B	-2.01	1.46	1.51

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	ACP	C5-C4-N3	-6.53	117.73	126.72
3	A	502	ACP	N3-C4-N9	5.05	135.76	127.17
3	A	502	ACP	C2-N3-C4	4.50	122.81	111.83
2	A	501	YQY	C4-C3-C2	3.78	124.39	119.55
2	A	501	YQY	C3-C2-I14	3.76	125.31	119.69
2	A	501	YQY	C19-N17-C21	3.63	98.29	95.12
3	A	502	ACP	C4-C5-N7	-3.55	106.53	110.58
3	A	502	ACP	N3-C2-N1	-3.51	123.27	128.58
3	A	502	ACP	C5-N7-C8	3.47	108.90	103.45
3	A	502	ACP	O2B-PB-C3B	3.46	121.04	106.73
2	A	501	YQY	F22-C10-C9	-3.45	113.65	118.32
2	A	501	YQY	C4-C5-N7	3.16	127.64	121.32
2	A	501	YQY	C3-C2-C1	-3.12	117.05	121.08
3	A	502	ACP	O1G-PG-C3B	-2.90	105.04	111.37
3	A	502	ACP	C2'-C1'-N9	-2.85	106.23	113.30
2	A	501	YQY	O18-C16-N17	-2.78	117.97	122.35
3	A	502	ACP	N9-C8-N7	-2.66	110.17	113.94
2	A	501	YQY	F22-C10-C11	2.50	124.46	118.65
3	A	502	ACP	O3'-C3'-C2'	-2.40	104.11	111.82
2	A	501	YQY	C1-C6-C5	2.40	125.88	123.43
3	A	502	ACP	O4'-C1'-N9	2.14	112.20	108.09
2	A	501	YQY	F26-C9-C10	-2.04	115.01	119.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	ACP	C4-N9-C8	2.03	107.87	105.74

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	ACP	PB-C3B-PG-O1G
2	A	501	YQY	C12-C13-C16-N17
2	A	501	YQY	C12-C13-C16-O18

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	YQY	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Continued on next page...

Continued from previous page...

Mol	Chain	Number of breaks
-----	-------	------------------

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	262:PRO	C	305:ARG	N	1.65

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/301 (93%)	0.74	27 (9%) 13 11	30, 57, 80, 96	8 (2%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	58	MET	8.4
1	A	353	LYS	5.3
1	A	104	LYS	5.2
1	A	73	GLU	5.0
1	A	113	ARG	4.8
1	A	70	LYS	4.7
1	A	224	VAL	4.2
1	A	324	LYS	3.8
1	A	379	ILE	3.3
1	A	216	ILE	3.2
1	A	381	LEU	3.2
1	A	240	TYR	3.0
1	A	316	TYR	2.9
1	A	380	GLY	2.8
1	A	136	ASP	2.7
1	A	101	LEU	2.5
1	A	82	VAL	2.5
1	A	225	GLY	2.5
1	A	243	GLN	2.4
1	A	107	ILE	2.3
1	A	223	PHE	2.3
1	A	242	VAL	2.3
1	A	239	HIS	2.2
1	A	59	ALA	2.2
1	A	229	TYR	2.1
1	A	122	ASN	2.1
1	A	311	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ACP	A	502	31/31	0.94	0.10	45,52,62,63	0
4	MG	A	503	1/1	0.95	0.05	58,58,58,58	0
2	YQY	A	501	26/26	0.98	0.07	40,45,51,55	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around YQY A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.