



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:59 AM UTC

PDB ID : 4ANL / pdb_00004anl
Title : Structure of G1269A Mutant Anaplastic Lymphoma Kinase
Authors : McTigue, M.; Deng, Y.; Liu, W.; Brooun, A.
Deposited on : 2012-03-20
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	12% 71% 15% 14%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ALK TYROSINE KINASE RECEPTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	1
			2305	1473	391	420	21			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1070	MET	-	expression tag	UNP Q9UM73
A	1071	ALA	-	expression tag	UNP Q9UM73
A	1072	HIS	-	expression tag	UNP Q9UM73
A	1073	HIS	-	expression tag	UNP Q9UM73
A	1074	HIS	-	expression tag	UNP Q9UM73
A	1075	HIS	-	expression tag	UNP Q9UM73
A	1076	HIS	-	expression tag	UNP Q9UM73
A	1077	HIS	-	expression tag	UNP Q9UM73
A	1078	ASP	-	expression tag	UNP Q9UM73
A	1079	TYR	-	expression tag	UNP Q9UM73
A	1080	GLY	-	expression tag	UNP Q9UM73
A	1081	ILE	-	expression tag	UNP Q9UM73
A	1082	PRO	-	expression tag	UNP Q9UM73
A	1083	THR	-	expression tag	UNP Q9UM73
A	1084	THR	-	expression tag	UNP Q9UM73
A	1085	GLU	-	expression tag	UNP Q9UM73
A	1086	ASN	-	expression tag	UNP Q9UM73
A	1087	LEU	-	expression tag	UNP Q9UM73
A	1088	TYR	-	expression tag	UNP Q9UM73
A	1089	PHE	-	expression tag	UNP Q9UM73
A	1090	GLN	-	expression tag	UNP Q9UM73
A	1091	GLY	-	expression tag	UNP Q9UM73
A	1092	SER	-	expression tag	UNP Q9UM73
A	1269	ALA	GLY	engineered mutation	UNP Q9UM73

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	274	Total 274	O 274	0	0

- Molecule 1: ALK TYROSINE KINASE RECEPTOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.84Å 57.59Å 105.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	52.58 – 1.70 52.58 – 1.70	Depositor EDS
% Data completeness (in resolution range)	99.3 (52.58-1.70) 99.3 (52.58-1.70)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 1.70Å)	Xtriage
Refinement program	CNX 2005	Depositor
R, R_{free}	0.202 , 0.216 0.200 , 0.208	Depositor DCC
R_{free} test set	1059 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.4	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 41.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2579	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	1/2360 (0.0%)	0.65	2/3197 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1401	TYR	C-N	-5.64	1.25	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1261	GLY	CA-C-N	6.47	126.97	119.47
1	A	1261	GLY	C-N-CA	6.47	126.97	119.47

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2305	0	2290	37	0
2	A	274	0	0	4	0
All	All	2579	0	2290	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1119:ILE:HD11	1:A:1134:GLN:HG3	1.59	0.84
1:A:1279:ARG:HA	1:A:1279:ARG:HE	1.56	0.71
1:A:1135:VAL:HG13	1:A:1145:LEU:HB3	1.75	0.69
1:A:1211:THR:HG23	1:A:1221:LEU:HD22	1.75	0.68
1:A:1383:ILE:O	1:A:1387:THR:HG23	1.97	0.64
1:A:1119:ILE:HD11	1:A:1134:GLN:CG	2.28	0.63
1:A:1367:GLN:HB2	1:A:1373:ARG:HG2	1.81	0.62
1:A:1387:THR:HG21	2:A:2133:HOH:O	2.00	0.62
1:A:1207:PHE:O	1:A:1211:THR:HG22	1.99	0.61
1:A:1231:ARG:HH11	1:A:1231:ARG:HG2	1.64	0.61
1:A:1248:ARG:CZ	1:A:1277:ILE:HG21	2.31	0.60
1:A:1365:CYS:O	1:A:1373:ARG:HD2	2.01	0.60
1:A:1135:VAL:CG1	1:A:1145:LEU:HB3	2.30	0.60
1:A:1180:VAL:HG13	1:A:1197:GLU:HB3	1.84	0.59
1:A:1214:ARG:HB3	1:A:1215:PRO:HD2	1.85	0.58
1:A:1298:PRO:HA	1:A:1301:PHE:CZ	2.38	0.58
1:A:1100:GLY:O	1:A:1101:LYS:HD2	2.09	0.51
1:A:1203:ASP:HA	1:A:1256:LEU:HD23	1.92	0.51
1:A:1247:HIS:O	1:A:1248:ARG:HB2	2.11	0.51
1:A:1118:LEU:N	1:A:1118:LEU:HD12	2.26	0.50
1:A:1244:HIS:HD2	2:A:2015:HOH:O	1.95	0.49
1:A:1241:GLU:HG3	1:A:1308:SER:OG	2.14	0.48
1:A:1293:VAL:HG21	1:A:1335:ASN:CB	2.44	0.48
1:A:1368:HIS:HD2	2:A:2226:HOH:O	1.97	0.48
1:A:1146:GLN:HG2	2:A:2072:HOH:O	2.12	0.47
1:A:1279:ARG:HA	1:A:1279:ARG:NE	2.28	0.46
1:A:1109:LYS:HE3	1:A:1111:VAL:HG22	1.98	0.46
1:A:1113:ARG:HA	1:A:1113:ARG:HD3	1.80	0.46
1:A:1339:LEU:C	1:A:1339:LEU:HD23	2.43	0.43
1:A:1113:ARG:NH2	1:A:1191:PRO:O	2.52	0.43
1:A:1152:LEU:HD12	1:A:1153:PRO:HD2	2.00	0.43
1:A:1293:VAL:HG21	1:A:1335:ASN:HB3	2.00	0.43
1:A:1306:PHE:HD2	1:A:1310:THR:HG21	1.84	0.42
1:A:1101:LYS:HA	1:A:1101:LYS:CE	2.47	0.42
1:A:1294:LYS:HG2	1:A:1338:VAL:HG21	2.01	0.42
1:A:1216:SER:C	1:A:1218:PRO:HD3	2.46	0.41
1:A:1183:ILE:HD12	1:A:1183:ILE:N	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	285/342 (83%)	282 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	255/298 (86%)	254 (100%)	1 (0%)	84	80

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1221	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1115	ASN
1	A	1188	GLN
1	A	1243	ASN
1	A	1368	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	293/342 (85%)	0.60	40 (13%) 6 6	14, 24, 44, 60	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1126	ALA	6.0
1	A	1127	PHE	5.7
1	A	1286	GLY	5.5
1	A	1287	GLY	4.2
1	A	1301	PHE	4.2
1	A	1128	GLY	3.8
1	A	1135	VAL	3.7
1	A	1216	SER	3.6
1	A	1123	GLY	3.5
1	A	1302	MET	3.5
1	A	1288	CYS	3.4
1	A	1279	ARG	3.4
1	A	1217	GLN	3.2
1	A	1101	LYS	3.1
1	A	1215	PRO	3.1
1	A	1278	TYR	3.1
1	A	1144	PRO	3.1
1	A	1402	GLY	3.0
1	A	1277	ILE	3.0
1	A	1214	ARG	2.7
1	A	1094	PRO	2.7
1	A	1211	THR	2.6
1	A	1308	SER	2.5
1	A	1218	PRO	2.5
1	A	1300	ALA	2.5
1	A	1100	GLY	2.3
1	A	1262	PRO	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1306	PHE	2.3
1	A	1401	TYR	2.3
1	A	1384	GLU	2.2
1	A	1097	CYS	2.2
1	A	1122	LEU	2.2
1	A	1129	GLU	2.1
1	A	1118	LEU	2.1
1	A	1364	GLN	2.1
1	A	1158	GLU	2.1
1	A	1130	VAL	2.0
1	A	1155	VAL	2.0
1	A	1332	SER	2.0
1	A	1145	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.