



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:00 PM UTC

PDB ID : 7ANA / pdb_00007ana
Title : A single sulfatase is required for metabolism of colonic mucin O-glycans and intestinal colonization by a symbiotic human gut bacterium (BT1622-S1_20)
Authors : Sofia de Jesus Vaz Luis, A.; Basle, A.; Martens, E.C.; Cartmell, A.
Deposited on : 2020-10-11
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	503	11% 88% 6% 6%
1	BBB	503	3% 89% 5% 6%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 7799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-acetylgalactosamine-6-sulfatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	475	Total	C	N	O	S	0	0	0
			3785	2412	646	705	22			
1	BBB	475	Total	C	N	O	S	0	1	0
			3796	2418	650	706	22			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-2	MET	-	initiating methionine	UNP Q8A7A3
AAA	-1	GLY	-	expression tag	UNP Q8A7A3
AAA	0	SER	-	expression tag	UNP Q8A7A3
AAA	1	SER	-	expression tag	UNP Q8A7A3
AAA	2	HIS	-	expression tag	UNP Q8A7A3
AAA	3	HIS	-	expression tag	UNP Q8A7A3
AAA	4	HIS	-	expression tag	UNP Q8A7A3
AAA	5	HIS	-	expression tag	UNP Q8A7A3
AAA	6	HIS	-	expression tag	UNP Q8A7A3
AAA	7	HIS	-	expression tag	UNP Q8A7A3
AAA	8	SER	-	expression tag	UNP Q8A7A3
AAA	9	SER	-	expression tag	UNP Q8A7A3
AAA	10	GLY	-	expression tag	UNP Q8A7A3
AAA	11	LEU	-	expression tag	UNP Q8A7A3
AAA	12	VAL	-	expression tag	UNP Q8A7A3
AAA	13	PRO	-	expression tag	UNP Q8A7A3
AAA	14	ARG	-	expression tag	UNP Q8A7A3
AAA	15	GLY	-	expression tag	UNP Q8A7A3
AAA	16	SER	-	expression tag	UNP Q8A7A3
AAA	17	HIS	-	expression tag	UNP Q8A7A3
AAA	18	MET	-	expression tag	UNP Q8A7A3
AAA	19	ALA	-	expression tag	UNP Q8A7A3
AAA	20	SER	-	expression tag	UNP Q8A7A3
BBB	-2	MET	-	initiating methionine	UNP Q8A7A3
BBB	-1	GLY	-	expression tag	UNP Q8A7A3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
BBB	0	SER	-	expression tag	UNP Q8A7A3
BBB	1	SER	-	expression tag	UNP Q8A7A3
BBB	2	HIS	-	expression tag	UNP Q8A7A3
BBB	3	HIS	-	expression tag	UNP Q8A7A3
BBB	4	HIS	-	expression tag	UNP Q8A7A3
BBB	5	HIS	-	expression tag	UNP Q8A7A3
BBB	6	HIS	-	expression tag	UNP Q8A7A3
BBB	7	HIS	-	expression tag	UNP Q8A7A3
BBB	8	SER	-	expression tag	UNP Q8A7A3
BBB	9	SER	-	expression tag	UNP Q8A7A3
BBB	10	GLY	-	expression tag	UNP Q8A7A3
BBB	11	LEU	-	expression tag	UNP Q8A7A3
BBB	12	VAL	-	expression tag	UNP Q8A7A3
BBB	13	PRO	-	expression tag	UNP Q8A7A3
BBB	14	ARG	-	expression tag	UNP Q8A7A3
BBB	15	GLY	-	expression tag	UNP Q8A7A3
BBB	16	SER	-	expression tag	UNP Q8A7A3
BBB	17	HIS	-	expression tag	UNP Q8A7A3
BBB	18	MET	-	expression tag	UNP Q8A7A3
BBB	19	ALA	-	expression tag	UNP Q8A7A3
BBB	20	SER	-	expression tag	UNP Q8A7A3

- Molecule 2 is 2-acetamido-2-deoxy- α -D-galactopyranose (CCD ID: A2G) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	0	0
			15	8	1	6		
2	BBB	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	1	Total	Ca	0	0
			1	1		
3	BBB	1	Total	Ca	0	0
			1	1		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-galactopyranose (CCD ID: NGA) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	BBB	1	Total	C	N	O	0	0
			15	8	1	6		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	BBB	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	52	Total	O	0	0
			52	52		
6	BBB	115	Total	O	0	0
			115	115		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-acetylgalactosamine-6-sulfatase



- Molecule 1: N-acetylgalactosamine-6-sulfatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	83.25Å 83.25Å 390.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	70.22 – 2.30 70.22 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (70.22-2.30) 99.8 (70.22-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.211 , 0.235 0.216 , 0.235	Depositor DCC
R_{free} test set	3083 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtriage
Anisotropy	0.515	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7799	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A2G, EDO, NGA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	AAA	0.98	0/3899	1.32	3/5289 (0.1%)
1	BBB	0.96	0/3910	1.33	2/5303 (0.0%)
All	All	0.97	0/7809	1.32	5/10592 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	389	GLN	CB-CA-C	-8.21	96.72	110.68
1	AAA	496	ASN	CB-CA-C	-6.15	99.42	110.36
1	BBB	496	ASN	CB-CA-C	-6.05	99.59	110.36
1	AAA	42	ILE	CA-C-N	5.34	126.03	119.99
1	AAA	42	ILE	C-N-CA	5.34	126.03	119.99

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3785	0	3612	14	0
1	BBB	3796	0	3624	13	0
2	AAA	15	0	12	0	0
2	BBB	15	0	12	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	AAA	1	0	0	0	0
3	BBB	1	0	0	0	0
4	BBB	15	0	15	1	0
5	BBB	4	0	6	1	0
6	AAA	52	0	0	0	0
6	BBB	115	0	0	1	0
All	All	7799	0	7281	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 28 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:BBB:133:MET:HE3	1:BBB:134:MET:HE3	1.77	0.66
5:BBB:602:EDO:H21	6:BBB:718:HOH:O	1.97	0.65
1:AAA:133:MET:HE3	1:AAA:134:MET:HE3	1.82	0.61
1:AAA:127:THR:O	1:AAA:132:ARG:NH1	2.41	0.54
1:BBB:127:THR:O	1:BBB:132:ARG:NH1	2.41	0.53

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	473/503 (94%)	457 (97%)	16 (3%)	0	100	100
1	BBB	474/503 (94%)	459 (97%)	15 (3%)	0	100	100
All	All	947/1006 (94%)	916 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	400/425 (94%)	395 (99%)	5 (1%)	61	77
1	BBB	401/425 (94%)	396 (99%)	5 (1%)	63	79
All	All	801/850 (94%)	791 (99%)	10 (1%)	61	79

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	BBB	329	ASN
1	BBB	375	LYS
1	BBB	424	GLU
1	AAA	401	ILE
1	AAA	492	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	A2G	AAA	701	-	15,15,15	0.46	0	21,21,21	1.02	1 (4%)
5	EDO	BBB	602	-	3,3,3	0.30	0	2,2,2	0.37	0
2	A2G	BBB	603	-	15,15,15	0.76	1 (6%)	21,21,21	0.97	1 (4%)
4	NGA	BBB	601	-	15,15,15	0.45	0	21,21,21	1.38	3 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	A2G	AAA	701	-	-	2/6/26/26	0/1/1/1
5	EDO	BBB	602	-	-	0/1/1/1	-
2	A2G	BBB	603	-	-	1/6/26/26	0/1/1/1
4	NGA	BBB	601	-	-	3/6/26/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	603	A2G	C1-C2	2.23	1.55	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	701	A2G	O5-C1-C2	3.40	112.94	109.52
4	BBB	601	NGA	C8-C7-N2	3.23	121.47	116.12
4	BBB	601	NGA	O7-C7-N2	-2.82	117.00	121.98
2	BBB	603	A2G	C1-C2-N2	-2.12	108.27	110.73
4	BBB	601	NGA	O5-C1-C2	2.07	111.59	109.52

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	BBB	601	NGA	C8-C7-N2-C2
4	BBB	601	NGA	O7-C7-N2-C2
2	AAA	701	A2G	O5-C5-C6-O6
2	BBB	603	A2G	O5-C5-C6-O6
2	AAA	701	A2G	C4-C5-C6-O6


There are no ring outliers.

3 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	BBB	602	EDO	1	0
2	BBB	603	A2G	1	0
4	BBB	601	NGA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand A2G AAA 701



Ligand A2G BBB 603



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	475/503 (94%)	0.87	57 (12%) 9 9	45, 68, 120, 143	0
1	BBB	475/503 (94%)	0.13	13 (2%) 56 58	28, 48, 69, 92	1 (0%)
All	All	950/1006 (94%)	0.50	70 (7%) 20 22	28, 55, 109, 143	1 (0%)

The worst 5 of 70 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	420	GLY	4.3
1	AAA	416	LEU	4.1
1	AAA	421	LYS	3.9
1	AAA	401	ILE	3.7
1	AAA	263	ARG	3.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	BBB	602	4/4	0.82	0.25	56,60,60,62	0
4	NGA	BBB	601	15/15	0.88	0.11	62,63,66,66	0
2	A2G	AAA	701	15/15	0.94	0.07	46,47,48,48	0
2	A2G	BBB	603	15/15	0.94	0.08	44,45,47,48	0
3	CA	AAA	702	1/1	0.97	0.16	69,69,69,69	0
3	CA	BBB	604	1/1	0.97	0.15	64,64,64,64	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A2G AAA 701:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A2G BBB 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.