



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 07:10 PM UTC

PDB ID : 7AO0 / pdb_00007ao0
Title : Crystal structure of CotB2 variant F107A in complex with alendronate
Authors : Dimos, N.; Driller, R.; Loll, B.
Deposited on : 2020-10-13
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	318	4% 86% 7% 7%
1	B	318	5% 82% 10% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cyclooctat-9-en-7-ol synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	295	Total	C	N	O	S	0	12	0
			2522	1595	438	472	17			
1	B	296	Total	C	N	O	S	0	5	0
			2460	1564	415	464	17			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	107	ALA	PHE	engineered mutation	UNP C9K1X5
A	308	ALA	-	expression tag	UNP C9K1X5
A	309	ALA	-	expression tag	UNP C9K1X5
A	310	ALA	-	expression tag	UNP C9K1X5
A	311	LEU	-	expression tag	UNP C9K1X5
A	312	GLU	-	expression tag	UNP C9K1X5
A	313	HIS	-	expression tag	UNP C9K1X5
A	314	HIS	-	expression tag	UNP C9K1X5
A	315	HIS	-	expression tag	UNP C9K1X5
A	316	HIS	-	expression tag	UNP C9K1X5
A	317	HIS	-	expression tag	UNP C9K1X5
A	318	HIS	-	expression tag	UNP C9K1X5
B	107	ALA	PHE	engineered mutation	UNP C9K1X5
B	308	ALA	-	expression tag	UNP C9K1X5
B	309	ALA	-	expression tag	UNP C9K1X5
B	310	ALA	-	expression tag	UNP C9K1X5
B	311	LEU	-	expression tag	UNP C9K1X5
B	312	GLU	-	expression tag	UNP C9K1X5
B	313	HIS	-	expression tag	UNP C9K1X5
B	314	HIS	-	expression tag	UNP C9K1X5
B	315	HIS	-	expression tag	UNP C9K1X5
B	316	HIS	-	expression tag	UNP C9K1X5
B	317	HIS	-	expression tag	UNP C9K1X5
B	318	HIS	-	expression tag	UNP C9K1X5

- Molecule 2 is 4-AMINO-1-HYDROXYBUTANE-1,1-DIYLDIPHOSPHONATE (CCD ID: AHD) (formula: $C_4H_9NO_7P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			14	4	1	7	2		
2	B	1	Total	C	N	O	P	0	0
			14	4	1	7	2		

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0
3	B	1	Total C O 8 6 2	0	0

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total Mg 3 3	0	0
4	B	3	Total Mg 3 3	0	0

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Cl 1 1	0	0
5	B	1	Total Cl 1 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	229	Total O 229 229	0	8
6	B	226	Total O 226 226	0	7


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cyclooctat-9-en-7-ol synthase



• Molecule 1: Cyclooctat-9-en-7-ol synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.81Å 98.05Å 106.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.02 – 1.93 49.02 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.3 (49.02-1.93) 99.6 (49.02-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 1.92Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.181 , 0.238 0.179 , 0.236	Depositor DCC
R_{free} test set	2100 reflections (4.16%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.633	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5505	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 60.55 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4957e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, CL, AHD, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2584	0.56	0/3499
1	B	0.39	0/2522	0.57	0/3420
All	All	0.39	0/5106	0.56	0/6919

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2522	0	2430	17	0
1	B	2460	0	2372	21	0
2	A	14	0	9	1	0
2	B	14	0	9	0	0
3	A	8	0	14	1	0
3	B	24	0	42	1	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	229	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	226	0	0	1	0
All	All	5505	0	4876	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:297:THR:HG22	1:B:300:ASN:HD22	1.43	0.81
1:A:294:ARG:HG3	1:A:303:ASN:HB3	1.62	0.80
1:A:227:ARG:NH1	6:A:501:HOH:O	2.13	0.79
1:A:28[B]:ARG:HH12	1:A:250:LYS:HD2	1.61	0.65
1:A:24[B]:GLU:CD	1:A:27[B]:ARG:HH11	2.05	0.64
1:B:112:MET:HE3	1:B:116:LEU:HB2	1.79	0.63
1:B:294:ARG:HG3	1:B:303:ASN:CB	2.36	0.55
1:A:294:ARG:O	1:A:303:ASN:ND2	2.38	0.54
1:B:137[A]:GLU:H	1:B:137[A]:GLU:CD	2.17	0.52
1:B:128:LYS:NZ	6:B:503:HOH:O	2.34	0.52
1:A:293:LYS:HG3	1:A:302:VAL:HG21	1.92	0.50
1:A:40:ARG:NH1	6:A:505:HOH:O	2.44	0.50
1:B:116:LEU:HD11	1:B:307:GLN:C	2.37	0.49
1:B:294:ARG:HG3	1:B:303:ASN:HB3	1.95	0.49
1:A:58:CYS:HB3	1:A:74:TRP:CH2	2.48	0.49
1:A:24[B]:GLU:OE2	1:A:27[B]:ARG:NH1	2.36	0.49
1:A:177:ARG:O	1:A:181:ILE:HG22	2.12	0.49
1:A:196:HIS:CD2	1:A:273:LEU:HD22	2.48	0.49
1:A:98:ALA:HB2	1:A:133:TYR:CB	2.43	0.48
1:B:191:TYR:CG	1:B:192:PRO:HD3	2.49	0.48
1:B:227:ARG:HD3	1:B:303:ASN:ND2	2.29	0.47
1:B:292:ASN:ND2	1:B:294:ARG:HB3	2.29	0.47
2:A:401:AHD:HN41	3:A:402:MPD:H11	1.81	0.46
1:B:42:THR:HB	1:B:87:PRO:HG2	1.98	0.46
1:A:178:VAL:HG21	3:B:402:MPD:HM2	1.98	0.45
1:A:246:GLU:O	1:A:250:LYS:HG2	2.17	0.45
1:A:191:TYR:CG	1:A:192:PRO:HD3	2.53	0.44
1:A:195[A]:ARG:NH1	6:A:502:HOH:O	2.34	0.44
1:B:112:MET:CE	1:B:116:LEU:HB2	2.44	0.44
1:B:184:ASP:HA	1:B:209:ALA:HB2	2.01	0.42
1:B:145:ALA:HB1	1:B:185:PHE:HA	2.00	0.42
1:B:76:SER:O	1:B:288[B]:TRP:HZ2	2.03	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:173:TYR:CE2	1:B:237:CYS:HB3	2.55	0.42
1:B:196:HIS:CD2	1:B:273:LEU:HD22	2.55	0.42
1:B:41:PRO:HD3	1:B:283:TYR:CD2	2.56	0.41
1:A:196:HIS:CG	1:A:273:LEU:HD22	2.56	0.41
1:B:246:GLU:O	1:B:250[A]:LYS:HG2	2.21	0.40
1:B:58:CYS:HB3	1:B:74:TRP:CH2	2.56	0.40
1:B:98:ALA:HB2	1:B:133:TYR:CB	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	305/318 (96%)	302 (99%)	3 (1%)	0	100	100
1	B	299/318 (94%)	296 (99%)	3 (1%)	0	100	100
All	All	604/636 (95%)	598 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/280 (98%)	275 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	270/280 (96%)	266 (98%)	4 (2%)	57	49
All	All	545/560 (97%)	541 (99%)	4 (1%)	78	74

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	88	ARG
1	B	116	LEU
1	B	137[A]	GLU
1	B	137[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	233	GLN
1	A	285	ASN
1	B	125	GLN
1	B	285	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 8 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MPD	A	402	-	7,7,7	0.40	0	9,10,10	0.46	0
3	MPD	B	401	-	7,7,7	0.38	0	9,10,10	0.43	0
2	AHD	B	407	4	13,13,13	1.34	1 (7%)	19,21,21	1.49	2 (10%)
3	MPD	B	408	-	7,7,7	0.32	0	9,10,10	0.47	0
3	MPD	B	402	-	7,7,7	0.43	0	9,10,10	0.76	0
2	AHD	A	401	4	13,13,13	1.52	2 (15%)	19,21,21	1.88	4 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	402	-	-	2/5/5/5	-
3	MPD	B	401	-	-	5/5/5/5	-
2	AHD	B	407	4	-	4/23/23/23	-
3	MPD	B	408	-	-	0/5/5/5	-
3	MPD	B	402	-	-	4/5/5/5	-
2	AHD	A	401	4	-	3/23/23/23	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	AHD	O13-C8	-4.10	1.39	1.44
2	B	407	AHD	O13-C8	-2.91	1.41	1.44
2	A	401	AHD	P14-O16	-2.33	1.48	1.54

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	AHD	C2-C7-C8	-5.99	105.61	116.02
2	B	407	AHD	C2-C7-C8	-4.68	107.88	116.02
2	A	401	AHD	O10-P9-C8	2.70	112.12	106.19
2	B	407	AHD	O16-P14-C8	2.37	111.41	106.19
2	A	401	AHD	O12-P9-C8	2.23	111.09	106.19
2	A	401	AHD	O11-P9-C8	-2.04	104.59	109.82

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	407	AHD	C2-C7-C8-P9
2	B	407	AHD	C2-C7-C8-P14
2	B	407	AHD	C2-C7-C8-O13
3	B	401	MPD	C2-C3-C4-O4
3	B	401	MPD	C2-C3-C4-C5
3	B	402	MPD	C1-C2-C3-C4
3	B	402	MPD	O2-C2-C3-C4
3	B	402	MPD	C2-C3-C4-C5
2	A	401	AHD	C2-C7-C8-P9
2	A	401	AHD	C2-C7-C8-P14
2	A	401	AHD	C2-C7-C8-O13
3	B	401	MPD	C1-C2-C3-C4
3	B	401	MPD	CM-C2-C3-C4
2	B	407	AHD	C3-C2-C7-C8
3	A	402	MPD	O2-C2-C3-C4
3	B	401	MPD	O2-C2-C3-C4
3	A	402	MPD	C1-C2-C3-C4
3	B	402	MPD	CM-C2-C3-C4

There are no ring outliers.

3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	402	MPD	1	0
3	B	402	MPD	1	0
2	A	401	AHD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	295/318 (92%)	-0.20	14 (4%) 36 41	7, 16, 43, 86	12 (4%)
1	B	296/318 (93%)	-0.15	16 (5%) 31 35	7, 17, 46, 92	5 (1%)
All	All	591/636 (92%)	-0.17	30 (5%) 33 38	7, 17, 46, 92	17 (2%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	298	ALA	5.9
1	B	302	VAL	5.6
1	A	308	ALA	5.6
1	B	303	ASN	5.4
1	A	306	ILE	5.2
1	A	297	THR	5.2
1	A	302	VAL	5.1
1	B	298	ALA	4.9
1	A	299	VAL	4.9
1	B	306	ILE	4.7
1	A	303	ASN	4.7
1	B	299	VAL	4.5
1	B	297	THR	4.3
1	B	13	ILE	4.1
1	B	304	SER	3.9
1	B	300	ASN	3.8
1	B	307	GLN	3.8
1	A	304	SER	3.5
1	A	305	ARG	3.3
1	A	307	GLN	2.9
1	A	14	GLY	2.9
1	B	12	ASP	2.7
1	B	305	ARG	2.6
1	B	301	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	301	ASP	2.4
1	B	14	GLY	2.4
1	A	300	ASN	2.3
1	A	67[A]	LYS	2.2
1	B	227	ARG	2.2
1	B	73	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MPD	B	401	8/8	0.84	0.14	25,40,48,50	0
3	MPD	A	402	8/8	0.85	0.12	15,27,28,37	0
3	MPD	B	402	8/8	0.86	0.15	17,33,50,63	0
3	MPD	B	408	8/8	0.91	0.08	14,21,27,33	0
4	MG	A	403	1/1	0.98	0.03	15,15,15,15	0
4	MG	A	405	1/1	0.98	0.03	19,19,19,19	0
4	MG	B	403	1/1	0.98	0.02	16,16,16,16	0
4	MG	B	404	1/1	0.98	0.04	18,18,18,18	0
2	AHD	B	407	14/14	0.99	0.05	11,15,31,34	0
2	AHD	A	401	14/14	0.99	0.04	10,15,33,34	0
4	MG	A	404	1/1	0.99	0.03	16,16,16,16	0
5	CL	A	406	1/1	0.99	0.07	21,21,21,21	0
5	CL	B	406	1/1	0.99	0.10	19,19,19,19	0
4	MG	B	405	1/1	1.00	0.01	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.