



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 10:33 AM UTC

PDB ID : 3APR / pdb_00003apr
Title : BINDING OF A REDUCED PEPTIDE INHIBITOR TO THE ASPARTIC
PROTEINASE FROM RHIZOPUS CHINENSIS. IMPLICATIONS FOR A
MECHANISM OF ACTION
Authors : Suguna, K.; Davies, D.R.
Deposited on : 1987-06-22
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	325	 88% 11% .
2	I	7	 43% 29% 14% 14%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2818 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RHIZOPUSPEPSIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	325	Total	C	N	O	S	0	4	0
			2417	1524	399	488	6			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	15	ILE	VAL	conflict	UNP P06026
E	54	GLY	ARG	conflict	UNP P06026
E	61	ASN	LYS	conflict	UNP P06026
E	116	SER	ASN	conflict	UNP P06026
E	162	LYS	SER	conflict	UNP P06026
E	230	ILE	VAL	conflict	UNP P06026
E	256	ALA	ARG	conflict	UNP P06026
E	281	PHE	TYR	conflict	UNP P06026
E	294	TRP	PHE	conflict	UNP P06026
E	295	GLY	ASP	conflict	UNP P06026

- Molecule 2 is a protein called REDUCED PEPTIDE INHIBITOR.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	I	6	Total	C	N	O	0	0	1
			57	43	9	5			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	338	Total	O	0	0
			338	338		
3	I	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RHIZOPUSPEPSIN



- Molecule 2: REDUCED PEPTIDE INHIBITOR



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.37Å 60.61Å 106.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.80 53.38 – 1.80	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-1.80) 80.4 (53.38-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	PROLSQ	Depositor
R, R_{free}	0.147 , (Not available) 0.194 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	14.0	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 68.3	EDS
L-test for twinning ¹	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.036 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2818	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PUK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	E	0.97	0/2488	1.66	26/3381 (0.8%)
2	I	0.82	0/37	1.47	0/48
All	All	0.97	0/2525	1.65	26/3429 (0.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	I	0	3

There are no bond length outliers.

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	325	GLU	CB-CG-CD	10.20	129.94	112.60
1	E	141	ASP	CA-CB-CG	6.88	119.48	112.60
1	E	108	LYS	N-CA-C	-6.61	105.75	113.88
1	E	52	GLY	CA-C-O	-6.53	115.63	121.64
1	E	294	TRP	N-CA-CB	6.27	120.50	110.85
1	E	35	ASP	N-CA-C	6.25	118.96	108.02
1	E	271	VAL	N-CA-C	-6.08	98.37	107.37
1	E	120	ASP	N-CA-C	-6.01	105.98	113.38
1	E	62	GLN	CG-CD-NE2	-5.99	107.42	116.40
1	E	257	PHE	CA-CB-CG	5.98	119.78	113.80
1	E	269	PHE	CA-C-N	-5.57	113.46	122.82
1	E	269	PHE	C-N-CA	-5.57	113.46	122.82
1	E	284	GLN	OE1-CD-NE2	-5.56	117.04	122.60
1	E	134	ARG	CD-NE-CZ	-5.48	116.72	124.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	40	ASP	CA-CB-CG	5.46	118.06	112.60
1	E	156	VAL	N-CA-C	5.45	115.71	107.75
1	E	324	ALA	N-CA-C	-5.40	100.98	109.25
1	E	262	PHE	CA-CB-CG	5.40	119.20	113.80
1	E	131	THR	N-CA-C	-5.40	103.01	110.35
1	E	286	ILE	CA-C-O	-5.39	114.80	120.57
1	E	225	ILE	CB-CG1-CD1	-5.29	102.70	113.80
1	E	323	VAL	CB-CA-C	-5.27	104.00	111.28
1	E	154	PHE	CA-CB-CG	5.25	119.05	113.80
1	E	20	GLN	OE1-CD-NE2	5.14	127.75	122.60
1	E	167	GLY	N-CA-C	-5.10	104.31	111.09
1	E	319	GLN	CA-C-O	-5.06	114.71	120.32

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	I	5	PUK	Mainchain,Peptide
2	I	6	VAL	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	2417	0	2331	8	0
2	I	57	0	46	2	0
3	E	338	0	0	1	0
3	I	6	0	0	1	0
All	All	2818	0	2377	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:20:GLN:NE2	1:E:29:LYS:HD3	2.10	0.66
1:E:20:GLN:HE22	1:E:29:LYS:HD3	1.60	0.65
1:E:162:LYS:HG3	1:E:163:ASN:ND2	2.27	0.48
1:E:256:ALA:HB2	3:E:664:HOH:O	2.13	0.48
1:E:294:TRP:CZ2	2:I:5:PUK:H20	2.49	0.47
1:E:292:GLY:HA3	1:E:294:TRP:CZ3	2.51	0.46
2:I:2:PRO:N	3:I:876:HOH:O	2.51	0.44
1:E:226:LEU:HA	1:E:227:PRO:HD3	1.92	0.43
1:E:176:SER:HA	1:E:179:PHE:CE1	2.54	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	327/325 (101%)	323 (99%)	4 (1%)	0	100	100
2	I	3/7 (43%)	2 (67%)	1 (33%)	0	100	100
All	All	330/332 (99%)	325 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	259/255 (102%)	253 (98%)	6 (2%)	44	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	I	4/5 (80%)	4 (100%)	0	100	100
All	All	263/260 (101%)	257 (98%)	6 (2%)	44	33

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	41	LEU
1	E	43	ILE
1	E	123	LEU
1	E	241	SER
1	E	293	ASN
1	E	323	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	13	ASN
1	E	20	GLN
1	E	163	ASN
1	E	243	ASN
1	E	284	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PUK	I	5	2	21,22,23	0.74	0	23,27,29	1.23	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PUK	I	5	2	-	3/14/15/17	0/2/2/2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	5	PUK	C4-C3-CA	2.74	118.51	113.23
2	I	5	PUK	C8-C9-C7	2.12	122.78	119.87
2	I	5	PUK	C18-C17-C15	-2.10	117.65	120.24

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	5	PUK	CA-C2-N2-C10
2	I	5	PUK	CA-C3-C4-C6
2	I	5	PUK	CA-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	I	5	PUK	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	325/325 (100%)	0.09	1 (0%) 90 90	8, 14, 25, 41	4 (1%)
2	I	5/7 (71%)	0.41	0 100 100	20, 21, 22, 23	0
All	All	330/332 (99%)	0.09	1 (0%) 90 90	8, 14, 25, 41	4 (1%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	294	TRP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PUK	I	5	21/22	0.84	0.09	16,17,20,20	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.