



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:07 AM UTC

PDB ID : 5AP0 / pdb_00005ap0
Title : Naturally Occurring Mutations in the MPS1 Gene Predispose Cells to Kinase Inhibitor Drug Resistance.
Authors : Gurden, M.D.; Westwood, I.M.; Faisal, A.; Naud, S.; Cheung, K.J.; McAndrew, C.; Wood, A.; Schmitt, J.; Boxall, K.; Mak, G.; Workman, P.; Burke, R.; Hoelder, S.; Blagg, J.; van Montfort, R.L.M.; Linardopoulos, S.
Deposited on : 2015-09-14
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	9% 72% 9% 18%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2219 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DUAL SPECIFICITY PROTEIN KINASE TTK.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	256	2025	1303	334	375	1	12	0	0	1

There are 23 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	496	MET	-	expression tag	UNP P33981
A	497	HIS	-	expression tag	UNP P33981
A	498	HIS	-	expression tag	UNP P33981
A	499	HIS	-	expression tag	UNP P33981
A	500	HIS	-	expression tag	UNP P33981
A	501	HIS	-	expression tag	UNP P33981
A	502	HIS	-	expression tag	UNP P33981
A	503	SER	-	expression tag	UNP P33981
A	504	SER	-	expression tag	UNP P33981
A	505	GLY	-	expression tag	UNP P33981
A	506	VAL	-	expression tag	UNP P33981
A	507	ASP	-	expression tag	UNP P33981
A	508	LEU	-	expression tag	UNP P33981
A	509	GLY	-	expression tag	UNP P33981
A	510	THR	-	expression tag	UNP P33981
A	511	GLU	-	expression tag	UNP P33981
A	512	ASN	-	expression tag	UNP P33981
A	513	LEU	-	expression tag	UNP P33981
A	514	TYR	-	expression tag	UNP P33981
A	515	PHE	-	expression tag	UNP P33981
A	516	GLN	-	expression tag	UNP P33981
A	517	SER	-	expression tag	UNP P33981
A	518	MET	-	expression tag	UNP P33981

- Molecule 2 is 9-CYCLOPENTYL-2-[[2-METHOXY-4-[(1-METHYLPYPERIDIN-4-YL)OXY]-PHENYL]AMINO]-7-METHYL-7,9-DIHYDRO-8H-PURIN-8-ONE (CCD ID: AU5) (formula: C₂₄H₃₂N₆O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	24	6	3		

- Molecule 3 is O-ACETALDEHYDYL-HEXAETHYLENE GLYCOL (CCD ID: P4C) (formula: $C_{14}H_{28}O_8$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			22	14	8		
3	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	59	Total O 59 59	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DUAL SPECIFICITY PROTEIN KINASE TTK



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	70.93Å 111.96Å 113.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.12 – 2.15 41.12 – 2.15	Depositor EDS
% Data completeness (in resolution range)	95.2 (41.12-2.15) 95.1 (41.12-2.15)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.34 (at 2.16Å)	Xtriage
Refinement program	BUSTER 2.11.4	Depositor
R, R_{free}	0.184 , 0.220 0.197 , 0.225	Depositor DCC
R_{free} test set	1209 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	55.0	Xtriage
Anisotropy	0.380	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 65.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2219	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, P4C, TPO, AU5, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	4/2058 (0.2%)	1.27	2/2792 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	794	GLN	C-N	-5.46	1.25	1.33
1	A	607	ILE	CA-CB	5.33	1.59	1.55
1	A	631	MET	SD-CE	-5.22	1.66	1.79
1	A	600	MET	SD-CE	-5.18	1.66	1.79

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	561	ASP	CA-CB-CG	5.08	117.69	112.60
1	A	544	ASN	CA-CB-CG	5.08	117.68	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2025	0	1945	14	0
2	A	33	0	32	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	38	0	49	5	0
4	A	12	0	18	1	0
5	A	52	0	78	5	0
6	A	59	0	0	0	0
All	All	2219	0	2122	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:606:ASN:HA	4:A:1798:DMS:H22	1.61	0.81
1:A:663:ILE:HD12	3:A:1796:P4C:H52	1.81	0.61
1:A:574:TYR:CD1	1:A:643:ILE:HD11	2.35	0.61
2:A:1795:AU5:O	3:A:1796:P4C:H51	2.00	0.60
1:A:571:GLU:HG2	3:A:1796:P4C:H112	1.86	0.58
1:A:568:TYR:CE1	3:A:1796:P4C:H151	2.40	0.57
5:A:1800:EDO:H12	5:A:1803:EDO:H12	1.88	0.55
1:A:639:HIS:HB3	5:A:1800:EDO:H11	1.90	0.52
1:A:586:ILE:HD11	2:A:1795:AU5:H103	1.92	0.52
1:A:528:LEU:HD11	1:A:543:LEU:HG	1.95	0.48
1:A:641:HIS:HB3	5:A:1810:EDO:H11	2.01	0.42
1:A:555:VAL:HG21	3:A:1796:P4C:H141	2.00	0.42
1:A:774:ARG:HB2	5:A:1805:EDO:H12	2.02	0.42
1:A:586:ILE:CD1	2:A:1795:AU5:H103	2.49	0.42
1:A:603:GLU:O	2:A:1795:AU5:H3	2.19	0.42
1:A:783:PRO:HG2	5:A:1803:EDO:H11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	249/313 (80%)	245 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/287 (74%)	202 (96%)	9 (4%)	26	24

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	521	LYS
1	A	523	ARG
1	A	537	SER
1	A	549	ILE
1	A	731	LYS
1	A	738	ILE
1	A	741	ILE
1	A	742	SER
1	A	794	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	548	GLN
1	A	640	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	686	1	8,10,11	0.90	0	10,14,16	1.77	3 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	686	1	-	4/9/11/13	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	686	TPO	P-OG1-CB	-3.55	113.69	123.33
1	A	686	TPO	O3P-P-OG1	2.72	116.46	105.85
1	A	686	TPO	O3P-P-O1P	-2.08	102.74	110.83

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	686	TPO	N-CA-CB-OG1
1	A	686	TPO	O-C-CA-CB
1	A	686	TPO	CB-OG1-P-O2P
1	A	686	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	1801	-	3,3,3	0.57	0	2,2,2	0.19	0
4	DMS	A	1799	-	3,3,3	0.26	0	3,3,3	0.28	0
5	EDO	A	1800	-	3,3,3	0.59	0	2,2,2	0.33	0
5	EDO	A	1812	-	3,3,3	0.51	0	2,2,2	0.37	0
5	EDO	A	1810	-	3,3,3	0.57	0	2,2,2	0.36	0
2	AU5	A	1795	-	37,37,37	1.07	3 (8%)	44,53,53	1.44	6 (13%)
3	P4C	A	1796	-	21,21,21	0.88	1 (4%)	20,20,20	1.13	1 (5%)
5	EDO	A	1805	-	3,3,3	0.63	0	2,2,2	0.23	0
5	EDO	A	1811	-	3,3,3	0.57	0	2,2,2	0.21	0
4	DMS	A	1798	-	3,3,3	0.60	0	3,3,3	0.56	0
5	EDO	A	1806	-	3,3,3	0.65	0	2,2,2	0.12	0
5	EDO	A	1804	-	3,3,3	0.60	0	2,2,2	0.24	0
5	EDO	A	1803	-	3,3,3	0.54	0	2,2,2	0.26	0
3	P4C	A	1797	-	15,15,21	0.11	0	14,14,20	0.13	0
5	EDO	A	1809	-	3,3,3	0.53	0	2,2,2	0.33	0
5	EDO	A	1802	-	3,3,3	0.39	0	2,2,2	0.50	0
5	EDO	A	1807	-	3,3,3	0.48	0	2,2,2	0.38	0
5	EDO	A	1808	-	3,3,3	0.52	0	2,2,2	0.34	0
4	DMS	A	1813	-	3,3,3	0.30	0	3,3,3	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1804	-	-	0/1/1/1	-
5	EDO	A	1802	-	-	0/1/1/1	-
5	EDO	A	1809	-	-	0/1/1/1	-
5	EDO	A	1807	-	-	0/1/1/1	-
5	EDO	A	1801	-	-	0/1/1/1	-
5	EDO	A	1808	-	-	0/1/1/1	-
3	P4C	A	1796	-	-	5/18/19/19	-
5	EDO	A	1800	-	-	1/1/1/1	-
5	EDO	A	1803	-	-	0/1/1/1	-
5	EDO	A	1805	-	-	1/1/1/1	-
3	P4C	A	1797	-	-	2/13/13/19	-
5	EDO	A	1811	-	-	0/1/1/1	-
5	EDO	A	1812	-	-	0/1/1/1	-
5	EDO	A	1806	-	-	0/1/1/1	-
5	EDO	A	1810	-	-	0/1/1/1	-
2	AU5	A	1795	-	-	4/14/31/31	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1796	P4C	O22-C21	3.94	1.42	1.20
2	A	1795	AU5	C4-N2	3.03	1.41	1.37
2	A	1795	AU5	C-N	2.98	1.38	1.34
2	A	1795	AU5	C4-N3	2.57	1.42	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1796	P4C	O22-C21-C20	-4.94	111.50	126.10
2	A	1795	AU5	O1-C16-C11	4.19	120.03	114.81
2	A	1795	AU5	O1-C16-C15	-3.45	118.13	124.08
2	A	1795	AU5	C17-O1-C16	-3.12	112.94	117.51
2	A	1795	AU5	C2-N2-C4	-2.92	108.59	110.10
2	A	1795	AU5	C2-C1-N1	-2.29	124.89	126.50
2	A	1795	AU5	C19-C20-N5	2.24	114.08	111.20

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1795	AU5	C11-C16-O1-C17

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1795	AU5	C15-C16-O1-C17
5	A	1805	EDO	O1-C1-C2-O2
2	A	1795	AU5	C15-C14-O2-C18
3	A	1797	P4C	C2-C3-O4-C5
2	A	1795	AU5	C13-C14-O2-C18
3	A	1797	P4C	C9-C8-O7-C6
3	A	1796	P4C	C8-C9-O10-C11
3	A	1796	P4C	C15-C14-O13-C12
3	A	1796	P4C	C18-C17-O16-C15
3	A	1796	P4C	C21-C20-O19-C18
5	A	1800	EDO	O1-C1-C2-O2
3	A	1796	P4C	O10-C11-C12-O13

There are no ring outliers.

7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1800	EDO	2	0
5	A	1810	EDO	1	0
2	A	1795	AU5	4	0
3	A	1796	P4C	5	0
5	A	1805	EDO	1	0
4	A	1798	DMS	1	0
5	A	1803	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	255/313 (81%)	0.74	29 (11%)	10 11	49, 70, 105, 137	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	515	PHE	4.5
1	A	738	ILE	4.4
1	A	726	TYR	4.4
1	A	795	THR	4.0
1	A	516	GLN	3.9
1	A	711	ILE	3.7
1	A	698	MET	3.3
1	A	669	ASN	3.2
1	A	612	TRP	3.1
1	A	748	ILE	3.0
1	A	622	TRP	2.9
1	A	547	LYS	2.7
1	A	615	LYS	2.6
1	A	614	LYS	2.4
1	A	536	SER	2.4
1	A	528	LEU	2.3
1	A	619	ILE	2.2
1	A	627	TYR	2.2
1	A	729	TYR	2.2
1	A	527	ILE	2.2
1	A	535	GLY	2.2
1	A	618	SER	2.2
1	A	530	GLN	2.1
1	A	540	PHE	2.1
1	A	684	VAL	2.1
1	A	522	GLY	2.1
1	A	537	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	762	LYS	2.0
1	A	685	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TPO	A	686	11/12	0.88	0.14	92,98,108,113	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DMS	A	1798	4/4	0.72	0.31	122,122,123,123	0
5	EDO	A	1804	4/4	0.73	0.18	85,86,88,88	0
5	EDO	A	1812	4/4	0.75	0.29	105,105,105,105	0
5	EDO	A	1805	4/4	0.79	0.15	79,81,81,82	0
4	DMS	A	1813	4/4	0.79	0.25	108,108,108,109	0
5	EDO	A	1810	4/4	0.81	0.19	77,78,80,81	0
5	EDO	A	1809	4/4	0.82	0.15	114,115,116,117	0
5	EDO	A	1806	4/4	0.84	0.24	87,87,88,88	0
5	EDO	A	1811	4/4	0.89	0.19	83,85,87,87	0
5	EDO	A	1808	4/4	0.89	0.17	98,99,99,99	0
5	EDO	A	1807	4/4	0.90	0.21	94,96,98,101	0
5	EDO	A	1802	4/4	0.90	0.15	60,63,66,71	0
3	P4C	A	1797	16/22	0.90	0.14	82,92,98,100	0
5	EDO	A	1803	4/4	0.91	0.16	83,84,84,86	0
3	P4C	A	1796	22/22	0.91	0.15	64,70,90,91	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	A	1801	4/4	0.92	0.15	93,93,93,93	0
4	DMS	A	1799	4/4	0.92	0.18	109,110,111,112	0
2	AU5	A	1795	33/33	0.93	0.11	48,56,85,88	0
5	EDO	A	1800	4/4	0.94	0.12	50,57,57,58	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around P4C A 1796:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.