



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:07 AM UTC

PDB ID : 7AP4 / pdb_00007ap4
Title : Thermus thermophilus Aspartyl-tRNA Synthetase in Complex with Compound AspS7HMDDA
Authors : De Graef, S.; Pang, L.; Strelkov, S.V.; Weeks, S.D.
Deposited on : 2020-10-15
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	581	8% 96% ..
1	B	581	24% 92% 8%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 18134 atoms, of which 8807 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aspartate-tRNA(Asp) ligase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	573	Total	C	H	N	O	S	0	0	0
			8944	2883	4447	797	806	11			
1	B	580	Total	C	H	N	O	S	0	2	0
			8783	2860	4316	797	799	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q5SKD2
B	0	GLY	-	expression tag	UNP Q5SKD2

- Molecule 2 is (3 {S})-3-azanyl-4-[[[(2 {R}),3 {S},4 {R},5 {R})-5-[7-azanyl-5-(hydroxymethyl) benzimidazol-1-yl]-3,4-bis(oxidanyl)oxolan-2-yl]methoxysulfonylamino]-4-oxidanylidene-b utanoic acid (CCD ID: RSN) (formula: C₁₇H₂₃N₅O₁₀S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	S	0	0
			55	17	22	5	10	1		
2	B	1	Total	C	H	N	O	S	0	0
			55	17	22	5	10	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	166	Total	O	0	0
			166	166		
3	B	131	Total	O	0	0
			131	131		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aspartate-tRNA(Asp) ligase



- Molecule 1: Aspartate-tRNA(Asp) ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.28Å 112.53Å 88.25Å 90.00° 104.86° 90.00°	Depositor
Resolution (Å)	79.53 – 2.15 79.53 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.8 (79.53-2.15) 98.9 (79.53-2.15)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.07Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.199 , 0.240 (Not available) , 0.265	Depositor DCC
R_{free} test set	2340 reflections (2.44%)	wwPDB-VP
Wilson B-factor (Å ²)	47.4	Xtriage
Anisotropy	0.344	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	18134	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RSN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/4603	0.33	0/6243
1	B	0.12	0/4585	0.33	0/6235
All	All	0.13	0/9188	0.33	0/12478

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4497	4447	4447	7	0
1	B	4467	4316	4304	27	0
2	A	33	22	0	1	0
2	B	33	22	0	0	0
3	A	166	0	0	1	0
3	B	131	0	0	1	0
All	All	9327	8807	8751	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:302:VAL:HG13	1:B:305:LEU:HD12	1.56	0.88
2:A:601:RSN:N6	3:A:701:HOH:O	2.29	0.65
1:B:302:VAL:CG1	1:B:305:LEU:HD12	2.26	0.64
1:B:329:LEU:HG	1:B:385:THR:HG21	1.82	0.61
1:A:501:GLU:N	1:A:501:GLU:OE1	2.38	0.57
1:B:421:LEU:C	1:B:421:LEU:HD12	2.30	0.56
1:B:338:GLU:N	1:B:349:LEU:HD22	2.24	0.53
1:B:25:VAL:HG21	1:B:63:VAL:CG1	2.38	0.53
1:B:251:LEU:O	1:B:255:GLU:HG2	2.10	0.52
1:B:363:VAL:HG12	1:B:363:VAL:O	2.10	0.52
1:B:312:ARG:HA	1:B:315:GLN:HG3	1.92	0.51
1:A:349:LEU:HD11	1:A:387:LEU:HB3	1.92	0.51
1:A:349:LEU:HD12	1:A:388:PHE:O	2.12	0.50
1:B:302:VAL:HG13	1:B:305:LEU:CD1	2.35	0.50
1:B:161:GLU:OE2	1:B:256:ARG:NH2	2.40	0.49
1:B:228:ARG:NH2	3:B:708:HOH:O	2.45	0.48
1:B:240:LEU:C	1:B:240:LEU:HD12	2.39	0.48
1:B:326:PRO:HA	1:B:384:ASP:OD1	2.16	0.46
1:B:496:LEU:HD13	1:B:496:LEU:O	2.15	0.46
1:A:240:LEU:HD12	1:A:240:LEU:C	2.41	0.45
1:B:290:LYS:HA	1:B:471:VAL:HG21	1.98	0.45
1:B:202:LEU:HD21	1:B:445:PHE:CZ	2.51	0.45
1:B:363:VAL:HG12	1:B:366:PHE:HB2	1.99	0.44
1:B:446:THR:HG23	1:B:481:SER:HB2	1.98	0.44
1:B:7:ALA:N	1:B:41:ASP:OD2	2.41	0.44
1:A:204:LYS:HD3	1:A:239:ASP:OD1	2.17	0.43
1:A:228:ARG:O	1:A:229:ALA:HB3	2.19	0.43
1:B:351:TRP:HA	1:B:386:LEU:O	2.19	0.43
1:B:173:LYS:O	1:B:174:SER:C	2.62	0.42
1:B:25:VAL:HG21	1:B:63:VAL:HG13	2.01	0.42
1:B:202:LEU:HD21	1:B:445:PHE:CE2	2.55	0.42
1:A:297:LEU:HB3	1:A:323:LEU:HD11	2.02	0.41
1:B:385:THR:O	1:B:385:THR:HG23	2.21	0.41
1:B:78:ARG:HD2	1:B:90:VAL:O	2.22	0.40
1:B:204:LYS:HD2	1:B:241:GLU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	567/581 (98%)	544 (96%)	22 (4%)	1 (0%)	43	44
1	B	578/581 (100%)	558 (96%)	18 (3%)	2 (0%)	36	34
All	All	1145/1162 (98%)	1102 (96%)	40 (4%)	3 (0%)	36	34

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	371	ARG
1	B	227	LEU
1	A	362	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	454/483 (94%)	449 (99%)	5 (1%)	65	72
1	B	436/483 (90%)	428 (98%)	8 (2%)	51	58
All	All	890/966 (92%)	877 (98%)	13 (2%)	57	64

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	215	TYR
1	A	226	ASP
1	A	355	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	379	GLU
1	A	441	MET
1	B	215	TYR
1	B	225	GLU
1	B	253	LEU
1	B	315	GLN
1	B	413	LEU
1	B	462	ARG
1	B	488	ARG
1	B	496	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	47	GLN
1	A	308	GLN
1	A	315	GLN
1	A	504	GLN
1	B	217	GLN
1	B	315	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RSN	B	601	-	35,35,35	0.92	1 (2%)	45,52,52	0.73	1 (2%)
2	RSN	A	601	-	35,35,35	0.97	1 (2%)	45,52,52	0.72	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RSN	B	601	-	-	1/24/41/41	0/3/3/3
2	RSN	A	601	-	-	6/24/41/41	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	RSN	C-N3S	5.07	1.46	1.37
2	B	601	RSN	C-N3S	4.74	1.46	1.37

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	RSN	O-C-N3S	-3.41	116.80	122.98
2	A	601	RSN	O-C-N3S	-3.39	116.83	122.98

There are no chirality outliers.

All (7) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	RSN	N-CA-CB-CG
2	A	601	RSN	O-C-CA-CB
2	A	601	RSN	O-C-CA-N
2	A	601	RSN	N3S-C-CA-CB
2	A	601	RSN	C-CA-CB-CG
2	A	601	RSN	C3'-C4'-C5'-O5'
2	B	601	RSN	C3'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	RSN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	573/581 (98%)	0.62	46 (8%)	18 20	38, 60, 92, 119	0
1	B	580/581 (99%)	1.39	141 (24%)	2 2	30, 64, 173, 277	1 (0%)
All	All	1153/1162 (99%)	1.01	187 (16%)	4 5	30, 62, 159, 277	1 (0%)

All (187) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	176	PRO	11.1
1	B	325	LEU	10.3
1	B	323	LEU	9.4
1	A	432	ASP	8.8
1	B	227	LEU	8.5
1	B	387	LEU	8.5
1	B	360	SER	6.8
1	B	314	PHE	6.7
1	B	496	LEU	6.6
1	B	359	PHE	6.5
1	B	320	VAL	6.4
1	A	437	ALA	6.2
1	B	337	LEU	6.2
1	B	389	VAL	6.1
1	B	373	ALA	5.8
1	B	375	LEU	5.7
1	B	326	PRO	5.7
1	B	366	PHE	5.6
1	B	386	LEU	5.5
1	A	176	PRO	5.4
1	B	399	ALA	5.4
1	B	334	VAL	5.3
1	B	385	THR	5.2
1	A	431	TRP	5.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	336	GLU	5.2
1	B	351	TRP	5.1
1	B	371	ARG	5.1
1	B	324	ALA	4.8
1	B	306	PHE	4.8
1	B	369	PRO	4.8
1	B	388	PHE	4.7
1	B	312	ARG	4.7
1	B	363	VAL	4.7
1	B	440	TYR	4.7
1	B	370	VAL	4.6
1	B	329	LEU	4.6
1	B	178	GLY	4.5
1	B	367	LEU	4.5
1	B	365	LYS	4.5
1	B	340	VAL	4.5
1	B	348	GLY	4.4
1	B	349	LEU	4.4
1	B	374	LEU	4.4
1	B	310	GLY	4.4
1	B	356	GLU	4.3
1	B	382	PRO	4.3
1	B	352	ALA	4.3
1	B	174	SER	4.3
1	B	335	ALA	4.2
1	B	378	THR	4.2
1	B	322	ALA	4.1
1	B	350	ALA	4.1
1	B	439	THR	4.0
1	B	364	ALA	4.0
1	B	377	ALA	4.0
1	B	338	GLU	4.0
1	B	372	GLU	4.0
1	B	311	PHE	4.0
1	B	381	ARG	4.0
1	B	354	VAL	4.0
1	B	390	ALA	3.9
1	B	305	LEU	3.9
1	B	342	LYS	3.9
1	B	376	GLN	3.9
1	B	400	LEU	3.8
1	B	383	GLY	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	433	GLU	3.8
1	B	410	LEU	3.8
1	B	353	ARG	3.8
1	B	411	LEU	3.7
1	B	341	ALA	3.7
1	B	307	ARG	3.7
1	B	332	LYS	3.6
1	B	346	ALA	3.6
1	A	362	GLY	3.6
1	A	502	GLU	3.5
1	B	339	GLU	3.5
1	B	407	ALA	3.5
1	B	316	GLU	3.5
1	B	380	ALA	3.5
1	B	304	PRO	3.4
1	B	300	LYS	3.4
1	B	328	ALA	3.4
1	B	229	ALA	3.4
1	B	343	ARG	3.3
1	B	368	GLU	3.3
1	B	309	SER	3.3
1	B	361	GLY	3.3
1	B	393	ARG	3.2
1	B	355	GLU	3.2
1	B	497	LEU	3.1
1	B	391	GLY	3.1
1	B	435	GLU	3.1
1	A	442	HIS	3.0
1	B	362	GLY	3.0
1	B	392	PRO	3.0
1	B	302	VAL	3.0
1	B	396	ALA	3.0
1	B	318	GLU	3.0
1	B	345	LYS	3.0
1	B	358	GLY	2.9
1	B	498	GLY	2.9
1	B	327	LYS	2.9
1	B	276	ARG	2.9
1	B	495	ARG	2.9
1	B	431	TRP	2.9
1	B	331	ARG	2.9
1	A	508	PHE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	228	ARG	2.8
1	A	31	LEU	2.8
1	A	429	LEU	2.8
1	B	434	GLU	2.8
1	B	347	GLN	2.8
1	B	344	HIS	2.8
1	B	177	GLU	2.8
1	B	321	LYS	2.8
1	B	357	GLY	2.7
1	A	507	LYS	2.7
1	A	440	TYR	2.7
1	B	508	PHE	2.7
1	B	395	VAL	2.7
1	B	458	LYS	2.7
1	A	359	PHE	2.7
1	B	319	SER	2.7
1	B	413	LEU	2.6
1	A	33	GLY	2.6
1	A	443	HIS	2.6
1	A	366	PHE	2.5
1	A	445	PHE	2.5
1	B	179	ALA	2.5
1	B	308	GLN	2.5
1	B	405	LEU	2.5
1	B	330	SER	2.5
1	A	179	ALA	2.5
1	B	408	ALA	2.5
1	A	311	PHE	2.5
1	B	170	PHE	2.5
1	A	0	GLY	2.5
1	A	361	GLY	2.5
1	A	578	VAL	2.4
1	A	372	GLU	2.4
1	A	506	GLU	2.4
1	B	442	HIS	2.4
1	B	500	GLY	2.4
1	B	444	PRO	2.4
1	B	397	ALA	2.4
1	A	358	GLY	2.4
1	A	226	ASP	2.4
1	A	363	VAL	2.4
1	A	309	SER	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	381	ARG	2.4
1	B	507	LYS	2.4
1	B	398	THR	2.3
1	B	303	GLY	2.3
1	B	313	VAL	2.3
1	A	410	LEU	2.3
1	A	438	TRP	2.3
1	B	571	ARG	2.3
1	B	173	LYS	2.3
1	A	310	GLY	2.3
1	A	441	MET	2.3
1	B	579	ARG	2.2
1	B	196	LEU	2.2
1	A	313	VAL	2.2
1	A	505	ARG	2.2
1	A	370	VAL	2.2
1	B	545	VAL	2.2
1	B	279	TYR	2.2
1	B	499	ILE	2.1
1	A	565	VAL	2.1
1	A	328	ALA	2.1
1	A	576	MET	2.1
1	B	429	LEU	2.1
1	A	181	ASP	2.1
1	B	403	VAL	2.1
1	A	421	LEU	2.1
1	B	299	LEU	2.1
1	B	428	LEU	2.1
1	A	105	THR	2.1
1	B	477	VAL	2.1
1	B	502	GLU	2.0
1	A	411	LEU	2.0
1	A	331	ARG	2.0
1	A	360	SER	2.0
1	B	438	TRP	2.0
1	B	394	LYS	2.0
1	A	357	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	RSN	A	601	33/33	0.86	0.13	53,82,98,114	0
2	RSN	B	601	33/33	0.94	0.09	50,62,75,93	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.