



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 10:00 AM UTC

PDB ID : 5AQL / pdb_00005aql
Title : Fragment-based screening of HSP70 sheds light on the functional role of ATP-binding site residues
Authors : Jones, A.M.; Westwood, I.M.; Osborne, J.D.; Matthews, T.P.; Cheeseman, M.D.; Rowlands, M.G.; Jeganathan, F.; Burke, R.; Lee, D.; Kadi, N.; Liu, M.; Richards, M.; McAndrew, C.; Yahya, N.; Dobson, S.E.; Jones, K.; Workman, P.; Collins, I.; van Montfort, R.L.M.
Deposited on : 2015-09-22
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	94% 5%
1	C	386	94% 5%
2	B	118	93%
2	D	118	20% 87% 8%

Validation Pipeline (wwPDB-VP) : 2.49

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 8593 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEAT SHOCK COGNATE 71 KDA PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	381	Total	C	N	O	S	0	1	0
			2927	1844	506	568	9			
1	C	381	Total	C	N	O	S	0	3	0
			2962	1864	517	571	10			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	GLY	-	expression tag	UNP P11142
A	-3	PRO	-	expression tag	UNP P11142
A	-2	LEU	-	expression tag	UNP P11142
A	-1	GLY	-	expression tag	UNP P11142
A	0	SER	-	expression tag	UNP P11142
A	275	TRP	SER	engineered mutation	UNP P11142
C	-4	GLY	-	expression tag	UNP P11142
C	-3	PRO	-	expression tag	UNP P11142
C	-2	LEU	-	expression tag	UNP P11142
C	-1	GLY	-	expression tag	UNP P11142
C	0	SER	-	expression tag	UNP P11142
C	275	TRP	SER	engineered mutation	UNP P11142


- Molecule 2 is a protein called BAG FAMILY MOLECULAR CHAPERONE REGULATOR 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	113	Total	C	N	O	S	0	0	0
			876	555	148	169	4			
2	D	113	Total	C	N	O	S	0	1	0
			858	544	144	166	4			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	146	GLY	-	expression tag	UNP Q99933
B	147	PRO	-	expression tag	UNP Q99933
B	148	LEU	-	expression tag	UNP Q99933
B	149	GLY	-	expression tag	UNP Q99933
B	150	SER	-	expression tag	UNP Q99933
D	146	GLY	-	expression tag	UNP Q99933
D	147	PRO	-	expression tag	UNP Q99933
D	148	LEU	-	expression tag	UNP Q99933
D	149	GLY	-	expression tag	UNP Q99933
D	150	SER	-	expression tag	UNP Q99933

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	A	1	Total	C	N	O	0	0
			8	4	1	3		
3	C	1	Total	C	N	O	0	0
			8	4	1	3		
3	C	1	Total	C	N	O	0	0
			8	4	1	3		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	367	Total	O	0	0
			367	367		
5	B	107	Total	O	0	0
			107	107		
5	C	361	Total	O	0	0
			361	361		
5	D	73	Total	O	0	0
			73	73		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HEAT SHOCK COGNATE 71 KDA PROTEIN

Chain A: 





- Molecule 1: HEAT SHOCK COGNATE 71 KDA PROTEIN

Chain C: 




- Molecule 2: BAG FAMILY MOLECULAR CHAPERONE REGULATOR 1

Chain B: 



- Molecule 2: BAG FAMILY MOLECULAR CHAPERONE REGULATOR 1

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	232.03Å 40.93Å 116.55Å 90.00° 90.58° 90.00°	Depositor
Resolution (Å)	58.27 – 1.69 58.27 – 1.69	Depositor EDS
% Data completeness (in resolution range)	97.4 (58.27-1.69) 97.4 (58.27-1.69)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.69Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.169 , 0.197 0.178 , 0.206	Depositor DCC
R_{free} test set	6048 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.4	Xtriage
Anisotropy	0.123	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8593	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	0/2975	0.98	5/4026 (0.1%)
1	C	0.73	0/3010	0.97	5/4069 (0.1%)
2	B	0.80	0/883	1.08	0/1187
2	D	0.80	0/865	1.17	3/1167 (0.3%)
All	All	0.75	0/7733	1.01	13/10449 (0.1%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	151	ASN	N-CA-C	7.23	120.06	111.02
1	C	29	ILE	N-CA-C	6.59	118.07	108.58
1	A	29	ILE	N-CA-C	6.38	117.77	108.58
1	A	47	THR	N-CA-C	5.85	121.45	113.97
1	A	254	SER	N-CA-C	5.52	118.06	111.71
1	C	254	SER	N-CA-C	5.41	117.93	111.71
1	A	143	VAL	N-CA-C	-5.41	100.60	108.17
1	A	332	HIS	N-CA-C	5.37	117.21	111.36
1	C	332	HIS	N-CA-C	5.34	117.18	111.36
2	D	184	THR	CA-C-N	5.22	125.77	119.98
2	D	184	THR	C-N-CA	5.22	125.77	119.98
1	C	143	VAL	N-CA-C	-5.21	100.88	108.17
1	C	172	ILE	N-CA-C	-5.15	100.33	107.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2927	0	2879	8	0
1	C	2962	0	2943	10	0
2	B	876	0	875	4	0
2	D	858	0	839	6	0
3	A	16	0	24	3	0
3	C	16	0	24	2	0
4	B	18	0	24	1	0
4	C	6	0	8	0	0
4	D	6	0	8	0	0
5	A	367	0	0	2	0
5	B	107	0	0	0	0
5	C	361	0	0	0	0
5	D	73	0	0	0	0
All	All	8593	0	7624	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:315:ASP:HB2	1:C:316:PRO:HD3	1.69	0.74
1:A:315:ASP:HB2	1:A:316:PRO:HD3	1.72	0.71
2:B:250:GLU:HA	4:B:1263:GOL:H32	1.74	0.69
1:A:42:VAL:HG13	1:A:51:ILE:HD13	1.92	0.52
2:D:183:LEU:HD12	2:D:200:LEU:HD22	1.92	0.51
1:C:42:VAL:HG13	1:C:51:ILE:HD13	1.94	0.50
1:C:279:GLN:HG2	1:C:280:ALA:N	2.26	0.49
2:D:227:PRO:HB2	2:D:229:ASN:OD1	2.13	0.49
2:B:153:PRO:HB2	2:D:190:PHE:HZ	1.78	0.48
1:C:171:ARG:NE	1:C:173:ILE:HG22	2.28	0.48
3:A:1383:TRS:H31	2:B:245:GLN:NE2	2.29	0.47
1:C:200:LEU:HG	1:C:340:SER:HB2	1.96	0.47
1:A:363:ILE:O	1:A:364:ASN:C	2.59	0.45
1:A:200:LEU:HG	1:A:340:SER:HB2	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:240:HIS:CE1	1:C:301:ARG:HD3	2.51	0.45
3:A:1383:TRS:H31	2:B:245:GLN:HE21	1.83	0.44
1:A:23:HIS:HE1	5:A:2016:HOH:O	2.01	0.43
1:A:68:PHE:O	1:A:69:ASP:HB2	2.18	0.43
3:A:1383:TRS:H32	5:A:2278:HOH:O	2.17	0.43
1:C:363:ILE:O	1:C:364:ASN:C	2.62	0.42
3:C:1383:TRS:H11	2:D:245:GLN:NE2	2.35	0.42
1:C:315:ASP:HB2	1:C:316:PRO:CD	2.45	0.42
3:C:1383:TRS:H11	2:D:245:GLN:HE21	1.84	0.41
1:A:75:GLY:HA3	1:A:154:GLN:HA	2.02	0.41
2:D:203:LEU:O	2:D:207[A]:VAL:HG23	2.21	0.41
1:C:279:GLN:HG2	1:C:280:ALA:H	1.85	0.40
1:C:315:ASP:CB	1:C:316:PRO:HD3	2.46	0.40
1:A:237:MET:HE2	1:A:237:MET:HB3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/386 (98%)	377 (99%)	3 (1%)	0	100	100
1	C	382/386 (99%)	379 (99%)	3 (1%)	0	100	100
2	B	111/118 (94%)	111 (100%)	0	0	100	100
2	D	112/118 (95%)	109 (97%)	3 (3%)	0	100	100
All	All	985/1008 (98%)	976 (99%)	9 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	304/319 (95%)	304 (100%)	0	100	100
1	C	312/319 (98%)	312 (100%)	0	100	100
2	B	91/107 (85%)	91 (100%)	0	100	100
2	D	87/107 (81%)	87 (100%)	0	100	100
All	All	794/852 (93%)	794 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	22	GLN
1	A	33	GLN
1	A	168	ASN
1	A	194	ASN
1	A	227	HIS
2	B	196	GLN
2	B	245	GLN
1	C	22	GLN
1	C	104	GLN
1	C	168	ASN
2	D	245	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	B	1262	-	5,5,5	0.06	0	5,5,5	0.35	0
4	GOL	C	1384	-	5,5,5	0.05	0	5,5,5	0.14	0
3	TRS	A	1382	-	7,7,7	0.30	0	9,9,9	0.45	0
3	TRS	C	1383	-	7,7,7	0.18	0	9,9,9	0.16	0
4	GOL	B	1261	-	5,5,5	0.15	0	5,5,5	0.44	0
3	TRS	A	1383	-	7,7,7	0.18	0	9,9,9	0.27	0
4	GOL	B	1263	-	5,5,5	0.06	0	5,5,5	0.14	0
4	GOL	D	1261	-	5,5,5	0.15	0	5,5,5	0.42	0
3	TRS	C	1382	-	7,7,7	0.26	0	9,9,9	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	1262	-	-	0/4/4/4	-
4	GOL	C	1384	-	-	0/4/4/4	-
3	TRS	A	1382	-	-	3/9/9/9	-
3	TRS	C	1383	-	-	3/9/9/9	-
4	GOL	B	1261	-	-	0/4/4/4	-
3	TRS	A	1383	-	-	3/9/9/9	-
4	GOL	B	1263	-	-	0/4/4/4	-
4	GOL	D	1261	-	-	0/4/4/4	-
3	TRS	C	1382	-	-	3/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1382	TRS	N-C-C2-O2
3	A	1383	TRS	N-C-C3-O3
3	C	1382	TRS	N-C-C2-O2
3	C	1383	TRS	N-C-C1-O1
3	A	1382	TRS	C1-C-C2-O2
3	A	1383	TRS	C2-C-C3-O3
3	C	1382	TRS	C1-C-C2-O2
3	C	1383	TRS	C2-C-C1-O1
3	C	1383	TRS	C3-C-C1-O1
3	A	1383	TRS	C1-C-C3-O3
3	A	1382	TRS	C3-C-C2-O2
3	C	1382	TRS	C3-C-C2-O2

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1383	TRS	2	0
3	A	1383	TRS	3	0
4	B	1263	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	381/386 (98%)	0.01	4 (1%) 79 82	11, 28, 46, 62	1 (0%)
1	C	381/386 (98%)	0.04	8 (2%) 63 68	11, 28, 45, 60	3 (0%)
2	B	113/118 (95%)	0.12	5 (4%) 39 42	18, 29, 47, 56	0
2	D	113/118 (95%)	0.82	24 (21%) 2 2	11, 34, 76, 85	1 (0%)
All	All	988/1008 (98%)	0.13	41 (4%) 41 45	11, 28, 49, 85	5 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	190	PHE	6.4
2	D	195	LEU	5.1
2	D	260	GLN	4.9
2	D	148	LEU	4.6
2	B	148	LEU	4.4
2	D	194	ASP	4.2
2	D	186	ILE	3.9
2	D	199	ALA	3.9
2	D	192	PRO	3.8
2	D	200	LEU	3.7
2	D	193	LYS	3.4
2	D	191	LEU	3.2
2	D	259	CYS	3.2
1	C	89	HIS	3.1
2	D	188	GLN	3.1
2	D	201	CYS	3.0
2	D	187	GLN	2.8
1	A	189	VAL	2.8
2	D	183	LEU	2.7
2	D	185	GLY	2.7
2	D	197	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	149	GLY	2.6
1	C	93	MET	2.5
2	B	195	LEU	2.5
1	C	88	LYS	2.4
2	D	149	GLY	2.4
1	A	89	HIS	2.4
2	B	259	CYS	2.2
1	C	381	SER	2.2
2	D	150	SER	2.2
2	D	189	GLY	2.2
1	A	381	SER	2.2
1	C	201	GLY	2.2
2	B	193	LYS	2.2
1	A	2	SER	2.1
1	C	2	SER	2.1
2	D	196	GLN	2.1
2	D	184	THR	2.1
1	C	191	ALA	2.1
2	D	203	LEU	2.1
1	C	189	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	C	1384	6/6	0.74	0.27	77,79,79,80	0
4	GOL	B	1263	6/6	0.79	0.21	76,77,77,78	0
3	TRS	C	1382	8/8	0.80	0.14	38,43,45,45	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	1262	6/6	0.83	0.16	36,50,54,56	0
4	GOL	D	1261	6/6	0.85	0.12	36,44,53,59	0
3	TRS	A	1382	8/8	0.89	0.11	32,39,41,42	0
4	GOL	B	1261	6/6	0.89	0.13	46,48,48,49	0
3	TRS	C	1383	8/8	0.90	0.13	25,38,45,52	0
3	TRS	A	1383	8/8	0.91	0.13	25,37,41,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.