



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 2, 2026 – 12:25 AM UTC

PDB ID : 3ARQ / pdb_00003arq
Title : Crystal Structure Analysis of Chitinase A from *Vibrio harveyi* with novel inhibitors - complex structure with IDARUBICIN
Authors : Pantoom, S.; Vetter, I.R.; Prinz, H.; Suginta, W.
Deposited on : 2010-12-09
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	584	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	3502	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5478 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Chitinase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	575	4592	2933	729	901	29	0	33	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	598	ARG	-	expression tag	UNP Q9AMP1
A	599	SER	-	expression tag	UNP Q9AMP1
A	600	HIS	-	expression tag	UNP Q9AMP1
A	601	HIS	-	expression tag	UNP Q9AMP1
A	602	HIS	-	expression tag	UNP Q9AMP1
A	603	HIS	-	expression tag	UNP Q9AMP1
A	604	HIS	-	expression tag	UNP Q9AMP1
A	605	HIS	-	expression tag	UNP Q9AMP1

- Molecule 2 is IDARUBICIN (CCD ID: DM5) (formula: C₂₆H₂₇NO₉).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			36	26	1	9		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	1
			12	6	6		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	826	Total	O	0	0
			826	826		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Chitinase A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.01Å 50.89Å 93.40Å 90.00° 99.51° 90.00°	Depositor
Resolution (Å)	19.93 – 1.50 19.93 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.93-1.50) 99.5 (19.93-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.70 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.6.0093	Depositor
R, R_{free}	0.145 , 0.184 (Not available) , 0.181	Depositor DCC
R_{free} test set	4808 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	10.0	Xtriage
Anisotropy	0.171	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5478	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, DM5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.58	28/4795 (0.6%)	1.35	19/6523 (0.3%)

All (28) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	526	TYR	N-CA	7.39	1.55	1.46
1	A	195	ILE	N-CA	6.94	1.54	1.46
1	A	202	VAL	CA-CB	6.84	1.64	1.54
1	A	247	THR	CA-C	6.75	1.60	1.53
1	A	522	PHE	C-O	6.45	1.31	1.23
1	A	328	ASP	N-CA	6.41	1.53	1.46
1	A	234	TYR	N-CA	6.27	1.53	1.46
1	A	560	LYS	C-O	6.06	1.31	1.24
1	A	345	VAL	N-CA	5.88	1.53	1.46
1	A	539	SER	N-CA	5.88	1.53	1.46
1	A	593	THR	CB-OG1	5.71	1.52	1.43
1	A	302	LYS	N-CA	5.70	1.53	1.46
1	A	501	VAL	CA-CB	-5.66	1.47	1.54
1	A	509	SER	CA-C	-5.57	1.45	1.52
1	A	535	VAL	N-CA	5.53	1.52	1.46
1	A	390	THR	N-CA	5.50	1.53	1.46
1	A	420	GLY	N-CA	5.45	1.53	1.45
1	A	295	ALA	CA-CB	5.45	1.61	1.53
1	A	482	GLY	N-CA	5.31	1.50	1.45
1	A	509	SER	N-CA	5.15	1.52	1.46
1	A	157	PRO	N-CA	-5.15	1.41	1.47
1	A	443	LEU	N-CA	5.09	1.52	1.46
1	A	191	GLY	CA-C	5.05	1.55	1.52
1	A	546	PHE	CA-CB	5.05	1.60	1.53
1	A	132	ALA	N-CA	5.05	1.52	1.46
1	A	24	THR	CA-C	5.01	1.59	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	547	ASP	C-O	5.01	1.30	1.24
1	A	526	TYR	CA-C	-5.01	1.46	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	290	ARG	NE-CZ-NH2	-7.08	112.83	119.20
1	A	316	PHE	CA-C-N	-6.65	112.81	119.93
1	A	316	PHE	C-N-CA	-6.65	112.81	119.93
1	A	508	LYS	CG-CD-CE	-6.47	96.43	111.30
1	A	557	ASN	CA-CB-CG	-6.19	106.41	112.60
1	A	560	LYS	N-CA-C	-6.01	104.73	111.28
1	A	30	MET	CA-C-O	5.76	126.52	120.42
1	A	115	ALA	CA-C-O	-5.62	114.63	120.70
1	A	538	ARG	NE-CZ-NH1	-5.43	116.07	121.50
1	A	26	PRO	CB-CA-C	-5.40	104.70	111.56
1	A	504	TYR	O-C-N	5.37	127.81	122.12
1	A	464	GLY	O-C-N	5.35	129.65	123.44
1	A	375	ASP	CA-C-O	5.33	127.55	121.58
1	A	193	ILE	CA-C-O	5.33	123.06	119.18
1	A	114	GLU	N-CA-CB	-5.14	102.41	110.69
1	A	560	LYS	CA-C-O	-5.13	115.11	120.55
1	A	504	TYR	CA-C-O	-5.09	115.14	120.63
1	A	344	ARG	O-C-N	5.07	128.15	122.22
1	A	305	LYS	CD-CE-NZ	-5.04	95.77	111.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4592	0	4398	66	0
2	A	36	0	25	5	0
3	A	24	0	32	8	0
4	A	826	0	0	17	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5478	0	4455	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:557:ASN:HB3	4:A:698:HOH:O	1.40	1.22
1:A:511[B]:MET:HE1	1:A:544:ILE:HD11	1.42	0.99
1:A:511[B]:MET:SD	1:A:544:ILE:HD12	2.06	0.95
1:A:511[B]:MET:SD	1:A:544:ILE:CD1	2.55	0.93
1:A:31:TYR:O	4:A:693:HOH:O	1.87	0.91
1:A:511[B]:MET:HE1	1:A:544:ILE:CD1	2.04	0.87
1:A:435[A]:TYR:CE2	2:A:606:DM5:H141	2.11	0.85
1:A:87[A]:VAL:HG21	1:A:103:TYR:CE1	2.13	0.82
1:A:79:ASN:OD1	1:A:89[B]:THR:HG22	1.81	0.81
1:A:45[B]:MET:HE2	1:A:250:LYS:NZ	1.97	0.79
1:A:511[B]:MET:CE	1:A:544:ILE:CD1	2.61	0.79
1:A:581:ASN:HB2	3:A:3502:GOL:H11	1.64	0.79
1:A:45[B]:MET:HE2	1:A:250:LYS:HZ1	1.49	0.76
1:A:34:ASN:HA	4:A:1361:HOH:O	1.86	0.75
1:A:570[B]:TRP:NE1	4:A:775:HOH:O	2.19	0.75
1:A:511[B]:MET:CE	1:A:544:ILE:HD11	2.18	0.73
1:A:87[A]:VAL:CG2	1:A:103:TYR:CE1	2.72	0.72
1:A:435[A]:TYR:HE2	2:A:606:DM5:H141	1.51	0.71
1:A:45[B]:MET:HE3	4:A:832:HOH:O	1.89	0.70
1:A:34:ASN:CA	4:A:1361:HOH:O	2.43	0.63
1:A:35[A]:ASN:OD1	1:A:37:GLN:HG2	1.98	0.63
1:A:30:MET:HE1	1:A:127:VAL:HG21	1.82	0.61
1:A:87[A]:VAL:CG2	1:A:103:TYR:HE1	2.13	0.61
1:A:581:ASN:CB	3:A:3502:GOL:H11	2.31	0.60
1:A:511[B]:MET:SD	1:A:544:ILE:HD13	2.42	0.59
1:A:435[A]:TYR:CZ	4:A:805:HOH:O	2.45	0.59
1:A:586:GLY:HA3	4:A:1076:HOH:O	2.02	0.58
1:A:389[B]:MET:HE3	1:A:391:TYR:HE1	1.69	0.56
1:A:435[A]:TYR:CD2	2:A:606:DM5:H141	2.41	0.55
1:A:151:LYS:HZ3	3:A:3502:GOL:H32	1.71	0.55
1:A:45[B]:MET:HE1	1:A:235:GLN:HA	1.90	0.54
1:A:248:PRO:HG3	4:A:740:HOH:O	2.08	0.54
1:A:581:ASN:HD22	3:A:3502:GOL:H31	1.73	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89[B]:THR:HG23	4:A:736:HOH:O	2.10	0.52
1:A:557:ASN:CB	4:A:698:HOH:O	2.23	0.52
1:A:151:LYS:NZ	3:A:3502:GOL:H32	2.24	0.51
1:A:592[B]:VAL:HG12	1:A:593:THR:N	2.26	0.51
1:A:581:ASN:HD22	3:A:3502:GOL:C3	2.25	0.49
1:A:46:GLU:CD	1:A:245:TYR:OH	2.55	0.49
1:A:570[B]:TRP:CD1	4:A:775:HOH:O	2.63	0.49
2:A:606:DM5:O6	2:A:606:DM5:O8	2.25	0.48
1:A:596:ASN:H	1:A:596:ASN:HD22	1.62	0.48
1:A:267:LYS:NZ	4:A:692:HOH:O	2.45	0.47
1:A:188:ILE:HD12	1:A:266:LEU:HD21	1.95	0.47
1:A:45[B]:MET:CE	1:A:250:LYS:NZ	2.74	0.47
1:A:154:ASN:N	4:A:814:HOH:O	2.10	0.46
1:A:159[B]:ILE:CD1	1:A:592[B]:VAL:HG23	2.46	0.46
1:A:169:GLY:HA3	4:A:885:HOH:O	2.16	0.44
1:A:549:HIS:CE1	1:A:553:LEU:HD11	2.53	0.44
1:A:596:ASN:HD22	1:A:596:ASN:N	2.16	0.44
1:A:159[B]:ILE:HD13	1:A:592[B]:VAL:HG23	2.01	0.43
1:A:305:LYS:HE3	4:A:630:HOH:O	2.18	0.42
1:A:392:ASP:O	2:A:606:DM5:H143	2.19	0.42
1:A:194:PRO:HG3	1:A:228[B]:HIS:HB2	2.02	0.42
1:A:407[B]:LEU:HD11	1:A:558:TYR:CD1	2.55	0.42
1:A:285:VAL:HG13	4:A:750:HOH:O	2.18	0.42
1:A:407[B]:LEU:HD11	1:A:558:TYR:CG	2.55	0.42
1:A:151:LYS:NZ	3:A:3502:GOL:C3	2.83	0.41
1:A:494:GLN:HA	1:A:510:PHE:CE2	2.55	0.41
1:A:151:LYS:HZ3	3:A:3502:GOL:C3	2.33	0.41
1:A:407[A]:LEU:HA	1:A:437:ALA:HB3	2.03	0.41
1:A:30:MET:HE1	1:A:127:VAL:CG2	2.50	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:688:HOH:O	4:A:1382:HOH:O[2_556]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	606/584 (104%)	588 (97%)	16 (3%)	2 (0%)	36 17

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	35[A]	ASN
1	A	35[B]	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	493/469 (105%)	487 (99%)	6 (1%)	63 38

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35[A]	ASN
1	A	35[B]	ASN
1	A	62	LYS
1	A	71	SER
1	A	141	PRO
1	A	596	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	34	ASN
1	A	37	GLN
1	A	52	ASN
1	A	96	GLN
1	A	105	GLN
1	A	154	ASN
1	A	199	ASN
1	A	447	GLN
1	A	596	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	3501	-	5,5,5	0.65	0	5,5,5	0.67	0
3	GOL	A	3502	-	5,5,5	0.84	0	5,5,5	1.15	0
3	GOL	A	3500[A]	-	5,5,5	0.51	0	5,5,5	1.27	1 (20%)
3	GOL	A	3500[B]	-	5,5,5	0.38	0	5,5,5	0.66	0
2	DM5	A	606	-	38,40,40	2.30	14 (36%)	50,63,63	1.51	10 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	3501	-	-	0/4/4/4	-
3	GOL	A	3502	-	-	2/4/4/4	-
3	GOL	A	3500[A]	-	-	0/4/4/4	-
3	GOL	A	3500[B]	-	-	0/4/4/4	-
2	DM5	A	606	-	-	0/10/56/56	0/5/5/5

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	606	DM5	O8-C8	-5.21	1.25	1.36
2	A	606	DM5	O10-C10	4.55	1.52	1.44
2	A	606	DM5	C5-C6	-4.32	1.39	1.48
2	A	606	DM5	O12-C12	4.00	1.50	1.43
2	A	606	DM5	C15-C16	3.72	1.57	1.51
2	A	606	DM5	C20-C19	-3.58	1.41	1.48
2	A	606	DM5	C2'-C1'	3.34	1.59	1.50
2	A	606	DM5	C14-C13	3.08	1.57	1.50
2	A	606	DM5	C18-C19	-2.95	1.40	1.47
2	A	606	DM5	O10-C1'	2.53	1.48	1.41
2	A	606	DM5	C9-C10	2.25	1.55	1.51
2	A	606	DM5	C2-C1	2.15	1.42	1.38
2	A	606	DM5	O5'-C1'	2.15	1.47	1.42
2	A	606	DM5	C7-C6	-2.12	1.42	1.47

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	606	DM5	O8-C8-C9	3.28	127.46	119.02
2	A	606	DM5	C11-C10-C9	3.10	117.97	110.71
2	A	606	DM5	O10-C10-C9	-3.08	100.62	107.54
2	A	606	DM5	O12-C12-C11	-2.70	103.25	109.40
2	A	606	DM5	O5'-C1'-C2'	2.68	114.99	110.84
2	A	606	DM5	O19-C19-C20	-2.56	117.30	120.99
2	A	606	DM5	O8-C8-C7	-2.55	115.35	120.95
2	A	606	DM5	C7-C8-C9	-2.25	117.17	121.28
2	A	606	DM5	C20-C19-C18	2.23	121.57	117.97
3	A	3500[A]	GOL	C3-C2-C1	2.11	119.55	111.80
2	A	606	DM5	O5'-C5'-C4'	2.10	113.34	109.55

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	3502	GOL	O1-C1-C2-C3
3	A	3502	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	3502	GOL	8	0
2	A	606	DM5	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	575/584 (98%)	-0.58	4 (0%) 84 87	3, 10, 24, 52	33 (5%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	70	TRP	3.0
1	A	35[A]	ASN	2.9
1	A	596	ASN	2.9
1	A	34	ASN	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	DM5	A	606	36/36	0.90	0.09	16,25,34,36	0
3	GOL	A	3501	6/6	0.94	0.08	17,19,20,20	0
3	GOL	A	3502	6/6	0.96	0.09	14,28,35,43	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	GOL	A	3500[A]	6/6	0.97	0.06	8,9,10,15	6
3	GOL	A	3500[B]	6/6	0.97	0.06	8,10,12,15	6

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.