



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 07:23 AM UTC

PDB ID : 6ARX / pdb_00006arx
Title : Crystal structure of an insecticide-resistant acetylcholinesterase mutant from the malaria vector *Anopheles gambiae* in the ligand-free state
Authors : Cheung, J.; Mahmood, A.; Kalathur, R.; Lixuan, L.; Carlier, P.R.
Deposited on : 2017-08-23
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	
1	B	542	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FLC	B	1002	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 9033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	537	Total	C	N	O	S	0	4	0
			4289	2729	754	793	13			
1	B	537	Total	C	N	O	S	0	4	0
			4289	2729	754	793	13			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	161	GLY	-	expression tag	UNP Q869C3
A	280	SER	GLY	engineered mutation	UNP Q869C3
B	161	GLY	-	expression tag	UNP Q869C3
B	280	SER	GLY	engineered mutation	UNP Q869C3

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is CITRATE ANION (CCD ID: FLC) (formula: $C_6H_5O_7^-$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	6	7		
3	A	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		
3	B	1	Total	C	O	0	0
			13	6	7		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl^-).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	185	Total 185	O 185	0	0
5	B	175	Total 175	O 175	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	149.60Å 149.60Å 225.86Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.10 – 2.30 49.10 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.10-2.30) 99.7 (49.10-2.30)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.29Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.178 , 0.203 0.180 , 0.204	Depositor DCC
R_{free} test set	6262 reflections (4.62%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.504	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9033	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, FLC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/4414	0.69	3/6022 (0.0%)
1	B	0.55	0/4414	0.68	1/6022 (0.0%)
All	All	0.56	0/8828	0.69	4/12044 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	616	LEU	CA-C-N	-6.32	109.03	121.41
1	A	616	LEU	C-N-CA	-6.32	109.03	121.41
1	A	475	LYS	CA-CB-CG	-6.16	101.77	114.10
1	B	465	GLN	CA-CB-CG	6.05	126.21	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4289	0	4138	26	0
1	B	4289	0	4139	31	0
2	A	28	0	26	0	0
2	B	14	0	13	0	0
3	A	26	0	10	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	26	0	10	2	0
4	A	1	0	0	0	0
5	A	185	0	0	0	0
5	B	175	0	0	7	0
All	All	9033	0	8336	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:292:ASP:O	5:B:1101:HOH:O	1.95	0.84
1:A:517:ARG:NH2	3:A:1004:FLC:OG1	2.15	0.79
1:B:676:ARG:NH2	3:B:1002:FLC:OB2	2.17	0.77
1:B:164:ASP:OD2	5:B:1102:HOH:O	2.05	0.72
1:A:506:THR:HB	1:A:544:PRO:HB2	1.74	0.70
1:A:274:LEU:HD21	1:A:307:LEU:HD23	1.75	0.68
1:B:570[A]:ARG:NH1	1:B:573:GLU:OE1	2.31	0.62
1:B:274:LEU:HD21	1:B:307:LEU:HD23	1.81	0.62
1:B:506:THR:HB	1:B:544:PRO:HB2	1.82	0.61
1:B:190:VAL:HG22	1:B:261:VAL:HG22	1.81	0.61
1:A:190:VAL:HG22	1:A:261:VAL:HG22	1.86	0.57
1:B:393:TRP:HB3	1:B:447:CYS:O	2.06	0.55
1:A:338[B]:ARG:CZ	1:A:378:LEU:HD21	2.39	0.53
1:A:676:ARG:NH2	3:A:1003:FLC:OB1	2.39	0.53
1:B:323:GLU:HG3	5:B:1162:HOH:O	2.10	0.52
1:A:676:ARG:HH22	3:A:1003:FLC:CBC	2.23	0.51
1:A:421:LEU:O	1:A:425:VAL:HG23	2.10	0.51
3:B:1002:FLC:OA2	3:B:1002:FLC:OHB	2.26	0.50
1:A:611:PRO:HA	1:A:618:TYR:CD1	2.47	0.49
1:A:171:ASP:OD2	1:A:343:ASN:ND2	2.42	0.48
1:A:517:ARG:HH22	3:A:1004:FLC:CGC	2.24	0.47
1:A:569:GLN:HE21	1:A:573:GLU:CD	2.22	0.46
1:A:569:GLN:NE2	1:A:573:GLU:OE2	2.38	0.46
1:B:648:SER:HB2	1:B:651:PHE:HB2	1.97	0.46
1:B:525:GLY:O	1:B:529:GLN:HG2	2.16	0.45
1:A:491:ILE:CG2	1:A:499:LEU:HD22	2.47	0.45
1:B:187:LYS:O	1:B:264:ARG:HG3	2.16	0.45
1:B:258:ILE:HD11	1:B:336:ALA:HB1	1.99	0.44
1:B:619:THR:O	1:B:623:LYS:HG3	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:353:ARG:HA	1:A:380:GLN:OE1	2.18	0.44
1:B:611:PRO:HA	1:B:618:TYR:CD1	2.52	0.44
1:A:196:TYR:CD2	1:A:258:ILE:HG13	2.53	0.44
1:B:475:LYS:HG3	5:B:1252:HOH:O	2.18	0.43
3:A:1003:FLC:OA1	3:A:1003:FLC:OHB	2.35	0.43
1:B:280:SER:HB2	1:B:282:TYR:CE2	2.52	0.43
1:B:519:LEU:HD12	1:B:554:MET:HE2	1.99	0.43
1:B:421:LEU:O	1:B:425:VAL:HG23	2.19	0.43
1:B:485:GLU:O	1:B:599:MET:HA	2.18	0.43
1:A:394:ALA:C	1:A:395:LEU:HD12	2.43	0.43
1:A:680:LEU:HD23	1:A:680:LEU:HA	1.88	0.42
1:A:465:GLN:H	1:A:465:GLN:CD	2.27	0.42
1:B:319:LEU:HD12	1:B:425:VAL:HG22	2.00	0.42
1:B:523:VAL:HB	1:B:527:ALA:HB3	2.00	0.42
1:B:670:ASN:N	5:B:1104:HOH:O	2.35	0.42
1:B:669:LEU:HD23	1:B:669:LEU:HA	1.71	0.42
1:A:525:GLY:O	1:A:529:GLN:HG2	2.19	0.42
1:B:357:PHE:CB	1:B:383:ILE:HB	2.51	0.41
1:B:644:PRO:HB3	5:B:1238:HOH:O	2.20	0.41
1:A:523:VAL:HB	1:A:527:ALA:HB3	2.03	0.41
1:A:258:ILE:HD13	1:A:258:ILE:HG21	1.82	0.41
1:A:367:SER:HA	1:A:370:LEU:HD12	2.01	0.41
1:B:165:PRO:HD2	5:B:1102:HOH:O	2.20	0.41
1:B:569:GLN:HE21	1:B:573:GLU:CD	2.29	0.41
1:A:631:ARG:HB3	1:A:651:PHE:HE1	1.85	0.41
1:B:367:SER:HA	1:B:370:LEU:HD12	2.03	0.41
1:A:229:VAL:HG11	1:A:433:PRO:HB2	2.02	0.41
1:B:613:ASN:HB3	1:B:616:LEU:HD12	2.03	0.41
1:A:475:LYS:HE2	1:A:574:GLU:HG2	2.03	0.40
1:B:607:VAL:O	1:B:630:MET:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	539/542 (99%)	520 (96%)	19 (4%)	0	100	100
1	B	539/542 (99%)	520 (96%)	19 (4%)	0	100	100
All	All	1078/1084 (99%)	1040 (96%)	38 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	459/458 (100%)	451 (98%)	8 (2%)	53	72
1	B	459/458 (100%)	451 (98%)	8 (2%)	53	72
All	All	918/916 (100%)	902 (98%)	16 (2%)	51	72

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	SER
1	A	293	HIS
1	A	341	ARG
1	A	357	PHE
1	A	388	SER
1	A	465	GLN
1	A	529	GLN
1	A	629	ILE
1	B	190	VAL
1	B	213	GLU
1	B	293	HIS
1	B	341	ARG
1	B	357	PHE
1	B	388	SER
1	B	465	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	529	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	269	ASN
1	A	576	ASN
1	B	577	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	1002	1	14,14,15	0.83	1 (7%)	17,19,21	0.77	1 (5%)
3	FLC	A	1004	-	12,12,12	1.85	3 (25%)	17,17,17	2.10	4 (23%)
3	FLC	A	1003	-	12,12,12	1.39	1 (8%)	17,17,17	0.81	0
2	NAG	B	1001	1	14,14,15	1.13	1 (7%)	17,19,21	1.31	3 (17%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FLC	B	1003	-	12,12,12	1.62	1 (8%)	17,17,17	0.88	1 (5%)
3	FLC	B	1002	-	12,12,12	1.87	3 (25%)	17,17,17	2.07	6 (35%)
2	NAG	A	1001	1	14,14,15	1.16	1 (7%)	17,19,21	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1002	1	-	4/6/23/26	0/1/1/1
3	FLC	A	1004	-	-	7/16/16/16	-
3	FLC	A	1003	-	-	7/16/16/16	-
2	NAG	B	1001	1	-	2/6/23/26	0/1/1/1
3	FLC	B	1003	-	-	9/16/16/16	-
3	FLC	B	1002	-	-	11/16/16/16	-
2	NAG	A	1001	1	-	2/6/23/26	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1004	FLC	CB-CBC	4.64	1.58	1.53
3	B	1003	FLC	CB-CBC	4.35	1.58	1.53
2	A	1001	NAG	O5-C1	4.11	1.50	1.43
2	B	1001	NAG	O5-C1	4.04	1.50	1.43
3	A	1003	FLC	CB-CBC	3.92	1.57	1.53
3	B	1002	FLC	CB-CBC	3.86	1.57	1.53
3	B	1002	FLC	OB1-CBC	2.93	1.31	1.22
3	A	1004	FLC	OB1-CBC	2.58	1.30	1.22
2	A	1002	NAG	O5-C1	2.41	1.47	1.43
3	B	1002	FLC	OA1-CAC	2.21	1.29	1.22
3	A	1004	FLC	OA1-CAC	2.02	1.28	1.22

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1004	FLC	OB2-CBC-CB	5.73	124.13	113.14
3	A	1004	FLC	OB1-CBC-CB	-4.91	112.58	122.09
3	B	1002	FLC	OB1-CBC-CB	-4.44	113.48	122.09
3	B	1002	FLC	OB2-CBC-CB	4.40	121.59	113.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1001	NAG	C1-O5-C5	3.20	116.48	112.19
3	B	1002	FLC	OA1-CAC-CA	-2.91	114.70	122.95
3	A	1004	FLC	OA2-CAC-CA	2.54	122.41	114.35
2	A	1002	NAG	C1-O5-C5	2.53	115.58	112.19
3	B	1002	FLC	OA2-CAC-CA	2.48	122.21	114.35
3	B	1002	FLC	OG1-CGC-CG	-2.45	116.01	122.95
3	A	1004	FLC	OA1-CAC-CA	-2.37	116.24	122.95
3	B	1002	FLC	OG2-CGC-CG	2.31	121.68	114.35
2	B	1001	NAG	C2-N2-C7	2.30	125.98	122.90
2	B	1001	NAG	C1-C2-N2	2.17	113.85	110.43
3	B	1003	FLC	OG2-CGC-CG	2.14	121.13	114.35

There are no chirality outliers.

All (42) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1001	NAG	C1-C2-N2-C7
3	A	1003	FLC	CG-CB-CBC-OB1
3	A	1003	FLC	CG-CB-CBC-OB2
3	A	1003	FLC	OHB-CB-CBC-OB1
3	A	1003	FLC	OHB-CB-CBC-OB2
3	A	1004	FLC	CG-CB-CBC-OB1
3	A	1004	FLC	CG-CB-CBC-OB2
3	A	1004	FLC	OHB-CB-CBC-OB1
3	A	1004	FLC	OHB-CB-CBC-OB2
3	B	1002	FLC	CAC-CA-CB-CBC
3	B	1002	FLC	CAC-CA-CB-OHB
3	B	1002	FLC	CA-CB-CBC-OB1
3	B	1002	FLC	CA-CB-CBC-OB2
3	B	1002	FLC	OHB-CB-CBC-OB1
3	B	1002	FLC	OHB-CB-CBC-OB2
3	B	1003	FLC	CA-CB-CBC-OB1
3	B	1003	FLC	CA-CB-CBC-OB2
3	B	1003	FLC	OHB-CB-CBC-OB1
3	B	1003	FLC	OHB-CB-CBC-OB2
3	B	1003	FLC	CBC-CB-CG-CGC
3	B	1003	FLC	OHB-CB-CG-CGC
2	A	1002	NAG	C4-C5-C6-O6
2	A	1002	NAG	O5-C5-C6-O6
2	A	1001	NAG	O5-C5-C6-O6
2	A	1001	NAG	C4-C5-C6-O6
3	A	1004	FLC	CAC-CA-CB-CBC

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	1004	FLC	CAC-CA-CB-CG
3	A	1004	FLC	CAC-CA-CB-OHB
3	B	1002	FLC	CAC-CA-CB-CG
3	B	1003	FLC	CA-CB-CG-CGC
2	A	1002	NAG	C8-C7-N2-C2
2	A	1002	NAG	O7-C7-N2-C2
3	A	1003	FLC	CAC-CA-CB-CBC
3	A	1003	FLC	CAC-CA-CB-OHB
3	A	1003	FLC	CAC-CA-CB-CG
3	B	1002	FLC	CB-CA-CAC-OA2
2	B	1001	NAG	C3-C2-N2-C7
3	B	1003	FLC	CB-CA-CAC-OA2
3	B	1002	FLC	CB-CA-CAC-OA1
3	B	1003	FLC	CB-CA-CAC-OA1
3	B	1002	FLC	CB-CG-CGC-OG1
3	B	1002	FLC	CB-CG-CGC-OG2

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1004	FLC	2	0
3	A	1003	FLC	3	0
3	B	1002	FLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	537/542 (99%)	-0.28	17 (3%)	50 52	24, 50, 77, 120	4 (0%)
1	B	537/542 (99%)	-0.28	12 (2%)	62 64	25, 51, 75, 127	4 (0%)
All	All	1074/1084 (99%)	-0.28	29 (2%)	56 58	24, 50, 76, 127	8 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	647	ALA	5.7
1	B	647	ALA	5.6
1	B	660[A]	HIS	5.0
1	B	699	ASN	4.6
1	A	699	ASN	4.5
1	A	185	GLY	4.2
1	A	660[A]	HIS	4.0
1	B	338[A]	ARG	3.7
1	B	651	PHE	3.1
1	A	646	THR	3.0
1	B	649	SER	3.0
1	A	650	GLU	2.9
1	B	570[A]	ARG	2.8
1	B	650	GLU	2.8
1	A	338[A]	ARG	2.8
1	B	163	ASN	2.7
1	A	649	SER	2.6
1	B	268	LYS	2.5
1	B	698	SER	2.5
1	B	646	THR	2.5
1	A	184	SER	2.5
1	A	268	LYS	2.5
1	A	165	PRO	2.4
1	A	163	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	651	PHE	2.3
1	A	617	GLY	2.2
1	A	443	THR	2.1
1	A	183	PRO	2.1
1	A	648	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1002	14/15	0.48	0.21	102,121,131,133	0
3	FLC	B	1003	13/13	0.70	0.18	90,99,106,107	13
3	FLC	A	1003	13/13	0.72	0.22	68,83,91,95	13
3	FLC	B	1002	13/13	0.74	0.19	73,86,98,104	13
3	FLC	A	1004	13/13	0.75	0.17	83,92,99,103	13
2	NAG	B	1001	14/15	0.77	0.17	76,87,93,109	0
2	NAG	A	1001	14/15	0.79	0.14	72,89,96,97	0
4	CL	A	1005	1/1	0.97	0.09	57,57,57,57	0

6.5 Other polymers [i](#)

There are no such residues in this entry.