



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 4, 2026 – 09:39 PM UTC

PDB ID : 8ARP / pdb\_00008arp  
Title : Crystal structure of DEAD-box protein Dbp2 in complex with ADP  
Authors : Song, Q.X.; Rety, S.; Xi, X.G.  
Deposited on : 2022-08-17  
Resolution : 3.05 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Buster-report : wwPDB partial adaption of 1.1.7 (2018)  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	546	
1	B	546	
1	C	546	
1	D	546	
1	E	546	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	546	 53% 28% 19%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	E	2002	-	-	X	-

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 21294 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP-dependent RNA helicase DBP2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	443	Total 3505	C 2202	N 616	O 671	S 16	0	0	0
1	B	444	Total 3516	C 2208	N 620	O 672	S 16	0	0	0
1	C	445	Total 3528	C 2217	N 621	O 674	S 16	0	0	0
1	D	444	Total 3516	C 2208	N 620	O 672	S 16	0	0	0
1	E	444	Total 3516	C 2208	N 620	O 672	S 16	0	0	0
1	F	443	Total 3508	C 2204	N 618	O 670	S 16	0	0	0

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C<sub>10</sub>H<sub>15</sub>N<sub>5</sub>O<sub>10</sub>P<sub>2</sub>) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	B	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	C	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	D	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	E	1	Total 27	C 10	N 5	O 10	P 2	0	0
2	F	1	Total 27	C 10	N 5	O 10	P 2	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
3	A	1	Total 5	O S 4 1	0	0
3	B	1	Total 5	O S 4 1	0	0
3	C	1	Total 5	O S 4 1	0	0
3	D	1	Total 5	O S 4 1	0	0
3	E	1	Total 5	O S 4 1	0	0
3	E	1	Total 5	O S 4 1	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	E	1	Total	O	S	0	0
			5	4	1		
3	F	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		
4	E	1	Total	Mg	0	0
			1	1		






- Molecule 1: ATP-dependent RNA helicase DBP2


Chain C: 47% 34% 18%




- Molecule 1: ATP-dependent RNA helicase DBP2

Chain D: 45% 35% 19%






• Molecule 1: ATP-dependent RNA helicase DBP2



• Molecule 1: ATP-dependent RNA helicase DBP2



HIS  
PRO  
ARG  
TYR  
GLY  
GLY  
ARG  
GLY  
GLY  
ARG  
GLY  
TYR  
GLY  
ARG  
ARG  
GLY  
GLY  
TYR  
GLY  
GLY  
ARG  
GLY  
GLY  
TYR  
GLY  
GLY  
ASN  
ASN  
ARG  
GLN  
ARG  
ASP  
GLY  
GLY  
TRP  
GLY  
ASN  
ARG  
GLY  
ARG  
SER  
ASN  
TYR

## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.94Å 168.84Å 168.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	119.40 – 3.05 119.40 – 3.05	Depositor EDS
% Data completeness (in resolution range)	96.2 (119.40-3.05) 96.1 (119.40-3.05)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 3.07Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.223 , 0.270 0.227 , 0.273	Depositor DCC
$R_{free}$ test set	3779 reflections (4.70%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	77.6	Xtrriage
Anisotropy	0.481	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 125.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.40$ , $\langle L^2 \rangle = 0.22$	Xtrriage
Estimated twinning fraction	0.428 for -h,l,k	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	21294	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	110.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.01% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.18	0/3574	0.50	2/4840 (0.0%)
1	B	0.22	0/3585	0.55	0/4854
1	C	0.22	0/3598	0.61	2/4872 (0.0%)
1	D	0.20	0/3585	0.59	1/4854 (0.0%)
1	E	0.21	0/3585	0.53	0/4854
1	F	0.19	0/3577	0.50	0/4843
All	All	0.20	0/21504	0.55	5/29117 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	D	0	3
1	E	0	2
1	F	0	2
All	All	0	13

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	302	PRO	CA-C-N	8.13	136.33	121.70
1	C	302	PRO	C-N-CA	8.13	136.33	121.70
1	A	55	PRO	CA-C-N	7.25	137.78	125.02
1	A	55	PRO	C-N-CA	7.25	137.78	125.02

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	118	ALA	N-CA-C	-5.30	100.27	108.90

There are no chirality outliers.

5 of 13 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	56	GLN	Peptide
1	A	58	LEU	Peptide
1	B	324	LEU	Peptide
1	B	81	HIS	Peptide
1	C	56	GLN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3505	0	3485	113	0
1	B	3516	0	3498	163	0
1	C	3528	0	3508	170	0
1	D	3516	0	3498	175	0
1	E	3516	0	3497	118	0
1	F	3508	0	3493	122	0
2	A	27	0	12	5	0
2	B	27	0	12	0	0
2	C	27	0	12	1	0
2	D	27	0	12	2	0
2	E	27	0	12	0	0
2	F	27	0	12	1	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	15	0	0	3	0
3	F	5	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	21294	0	21051	821	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 821 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:78:TYR:OH	1:D:110:LYS:NZ	1.95	0.98
1:C:330:HIS:NE2	1:C:481:GLU:OE2	2.08	0.87
1:C:302:PRO:HB2	1:C:304:GLU:H	1.39	0.87
1:C:397:ARG:NH1	1:F:213:SER:O	2.09	0.85
1:E:59:ILE:O	1:E:287:GLN:NE2	2.14	0.80

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	441/546 (81%)	437 (99%)	4 (1%)	0	100 100
1	B	442/546 (81%)	440 (100%)	1 (0%)	1 (0%)	43 70
1	C	443/546 (81%)	432 (98%)	10 (2%)	1 (0%)	43 70
1	D	442/546 (81%)	437 (99%)	4 (1%)	1 (0%)	43 70
1	E	442/546 (81%)	439 (99%)	3 (1%)	0	100 100
1	F	441/546 (81%)	437 (99%)	4 (1%)	0	100 100
All	All	2651/3276 (81%)	2622 (99%)	26 (1%)	3 (0%)	48 75

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	80	GLU
1	C	327	SER
1	B	324	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/450 (85%)	383 (100%)	0	100	100
1	B	384/450 (85%)	384 (100%)	0	100	100
1	C	385/450 (86%)	385 (100%)	0	100	100
1	D	384/450 (85%)	383 (100%)	1 (0%)	86	85
1	E	384/450 (85%)	384 (100%)	0	100	100
1	F	383/450 (85%)	383 (100%)	0	100	100
All	All	2303/2700 (85%)	2302 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	422	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	430	ASN
1	D	106	HIS
1	F	391	HIS
1	D	483	ASN
1	E	446	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 3 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	E	2002	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	A	2002	-	4,4,4	0.22	0	6,6,6	0.12	0
3	SO4	E	2003	-	4,4,4	0.24	0	6,6,6	0.06	0
2	ADP	D	2001	-	28,29,29	1.42	4 (14%)	43,45,45	1.86	9 (20%)
2	ADP	B	2001	-	28,29,29	1.43	4 (14%)	43,45,45	1.90	11 (25%)
2	ADP	F	2001	-	28,29,29	1.42	4 (14%)	43,45,45	1.83	11 (25%)
2	ADP	C	2001	4	28,29,29	1.39	5 (17%)	43,45,45	1.80	9 (20%)
3	SO4	D	2002	-	4,4,4	0.23	0	6,6,6	0.09	0
3	SO4	C	2002	-	4,4,4	0.23	0	6,6,6	0.14	0
3	SO4	B	2002	-	4,4,4	0.24	0	6,6,6	0.09	0
3	SO4	F	2002	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	E	2004	-	4,4,4	0.23	0	6,6,6	0.10	0
2	ADP	E	2001	-	28,29,29	1.43	4 (14%)	43,45,45	1.86	9 (20%)
2	ADP	A	2001	4	28,29,29	1.42	4 (14%)	43,45,45	1.85	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	D	2001	-	-	5/16/32/32	0/3/3/3
2	ADP	B	2001	-	-	4/16/32/32	0/3/3/3
2	ADP	F	2001	-	-	4/16/32/32	0/3/3/3
2	ADP	C	2001	4	-	5/16/32/32	0/3/3/3
2	ADP	E	2001	-	-	4/16/32/32	0/3/3/3
2	ADP	A	2001	4	-	4/16/32/32	0/3/3/3

The worst 5 of 25 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	2001	ADP	C5-C4	4.86	1.47	1.39
2	A	2001	ADP	C5-C4	4.85	1.47	1.39
2	B	2001	ADP	C5-C4	4.83	1.47	1.39
2	D	2001	ADP	C5-C4	4.82	1.47	1.39
2	F	2001	ADP	C5-C4	4.69	1.47	1.39

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2001	ADP	C5-C4-N3	-6.16	118.24	126.72
2	E	2001	ADP	C5-C4-N3	-5.93	118.55	126.72
2	F	2001	ADP	C5-C4-N3	-5.85	118.66	126.72
2	D	2001	ADP	C5-C4-N3	-5.82	118.71	126.72
2	B	2001	ADP	C5-C4-N3	-5.62	118.98	126.72

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	ADP	C5'-O5'-PA-O1A
2	A	2001	ADP	C4'-C5'-O5'-PA
2	A	2001	ADP	O4'-C4'-C5'-O5'
2	A	2001	ADP	C3'-C4'-C5'-O5'
2	B	2001	ADP	C5'-O5'-PA-O1A

There are no ring outliers.

6 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	E	2002	SO4	2	0



*Continued on next page...*



Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	2001	ADP	2	0
2	F	2001	ADP	1	0
2	C	2001	ADP	1	0
3	E	2004	SO4	1	0
2	A	2001	ADP	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	443/546 (81%)	-0.92	0	100   100	44, 79, 135, 213	0
1	B	444/546 (81%)	-0.47	0	100   100	81, 127, 188, 295	0
1	C	445/546 (81%)	-0.61	1 (0%)	91   83	47, 104, 181, 267	0
1	D	444/546 (81%)	-0.63	1 (0%)	91   83	69, 118, 174, 280	0
1	E	444/546 (81%)	-0.66	0	100   100	53, 106, 166, 274	0
1	F	443/546 (81%)	-0.73	0	100   100	46, 93, 164, 266	0
All	All	2663/3276 (81%)	-0.67	2 (0%)	92   86	44, 106, 175, 295	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	327	SER	2.7
1	D	454	ALA	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)



In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	ADP	B	2001	27/27	0.98	0.05	76,105,130,146	0
3	SO4	B	2002	5/5	0.98	0.10	98,99,110,123	0
3	SO4	E	2003	5/5	0.98	0.08	116,119,133,138	0
3	SO4	E	2004	5/5	0.98	0.07	89,106,119,140	0
3	SO4	F	2002	5/5	0.98	0.09	52,63,70,72	0
4	MG	C	2003	1/1	0.98	0.08	141,141,141,141	0
3	SO4	A	2002	5/5	0.99	0.06	62,94,101,154	0
2	ADP	A	2001	27/27	0.99	0.05	56,72,87,92	0
3	SO4	C	2002	5/5	0.99	0.07	51,65,87,94	0
3	SO4	D	2002	5/5	0.99	0.06	62,74,81,84	0
3	SO4	E	2002	5/5	0.99	0.05	79,81,85,85	0
2	ADP	C	2001	27/27	0.99	0.06	71,91,108,111	0
2	ADP	D	2001	27/27	0.99	0.04	78,95,129,132	0
2	ADP	E	2001	27/27	0.99	0.05	81,94,101,103	0
4	MG	A	2003	1/1	0.99	0.04	47,47,47,47	0
2	ADP	F	2001	27/27	0.99	0.04	62,74,86,86	0
4	MG	E	2005	1/1	0.99	0.03	88,88,88,88	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around ADP B 2001:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around ADP A 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ADP C 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around ADP D 2001:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around ADP E 2001:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around MG A 2003:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.