



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 07:40 AM UTC

PDB ID : 4ATN / pdb_00004atn
Title : Crystal structure of C2498 2'-O-ribose methyltransferase RlmM from Escherichia coli
Authors : Puneekar, A.S.; Shepherd, T.R.; Liljeruhm, J.; Forster, A.C.; Selmer, M.
Deposited on : 2012-05-08
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3494 (1.96-1.96)
Clashscore	190562	3612 (1.96-1.96)
Ramachandran outliers	187476	3587 (1.96-1.96)
Sidechain outliers	187428	3587 (1.96-1.96)
RSRZ outliers	180081	3495 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBOSOMAL RNA LARGE SUBUNIT METHYLTRANSFERASE M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	7	0
			2916	1872	502	519	23			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	367	SER	-	expression tag	UNP P0ADR6
A	368	LYS	-	expression tag	UNP P0ADR6
A	369	GLY	-	expression tag	UNP P0ADR6
A	370	HIS	-	expression tag	UNP P0ADR6
A	371	HIS	-	expression tag	UNP P0ADR6
A	372	HIS	-	expression tag	UNP P0ADR6
A	373	HIS	-	expression tag	UNP P0ADR6
A	374	HIS	-	expression tag	UNP P0ADR6
A	375	HIS	-	expression tag	UNP P0ADR6

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	195	Total	O	0	0
			195	195		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RIBOSOMAL RNA LARGE SUBUNIT METHYLTRANSFERASE M



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	92.09Å 92.09Å 83.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.77 – 1.95 28.77 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (28.77-1.95) 99.7 (28.77-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 1.95Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.169 , 0.206 0.164 , 0.199	Depositor DCC
R_{free} test set	1502 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3245	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSS, SO4, EDO, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	0/3002	0.73	0/4068

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2916	0	2893	55	0
2	A	30	0	40	4	0
3	A	84	0	126	11	0
4	A	20	0	0	1	0
5	A	195	0	0	5	0
All	All	3245	0	3059	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:ASP:H	3:A:1371:EDO:H22	1.37	0.85
1:A:309:MET:HB2	3:A:1368:EDO:H12	1.56	0.85
1:A:61:PHE:H	1:A:167:ASN:HD21	1.34	0.74
4:A:1387:SO4:O4	5:A:2122:HOH:O	2.07	0.72
1:A:166:ASN:HD22	1:A:166:ASN:H	1.37	0.72
1:A:335:GLN:HE22	1:A:353:ARG:HH21	1.38	0.68
1:A:69:GLN:HE22	1:A:169:SER:H	1.44	0.67
1:A:61:PHE:H	1:A:167:ASN:ND2	1.97	0.63
1:A:241:ASN:O	1:A:258:ARG:NH1	2.33	0.61
1:A:69:GLN:HE21	1:A:70:TRP:H	1.48	0.61
1:A:270:ASN:H	1:A:270:ASN:HD22	1.47	0.60
1:A:117[A]:LEU:CD1	1:A:121:ARG:NH2	2.65	0.59
1:A:268:ARG:HH11	1:A:270:ASN:HD21	1.51	0.58
1:A:60:PRO:HB3	3:A:1364:EDO:H12	1.86	0.58
1:A:69:GLN:NE2	1:A:169:SER:H	2.00	0.57
1:A:107:VAL:HG22	1:A:117[A]:LEU:HD23	1.86	0.57
1:A:337:GLN:HE21	1:A:351:HIS:HD2	1.51	0.57
3:A:1375:EDO:O1	5:A:2195:HOH:O	2.17	0.56
1:A:106:GLU:OE1	1:A:149:HIS:HD2	1.90	0.55
1:A:337:GLN:HG3	5:A:2185:HOH:O	2.09	0.53
1:A:59:LEU:HD12	1:A:60:PRO:HD2	1.91	0.53
1:A:328:ASP:OD2	5:A:2183:HOH:O	2.19	0.52
1:A:15:GLU:OE2	1:A:312:ARG:HD2	2.11	0.51
1:A:211:ASN:HD22	1:A:211:ASN:C	2.20	0.50
1:A:50:ASP:HB2	3:A:1372:EDO:H21	1.93	0.50
1:A:69:GLN:HE22	1:A:168:ASN:HB2	1.77	0.50
1:A:72:VAL:O	1:A:161:GLY:HA3	2.12	0.49
1:A:77:LEU:HB2	1:A:92[B]:MET:SD	2.53	0.49
1:A:168:ASN:C	1:A:168:ASN:HD22	2.21	0.49
1:A:176:PRO:HG2	1:A:194:GLU:HG3	1.94	0.49
1:A:97:VAL:HA	2:A:1360:GOL:H11	1.95	0.48
1:A:58:GLU:OE1	5:A:2030:HOH:O	2.20	0.48
1:A:51:ASP:N	3:A:1372:EDO:H12	2.29	0.48
1:A:312:ARG:NH2	1:A:346[A]:GLU:OE2	2.47	0.48
1:A:320:LEU:HD13	2:A:1361:GOL:H11	1.96	0.48
1:A:168:ASN:C	1:A:168:ASN:ND2	2.73	0.47
1:A:149:HIS:HE1	1:A:163:SER:OG	1.99	0.46
1:A:171:PHE:CE1	2:A:1362:GOL:H2	2.51	0.46
1:A:88:PRO:O	1:A:92[B]:MET:HG3	2.16	0.46
1:A:166:ASN:HD21	3:A:1376:EDO:H12	1.80	0.46
1:A:117[A]:LEU:HD11	1:A:121:ARG:NH2	2.31	0.45
1:A:117[A]:LEU:HD11	1:A:121:ARG:HH21	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:332:ILE:HA	3:A:1381:EDO:H21	1.98	0.44
1:A:223:GLY:HA2	1:A:244[A]:MET:HE1	1.99	0.44
1:A:294:VAL:HA	1:A:332:ILE:HD13	2.00	0.43
1:A:117[A]:LEU:HD12	1:A:121:ARG:NH2	2.32	0.43
1:A:117[A]:LEU:CD1	1:A:121:ARG:CZ	2.97	0.43
1:A:270:ASN:HD22	1:A:270:ASN:N	2.11	0.43
1:A:117[A]:LEU:CD1	1:A:121:ARG:HH21	2.32	0.42
1:A:260:ASP:N	3:A:1371:EDO:H22	2.18	0.42
1:A:308:PRO:HB2	3:A:1368:EDO:H21	2.02	0.42
1:A:146:PRO:HG2	2:A:1360:GOL:H2	2.01	0.42
1:A:259:GLU:HB3	3:A:1371:EDO:O2	2.20	0.42
1:A:240:ASP:O	1:A:258:ARG:HA	2.19	0.41
1:A:181:PRO:HG2	1:A:184:ALA:HB2	2.01	0.41
1:A:1:MET:HE1	1:A:162:TYR:HE2	1.86	0.41
1:A:335:GLN:NE2	1:A:353:ARG:HE	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/375 (96%)	353 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/318 (98%)	297 (96%)	14 (4%)	24	14

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	94	GLN
1	A	142	THR
1	A	166	ASN
1	A	168	ASN
1	A	211	ASN
1	A	265	ARG
1	A	270	ASN
1	A	280	GLU
1	A	304	ASN
1	A	305	LEU
1	A	315	GLU
1	A	329	GLU
1	A	354	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	26	GLN
1	A	69	GLN
1	A	94	GLN
1	A	139	ASN
1	A	149	HIS
1	A	166	ASN
1	A	167	ASN
1	A	168	ASN
1	A	211	ASN
1	A	228	GLN
1	A	233	ASN
1	A	246	GLN
1	A	270	ASN
1	A	291	GLN
1	A	295	ASN
1	A	304	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	326	GLN
1	A	335	GLN
1	A	337	GLN
1	A	340	GLN
1	A	351	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSS	A	157	1	4,6,7	1.10	0	2,6,8	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	157	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	1379	-	3,3,3	0.50	0	2,2,2	0.25	0
3	EDO	A	1381	-	3,3,3	0.42	0	2,2,2	0.42	0
3	EDO	A	1382	-	3,3,3	0.43	0	2,2,2	0.46	0
3	EDO	A	1368	-	3,3,3	0.45	0	2,2,2	0.23	0
2	GOL	A	1358	-	5,5,5	0.31	0	5,5,5	0.42	0
3	EDO	A	1380	-	3,3,3	0.50	0	2,2,2	0.21	0
2	GOL	A	1359	-	5,5,5	0.44	0	5,5,5	0.28	0
3	EDO	A	1369	-	3,3,3	0.38	0	2,2,2	0.40	0
3	EDO	A	1373	-	3,3,3	0.49	0	2,2,2	0.31	0
3	EDO	A	1370	-	3,3,3	0.43	0	2,2,2	0.51	0
3	EDO	A	1364	-	3,3,3	0.42	0	2,2,2	0.45	0
3	EDO	A	1376	-	3,3,3	0.31	0	2,2,2	0.50	0
3	EDO	A	1365	-	3,3,3	0.45	0	2,2,2	0.27	0
3	EDO	A	1377	-	3,3,3	0.47	0	2,2,2	0.34	0
3	EDO	A	1378	-	3,3,3	0.42	0	2,2,2	0.31	0
3	EDO	A	1366	-	3,3,3	0.47	0	2,2,2	0.45	0
2	GOL	A	1360	-	5,5,5	0.31	0	5,5,5	0.55	0
3	EDO	A	1374	-	3,3,3	0.43	0	2,2,2	0.42	0
4	SO4	A	1384	-	4,4,4	0.84	0	6,6,6	0.14	0
4	SO4	A	1385	-	4,4,4	0.83	0	6,6,6	0.08	0
3	EDO	A	1375	-	3,3,3	0.39	0	2,2,2	0.43	0
4	SO4	A	1386	-	4,4,4	0.84	0	6,6,6	0.07	0
3	EDO	A	1371	-	3,3,3	0.48	0	2,2,2	0.32	0
4	SO4	A	1387	-	4,4,4	0.91	0	6,6,6	0.08	0
2	GOL	A	1362	-	5,5,5	0.37	0	5,5,5	0.25	0
3	EDO	A	1372	-	3,3,3	0.45	0	2,2,2	0.30	0
2	GOL	A	1361	-	5,5,5	0.42	0	5,5,5	0.29	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	A	1367	-	3,3,3	0.54	0	2,2,2	0.12	0
3	EDO	A	1383	-	3,3,3	0.44	0	2,2,2	0.39	0
3	EDO	A	1363	-	3,3,3	0.42	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	1379	-	-	0/1/1/1	-
3	EDO	A	1381	-	-	0/1/1/1	-
3	EDO	A	1382	-	-	0/1/1/1	-
3	EDO	A	1368	-	-	1/1/1/1	-
2	GOL	A	1358	-	-	0/4/4/4	-
3	EDO	A	1380	-	-	0/1/1/1	-
2	GOL	A	1359	-	-	2/4/4/4	-
3	EDO	A	1369	-	-	1/1/1/1	-
3	EDO	A	1373	-	-	0/1/1/1	-
3	EDO	A	1370	-	-	1/1/1/1	-
3	EDO	A	1364	-	-	0/1/1/1	-
3	EDO	A	1376	-	-	1/1/1/1	-
3	EDO	A	1365	-	-	1/1/1/1	-
3	EDO	A	1377	-	-	0/1/1/1	-
3	EDO	A	1378	-	-	0/1/1/1	-
3	EDO	A	1366	-	-	0/1/1/1	-
2	GOL	A	1360	-	-	2/4/4/4	-
3	EDO	A	1374	-	-	1/1/1/1	-
3	EDO	A	1375	-	-	0/1/1/1	-
3	EDO	A	1371	-	-	0/1/1/1	-
2	GOL	A	1362	-	-	4/4/4/4	-
3	EDO	A	1372	-	-	0/1/1/1	-
2	GOL	A	1361	-	-	0/4/4/4	-
3	EDO	A	1367	-	-	1/1/1/1	-
3	EDO	A	1383	-	-	1/1/1/1	-
3	EDO	A	1363	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1360	GOL	C1-C2-C3-O3
2	A	1362	GOL	O1-C1-C2-C3
2	A	1362	GOL	C1-C2-C3-O3
3	A	1376	EDO	O1-C1-C2-O2
2	A	1362	GOL	O1-C1-C2-O2
3	A	1369	EDO	O1-C1-C2-O2
2	A	1359	GOL	O1-C1-C2-C3
2	A	1359	GOL	O1-C1-C2-O2
2	A	1360	GOL	O2-C2-C3-O3
3	A	1368	EDO	O1-C1-C2-O2
3	A	1383	EDO	O1-C1-C2-O2
2	A	1362	GOL	O2-C2-C3-O3
3	A	1363	EDO	O1-C1-C2-O2
3	A	1370	EDO	O1-C1-C2-O2
3	A	1374	EDO	O1-C1-C2-O2
3	A	1365	EDO	O1-C1-C2-O2
3	A	1367	EDO	O1-C1-C2-O2

There are no ring outliers.

11 monomers are involved in 16 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1381	EDO	1	0
3	A	1368	EDO	2	0
3	A	1364	EDO	1	0
3	A	1376	EDO	1	0
2	A	1360	GOL	2	0
3	A	1375	EDO	1	0
3	A	1371	EDO	3	0
4	A	1387	SO4	1	0
2	A	1362	GOL	1	0
3	A	1372	EDO	2	0
2	A	1361	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	356/375 (94%)	-0.33	3 (0%) 82 86	18, 31, 49, 60	7 (1%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	140	TYR	2.6
1	A	355	ILE	2.2
1	A	267	THR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSS	A	157	7/8	0.93	0.07	22,27,49,62	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	1385	5/5	0.48	0.14	74,80,92,98	0
3	EDO	A	1382	4/4	0.64	0.19	52,60,62,65	0
3	EDO	A	1383	4/4	0.68	0.14	57,61,63,66	0
3	EDO	A	1377	4/4	0.71	0.13	51,52,54,55	0
4	SO4	A	1386	5/5	0.75	0.13	67,72,78,82	0
4	SO4	A	1387	5/5	0.76	0.10	68,70,83,89	0
3	EDO	A	1370	4/4	0.79	0.11	42,44,48,58	0
3	EDO	A	1371	4/4	0.79	0.15	47,54,56,58	0
3	EDO	A	1378	4/4	0.80	0.14	45,56,57,60	0
3	EDO	A	1373	4/4	0.81	0.15	51,51,54,54	0
3	EDO	A	1380	4/4	0.82	0.14	41,47,62,63	0
3	EDO	A	1374	4/4	0.82	0.11	52,53,55,59	0
3	EDO	A	1376	4/4	0.84	0.16	43,47,54,57	0
3	EDO	A	1381	4/4	0.85	0.13	42,53,54,67	0
3	EDO	A	1365	4/4	0.85	0.18	48,51,52,58	0
3	EDO	A	1379	4/4	0.87	0.12	47,50,52,53	0
2	GOL	A	1362	6/6	0.87	0.10	40,48,56,57	0
3	EDO	A	1367	4/4	0.88	0.18	32,47,49,49	0
3	EDO	A	1363	4/4	0.88	0.13	48,53,58,64	0
3	EDO	A	1364	4/4	0.89	0.14	40,45,50,55	0
3	EDO	A	1372	4/4	0.89	0.12	40,41,51,52	0
3	EDO	A	1368	4/4	0.89	0.11	46,48,50,52	0
3	EDO	A	1369	4/4	0.89	0.12	39,48,54,55	0
3	EDO	A	1366	4/4	0.89	0.13	43,50,51,63	0
2	GOL	A	1359	6/6	0.91	0.11	39,41,47,47	0
2	GOL	A	1360	6/6	0.91	0.10	35,39,44,45	0
3	EDO	A	1375	4/4	0.92	0.13	35,37,41,51	0
2	GOL	A	1361	6/6	0.93	0.11	32,36,41,42	0
2	GOL	A	1358	6/6	0.95	0.08	28,32,34,36	0
4	SO4	A	1384	5/5	0.97	0.07	31,35,37,37	5

6.5 Other polymers

There are no such residues in this entry.