



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:24 PM UTC

PDB ID : 4ATP / pdb_00004atp
Title : Structure of GABA-transaminase A1R958 from *Arthrobacter aureus* in complex with PLP
Authors : Bruce, H.; Tuan, A.N.; Mangas Sanchez, J.; Hart, S.; Turkenburg, J.P.; Grogan, G.
Deposited on : 2012-05-09
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	 87% 8% . .
1	B	456	 87% 8% . . .
1	C	456	 85% 11% . .
1	D	456	 85% 10% . .
1	E	456	 86% 9% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	456	85% 9%
1	G	456	% 84% 11%
1	H	456	% 85% 10%
1	I	456	86% 9%
1	J	456	% 85% 10%
1	K	456	3% 85% 9%
1	L	456	84% 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 39107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-AMINOBUTYRATE TRANSAMINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	442	Total	C	N	O	S	0	1	0
			3242	2056	565	607	14			
1	B	442	Total	C	N	O	S	0	0	0
			3217	2045	557	601	14			
1	C	442	Total	C	N	O	S	0	0	0
			3225	2046	561	604	14			
1	D	439	Total	C	N	O	S	0	0	0
			3205	2030	559	602	14			
1	E	442	Total	C	N	O	S	0	0	0
			3226	2044	563	605	14			
1	F	439	Total	C	N	O	S	0	1	0
			3206	2035	560	597	14			
1	G	440	Total	C	N	O	S	0	0	0
			3130	1984	542	590	14			
1	H	440	Total	C	N	O	S	0	0	0
			3161	2006	549	592	14			
1	I	441	Total	C	N	O	S	0	0	0
			3190	2023	556	597	14			
1	J	438	Total	C	N	O	S	0	0	0
			3152	2000	553	585	14			
1	K	439	Total	C	N	O	S	0	0	0
			3130	1980	543	593	14			
1	L	436	Total	C	N	O	S	0	0	0
			3097	1964	536	583	14			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	THR	ALA	SEE REMARK 999	UNP A1R958
B	113	THR	ALA	SEE REMARK 999	UNP A1R958
C	113	THR	ALA	SEE REMARK 999	UNP A1R958
D	113	THR	ALA	SEE REMARK 999	UNP A1R958
E	113	THR	ALA	SEE REMARK 999	UNP A1R958

Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	113	THR	ALA	SEE REMARK 999	UNP A1R958
G	113	THR	ALA	SEE REMARK 999	UNP A1R958
H	113	THR	ALA	SEE REMARK 999	UNP A1R958
I	113	THR	ALA	SEE REMARK 999	UNP A1R958
J	113	THR	ALA	SEE REMARK 999	UNP A1R958
K	113	THR	ALA	SEE REMARK 999	UNP A1R958
L	113	THR	ALA	SEE REMARK 999	UNP A1R958

- PLP
-
- The diagram shows the chemical structure of Pyridoxal Phosphate (PLP). It consists of a pyridine ring with a phosphate group at the 3-position and an aldehyde group at the 4-position. The pyridine ring is colored blue, with the nitrogen atom labeled N1. The carbon atoms are labeled C2, C3, C4, C5, and C6. The phosphate group is colored purple, with the phosphorus atom labeled P. The oxygen atoms are labeled O1P, O2P, O3P, O4P, and O5A. The aldehyde group is colored red, with the carbonyl oxygen labeled O4A and the hydroxyl group labeled O3. The hydroxyl group at the 3-position is colored red, with the oxygen labeled O3. The carbon atom of the hydroxyl group is labeled C3A. The carbon atom of the aldehyde group is labeled C4A. The carbon atom of the phosphate group is labeled C5A. The carbon atom of the pyridine ring is labeled C2A. The carbon atom of the pyridine ring is labeled C3. The carbon atom of the pyridine ring is labeled C4. The carbon atom of the pyridine ring is labeled C5. The carbon atom of the pyridine ring is labeled C6. The nitrogen atom of the pyridine ring is labeled N1. The phosphorus atom of the phosphate group is labeled P. The oxygen atoms of the phosphate group are labeled O1P, O2P, O3P, O4P, and O5A. The oxygen atom of the aldehyde group is labeled O4A. The oxygen atom of the hydroxyl group at the 3-position is labeled O3. The carbon atom of the hydroxyl group at the 3-position is labeled C3A. The carbon atom of the aldehyde group is labeled C4A. The carbon atom of the phosphate group is labeled C5A. The carbon atom of the pyridine ring is labeled C2A. The carbon atom of the pyridine ring is labeled C3. The carbon atom of the pyridine ring is labeled C4. The carbon atom of the pyridine ring is labeled C5. The carbon atom of the pyridine ring is labeled C6. The nitrogen atom of the pyridine ring is labeled N1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	C	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	D	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	E	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	F	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	G	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	H	1	Total 15	C 8	N 1	O 5	P 1	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	I	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	J	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	K	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	L	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total	O	0	0
			120	120		
3	B	105	Total	O	0	0
			105	105		
3	C	95	Total	O	0	0
			95	95		
3	D	63	Total	O	0	0
			63	63		
3	E	110	Total	O	0	0
			110	110		
3	F	85	Total	O	0	0
			85	85		
3	G	38	Total	O	0	0
			38	38		
3	H	28	Total	O	0	0
			28	28		
3	I	32	Total	O	0	0
			32	32		
3	J	24	Total	O	0	0
			24	24		
3	K	25	Total	O	0	0
			25	25		
3	L	21	Total	O	0	0
			21	21		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: 4-AMINOBTYRATE TRANSAMINASE




• Molecule 1: 4-AMINOBTYRATE TRANSAMINASE




• Molecule 1: 4-AMINOBTYRATE TRANSAMINASE



• Molecule 1: 4-AMINOBTYRATE TRANSAMINASE





• Molecule 1: 4-AMINO BUTYRATE TRANSAMINASE

Chain E: 86% 9% . .



• Molecule 1: 4-AMINO BUTYRATE TRANSAMINASE

Chain F: 85% 9% . .



• Molecule 1: 4-AMINO BUTYRATE TRANSAMINASE

Chain G: 84% 11% . .



• Molecule 1: 4-AMINO BUTYRATE TRANSAMINASE

Chain H: 85% 10% . .





• Molecule 1: 4-AMINOBUTYRATE TRANSAMINASE

Chain I: 86% 9% . .




• Molecule 1: 4-AMINOBUTYRATE TRANSAMINASE

Chain J: 85% 10% . .



• Molecule 1: 4-AMINOBUTYRATE TRANSAMINASE

Chain K: 85% 9% . .



• Molecule 1: 4-AMINOBUTYRATE TRANSAMINASE

Chain L: 84% 10% . .



RES
ALA

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	175.70Å 290.99Å 104.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	89.84 – 2.80 89.84 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (89.84-2.80) 100.0 (89.84-2.80)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.82Å)	Xtriage
Refinement program	REFMAC 5.6.0119	Depositor
R, R_{free}	0.193 , 0.229 0.193 , 0.226	Depositor DCC
R_{free} test set	6667 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	39107	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	4/3305 (0.1%)	0.91	2/4492 (0.0%)
1	B	0.95	0/3275	0.94	7/4452 (0.2%)
1	C	0.92	3/3283 (0.1%)	0.93	2/4462 (0.0%)
1	D	0.92	2/3264 (0.1%)	0.91	4/4439 (0.1%)
1	E	0.92	0/3285	0.93	3/4468 (0.1%)
1	F	0.92	3/3268 (0.1%)	0.90	1/4445 (0.0%)
1	G	0.87	2/3188 (0.1%)	0.91	2/4351 (0.0%)
1	H	0.85	1/3220 (0.0%)	0.90	1/4387 (0.0%)
1	I	0.84	2/3249 (0.1%)	0.89	1/4425 (0.0%)
1	J	0.85	1/3211 (0.0%)	0.89	1/4376 (0.0%)
1	K	0.85	0/3188	0.90	3/4349 (0.1%)
1	L	0.83	0/3155	0.89	2/4306 (0.0%)
All	All	0.89	18/38891 (0.0%)	0.91	29/52952 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	283	HIS	CG-CD2	7.05	1.43	1.35
1	A	455	HIS	CG-CD2	6.41	1.43	1.35
1	C	455	HIS	CG-CD2	6.13	1.42	1.35
1	D	455	HIS	ND1-CE1	5.89	1.38	1.32
1	A	455	HIS	ND1-CE1	5.78	1.38	1.32

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	50	VAL	CB-CA-C	-7.05	100.54	111.26
1	A	376	VAL	CB-CA-C	-6.92	102.19	110.84
1	D	50	VAL	CB-CA-C	-6.69	101.08	111.26
1	A	50	VAL	CB-CA-C	-6.57	101.28	111.26
1	J	50	VAL	CB-CA-C	-6.36	101.59	111.26

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3242	0	3239	31	0
1	B	3217	0	3214	33	0
1	C	3225	0	3218	33	0
1	D	3205	0	3188	39	0
1	E	3226	0	3218	46	0
1	F	3206	0	3191	46	0
1	G	3130	0	3047	46	0
1	H	3161	0	3114	44	0
1	I	3190	0	3157	42	0
1	J	3152	0	3117	45	0
1	K	3130	0	3036	40	0
1	L	3097	0	3017	42	0
2	A	15	0	7	0	0
2	B	15	0	6	0	0
2	C	15	0	6	0	0
2	D	15	0	6	0	0
2	E	15	0	6	0	0
2	F	15	0	6	0	0
2	G	15	0	6	2	0
2	H	15	0	6	0	0
2	I	15	0	6	2	0
2	J	15	0	6	0	0
2	K	15	0	6	1	0
2	L	15	0	6	0	0
3	A	120	0	0	3	0
3	B	105	0	0	1	0
3	C	95	0	0	2	0
3	D	63	0	0	3	0
3	E	110	0	0	1	0
3	F	85	0	0	2	0
3	G	38	0	0	3	0
3	H	28	0	0	2	0
3	I	32	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	24	0	0	1	0
3	K	25	0	0	1	0
3	L	21	0	0	4	0
All	All	39107	0	37829	389	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 389 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:281:VAL:HG22	1:E:286:VAL:HG22	1.34	1.09
1:C:248:PHE:C	1:C:249:LEU:CA	2.33	1.01
1:F:14:GLN:HE22	1:F:56:GLY:H	1.12	0.96
1:D:14:GLN:HE22	1:D:56:GLY:H	1.12	0.96
1:E:281:VAL:CG2	1:E:286:VAL:HG22	1.98	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/456 (96%)	417 (95%)	19 (4%)	3 (1%)	18	47
1	B	438/456 (96%)	417 (95%)	18 (4%)	3 (1%)	18	47
1	C	436/456 (96%)	415 (95%)	18 (4%)	3 (1%)	18	47
1	D	435/456 (95%)	414 (95%)	18 (4%)	3 (1%)	18	47
1	E	438/456 (96%)	417 (95%)	17 (4%)	4 (1%)	14	41
1	F	436/456 (96%)	415 (95%)	18 (4%)	3 (1%)	18	47
1	G	436/456 (96%)	413 (95%)	20 (5%)	3 (1%)	18	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	436/456 (96%)	414 (95%)	19 (4%)	3 (1%)	18	47
1	I	437/456 (96%)	414 (95%)	20 (5%)	3 (1%)	18	47
1	J	434/456 (95%)	412 (95%)	19 (4%)	3 (1%)	18	47
1	K	435/456 (95%)	413 (95%)	19 (4%)	3 (1%)	18	47
1	L	432/456 (95%)	411 (95%)	18 (4%)	3 (1%)	18	47
All	All	5232/5472 (96%)	4972 (95%)	223 (4%)	37 (1%)	18	47

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	367	ALA
1	A	295	LYS
1	B	294	ALA
1	B	295	LYS
1	C	294	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/336 (96%)	307 (96%)	14 (4%)	25	59
1	B	314/336 (94%)	301 (96%)	13 (4%)	27	62
1	C	316/336 (94%)	305 (96%)	11 (4%)	32	67
1	D	315/336 (94%)	304 (96%)	11 (4%)	32	67
1	E	318/336 (95%)	303 (95%)	15 (5%)	23	57
1	F	312/336 (93%)	302 (97%)	10 (3%)	34	70
1	G	297/336 (88%)	286 (96%)	11 (4%)	30	65
1	H	304/336 (90%)	294 (97%)	10 (3%)	33	69
1	I	309/336 (92%)	299 (97%)	10 (3%)	34	70
1	J	305/336 (91%)	294 (96%)	11 (4%)	31	66
1	K	297/336 (88%)	284 (96%)	13 (4%)	25	59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	295/336 (88%)	284 (96%)	11 (4%)	30	65
All	All	3703/4032 (92%)	3563 (96%)	140 (4%)	29	64

5 of 140 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	412	LEU
1	K	113	THR
1	L	61	VAL
1	D	398	GLU
1	D	393	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	283	HIS
1	J	19	ASN
1	K	353	HIS
1	E	345	HIS
1	E	283	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	J	500	1	15,15,16	3.39	4 (26%)	21,22,23	1.84	7 (33%)
2	PLP	F	500	1	15,15,16	2.85	3 (20%)	21,22,23	2.08	7 (33%)
2	PLP	K	500	1	15,15,16	3.43	3 (20%)	21,22,23	2.12	6 (28%)
2	PLP	G	500	1	15,15,16	3.07	4 (26%)	21,22,23	2.08	9 (42%)
2	PLP	C	500	1	15,15,16	2.81	3 (20%)	21,22,23	1.88	6 (28%)
2	PLP	E	500	1	15,15,16	3.42	3 (20%)	21,22,23	1.92	7 (33%)
2	PLP	H	500	1	15,15,16	3.70	4 (26%)	21,22,23	1.95	8 (38%)
2	PLP	L	500	1	15,15,16	2.82	3 (20%)	21,22,23	1.48	4 (19%)
2	PLP	A	500	1	15,15,16	2.93	3 (20%)	21,22,23	1.56	3 (14%)
2	PLP	I	500	1	15,15,16	2.75	3 (20%)	21,22,23	3.64	8 (38%)
2	PLP	D	500	1	15,15,16	3.69	5 (33%)	21,22,23	1.97	7 (33%)
2	PLP	B	500	1	15,15,16	3.27	4 (26%)	21,22,23	2.31	8 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	J	500	1	-	0/6/6/8	0/1/1/1
2	PLP	F	500	1	-	3/6/6/8	0/1/1/1
2	PLP	K	500	1	-	2/6/6/8	0/1/1/1
2	PLP	G	500	1	-	2/6/6/8	0/1/1/1
2	PLP	C	500	1	-	0/6/6/8	0/1/1/1
2	PLP	E	500	1	-	5/6/6/8	0/1/1/1
2	PLP	H	500	1	-	0/6/6/8	0/1/1/1
2	PLP	L	500	1	-	2/6/6/8	0/1/1/1
2	PLP	A	500	1	-	0/6/6/8	0/1/1/1
2	PLP	I	500	1	-	3/6/6/8	0/1/1/1
2	PLP	D	500	1	-	0/6/6/8	0/1/1/1
2	PLP	B	500	1	-	0/6/6/8	0/1/1/1

The worst 5 of 42 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	500	PLP	C3-C2	10.40	1.51	1.41
2	D	500	PLP	C3-C2	9.81	1.51	1.41
2	J	500	PLP	C5-C4	9.73	1.51	1.40
2	E	500	PLP	C3-C2	9.67	1.51	1.41
2	K	500	PLP	C5-C4	9.41	1.51	1.40

The worst 5 of 80 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	500	PLP	C4A-C4-C5	12.68	134.01	120.94
2	I	500	PLP	C3-C4-C5	-5.46	112.03	118.59
2	F	500	PLP	C3-C4-C5	-5.01	112.58	118.59
2	D	500	PLP	C6-C5-C4	4.83	122.06	118.10
2	I	500	PLP	C6-C5-C4	4.57	121.85	118.10

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	E	500	PLP	C4-C5-C5A-O4P
2	E	500	PLP	C6-C5-C5A-O4P
2	E	500	PLP	C5A-O4P-P-O1P
2	E	500	PLP	C5A-O4P-P-O2P
2	E	500	PLP	C5A-O4P-P-O3P

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	500	PLP	1	0
2	G	500	PLP	2	0
2	I	500	PLP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	442/456 (96%)	-0.58	0 100 100	9, 17, 35, 59	1 (0%)
1	B	442/456 (96%)	-0.56	1 (0%) 91 88	7, 16, 28, 47	0
1	C	442/456 (96%)	-0.61	0 100 100	9, 17, 32, 48	0
1	D	439/456 (96%)	-0.47	2 (0%) 87 82	9, 20, 35, 64	0
1	E	442/456 (96%)	-0.49	1 (0%) 91 88	10, 19, 33, 50	0
1	F	439/456 (96%)	-0.47	1 (0%) 91 88	9, 20, 35, 56	1 (0%)
1	G	440/456 (96%)	0.15	6 (1%) 73 64	15, 36, 54, 65	0
1	H	440/456 (96%)	0.12	5 (1%) 78 70	16, 36, 57, 69	0
1	I	441/456 (96%)	-0.20	0 100 100	14, 31, 51, 72	0
1	J	438/456 (96%)	-0.07	4 (0%) 81 74	14, 33, 50, 67	0
1	K	439/456 (96%)	0.25	12 (2%) 56 46	20, 39, 63, 87	0
1	L	436/456 (95%)	0.21	2 (0%) 87 82	20, 41, 62, 72	0
All	All	5280/5472 (96%)	-0.23	34 (0%) 85 80	7, 26, 53, 87	2 (0%)

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	367	ALA	5.2
1	K	366	ALA	4.2
1	J	366	ALA	3.5
1	J	93	ALA	3.3
1	D	366	ALA	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PLP	I	500	15/16	0.94	0.09	30,34,36,39	0
2	PLP	G	500	15/16	0.95	0.09	25,27,37,39	0
2	PLP	L	500	15/16	0.95	0.09	33,36,38,44	0
2	PLP	K	500	15/16	0.96	0.08	35,40,44,48	0
2	PLP	J	500	15/16	0.96	0.08	22,24,27,28	0
2	PLP	C	500	15/16	0.97	0.07	16,17,19,21	0
2	PLP	H	500	15/16	0.97	0.07	22,25,30,33	0
2	PLP	D	500	15/16	0.97	0.07	15,16,17,18	0
2	PLP	E	500	15/16	0.98	0.06	12,13,14,15	0
2	PLP	F	500	15/16	0.98	0.06	13,14,15,16	0
2	PLP	A	500	15/16	0.98	0.06	15,16,19,20	0
2	PLP	B	500	15/16	0.98	0.06	11,13,14,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.