



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:51 AM UTC

PDB ID : 7ATL / pdb_00007atl
Title : EstCE1, a hydrolase with promiscuous acyltransferase activity
Authors : Palm, G.J.; Lammers, M.; Berndt, L.
Deposited on : 2020-10-30
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7589 (2.50-2.46)
Clashscore	190562	8295 (2.50-2.46)
Ramachandran outliers	187476	8164 (2.50-2.46)
Sidechain outliers	187428	8166 (2.50-2.46)
RSRZ outliers	180081	7593 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	396	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6128 atoms, of which 2970 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Esterase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	381	5810	1850	2886	515	551	8	140	2	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	389	GLY	-	expression tag	UNP Q1I192
AAA	390	GLY	-	expression tag	UNP Q1I192
AAA	391	HIS	-	expression tag	UNP Q1I192
AAA	392	HIS	-	expression tag	UNP Q1I192
AAA	393	HIS	-	expression tag	UNP Q1I192
AAA	394	HIS	-	expression tag	UNP Q1I192
AAA	395	HIS	-	expression tag	UNP Q1I192
AAA	396	HIS	-	expression tag	UNP Q1I192

- Molecule 2 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	H	O	1	0
			17	4	10	3		
2	AAA	1	Total	C	H	O	1	0
			17	4	10	3		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	AAA	1	Total	C	H	O	1	0
			31	8	18	5		
3	AAA	1	Total	C	H	O	1	0
			31	8	18	5		

- Molecule 4 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	AAA	1	Total	C	H	O	1	0
			38	10	22	6		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	AAA	1	Total	C	H	O	1	0
			10	2	6	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	AAA	173	Total 174	O 174	0	1

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Esterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.43Å 99.43Å 228.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.62 – 2.48 48.62 – 2.48	Depositor EDS
% Data completeness (in resolution range)	99.0 (48.62-2.48) 98.0 (48.62-2.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.41	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.204 , 0.235 0.205 , 0.214	Depositor DCC
R_{free} test set	1209 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.125	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 40.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6128	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, EDO, PEG, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.05	2/2998 (0.1%)	1.36	3/4080 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	222	HIS	CE1-NE2	5.69	1.38	1.32
1	AAA	35	HIS	CE1-NE2	5.40	1.38	1.32

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	350	ASP	CB-CA-C	6.14	116.25	110.17
1	AAA	362	ASN	N-CA-C	-5.29	106.13	112.59
1	AAA	223	ASP	CA-CB-CG	5.25	117.85	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2924	2886	2876	17	0
2	AAA	14	20	20	0	0
3	AAA	26	36	36	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	AAA	16	22	22	0	0
5	AAA	4	6	5	1	0
6	AAA	174	0	0	0	0
All	All	3158	2970	2959	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:290:GLN:HA	1:AAA:290:GLN:HE21	1.51	0.75
1:AAA:222:HIS:HA	1:AAA:364:ALA:O	1.98	0.63
1:AAA:13:GLN:OE1	1:AAA:42:ASN:ND2	2.44	0.51
1:AAA:290:GLN:HA	1:AAA:290:GLN:NE2	2.24	0.51
1:AAA:290:GLN:HE21	1:AAA:290:GLN:CA	2.21	0.50
1:AAA:3:ILE:O	1:AAA:7:SER:HB2	2.12	0.50
1:AAA:33:ALA:HA	1:AAA:37:ARG:O	2.13	0.48
1:AAA:137:GLY:HA2	5:AAA:405:EDO:H12	1.96	0.46
1:AAA:200:GLN:HB2	1:AAA:201:PRO:HD3	2.00	0.43
1:AAA:224:THR:HG22	1:AAA:225:ALA:H	1.85	0.42
1:AAA:125:LEU:C	1:AAA:125:LEU:HD12	2.45	0.42
1:AAA:125:LEU:HD12	1:AAA:126:SER:N	2.35	0.42
1:AAA:24:ARG:O	1:AAA:363:THR:HG21	2.20	0.42
1:AAA:333:HIS:CG	1:AAA:352[B]:GLN:HG2	2.55	0.41
1:AAA:349:TYR:CD1	1:AAA:349:TYR:C	2.99	0.41
1:AAA:8:LEU:HD22	1:AAA:39:ILE:HD12	2.02	0.40
1:AAA:372:TYR:N	1:AAA:373:PRO:HD2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	381/396 (96%)	366 (96%)	13 (3%)	2 (0%)	24	40

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	225	ALA
1	AAA	344	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	300/311 (96%)	291 (97%)	9 (3%)	36	61

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	7	SER
1	AAA	52	LYS
1	AAA	123	TYR
1	AAA	143	GLU
1	AAA	176	ASP
1	AAA	224	THR
1	AAA	285	LYS
1	AAA	290	GLN
1	AAA	361	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PG4	AAA	403	-	12,12,12	1.10	2 (16%)	11,11,11	1.50	1 (9%)
5	EDO	AAA	405	-	3,3,3	0.26	0	2,2,2	0.36	0
4	1PE	AAA	404	-	15,15,15	0.42	0	14,14,14	0.28	0
2	PEG	AAA	406	-	6,6,6	0.30	0	5,5,5	0.15	0
2	PEG	AAA	401	-	6,6,6	0.40	0	5,5,5	0.29	0
3	PG4	AAA	402	-	12,12,12	0.58	0	11,11,11	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	AAA	403	-	-	6/10/10/10	-
5	EDO	AAA	405	-	-	0/1/1/1	-
4	1PE	AAA	404	-	-	7/13/13/13	-
2	PEG	AAA	406	-	-	2/4/4/4	-
2	PEG	AAA	401	-	-	3/4/4/4	-
3	PG4	AAA	402	-	-	8/10/10/10	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	403	PG4	O3-C5	2.33	1.52	1.42
3	AAA	403	PG4	O3-C4	2.30	1.52	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	AAA	403	PG4	C5-O3-C4	4.64	133.56	113.26

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	AAA	404	1PE	OH4-C13-C23-OH3
4	AAA	404	1PE	OH6-C15-C25-OH5
3	AAA	403	PG4	O2-C3-C4-O3
3	AAA	403	PG4	O3-C5-C6-O4
2	AAA	401	PEG	O1-C1-C2-O2
2	AAA	406	PEG	O2-C3-C4-O4
2	AAA	406	PEG	O1-C1-C2-O2
4	AAA	404	1PE	OH7-C16-C26-OH6
3	AAA	403	PG4	O1-C1-C2-O2
3	AAA	403	PG4	O4-C7-C8-O5
4	AAA	404	1PE	OH2-C12-C22-OH3
3	AAA	402	PG4	O2-C3-C4-O3
3	AAA	402	PG4	O3-C5-C6-O4
2	AAA	401	PEG	C1-C2-O2-C3
2	AAA	401	PEG	O2-C3-C4-O4
3	AAA	402	PG4	C4-C3-O2-C2
3	AAA	402	PG4	C5-C6-O4-C7
4	AAA	404	1PE	C14-C24-OH4-C13
3	AAA	402	PG4	C3-C4-O3-C5
3	AAA	402	PG4	C6-C5-O3-C4
4	AAA	404	1PE	C12-C22-OH3-C23
3	AAA	402	PG4	C1-C2-O2-C3
3	AAA	402	PG4	C8-C7-O4-C6
4	AAA	404	1PE	C15-C25-OH5-C14
3	AAA	403	PG4	C3-C4-O3-C5
3	AAA	403	PG4	C6-C5-O3-C4

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	AAA	405	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	381/396 (96%)	-0.52	6 (1%) 70 68	17, 30, 55, 99	2 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	225	ALA	7.0
1	AAA	224	THR	5.1
1	AAA	226	ALA	5.1
1	AAA	134[A]	HIS	3.2
1	AAA	3	ILE	2.8
1	AAA	286	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PEG	AAA	401	7/7	0.86	0.18	63,63,73,73	1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PEG	AAA	406	7/7	0.86	0.19	50,69,85,85	1
5	EDO	AAA	405	4/4	0.88	0.17	19,21,22,22	10
4	1PE	AAA	404	16/16	0.91	0.14	49,57,63,76	1
3	PG4	AAA	403	13/13	0.92	0.12	40,47,50,54	31
3	PG4	AAA	402	13/13	0.93	0.11	47,51,56,75	31

6.5 Other polymers [i](#)

There are no such residues in this entry.