



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:36 PM UTC

PDB ID : 7ATM / pdb_00007atm
Title : Structure of P. aeruginosa PBP3 in complex with a phenyl boronic acid (Compound 1)
Authors : Newman, H.; Bellini, B.; Dowson, C.G.
Deposited on : 2020-10-30
Resolution : 1.58 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.58 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	537	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	2101	-	-	X	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3926 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Peptidoglycan D,D-transpeptidase FtsI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	478	3669	2324	649	683	13	0	10	0

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	43	GLY	-	expression tag	UNP G3XD46
A	44	PRO	-	expression tag	UNP G3XD46
A	45	GLY	-	expression tag	UNP G3XD46
A	46	TYR	-	expression tag	UNP G3XD46
A	47	GLN	-	expression tag	UNP G3XD46
A	48	ASP	-	expression tag	UNP G3XD46
A	49	PRO	-	expression tag	UNP G3XD46

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is (3-(1H-tetrazol-5-yl)phenyl)boronic acid (CCD ID: RY2) (formula: $C_7H_7BN_4O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	B	C	N	0	0
			12	1	7	4		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	235	Total	O	0	0
			235	235		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Peptidoglycan D,D-transpeptidase FtsI



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	69.84Å 80.95Å 88.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.82 – 1.58 59.82 – 1.58	Depositor EDS
% Data completeness (in resolution range)	95.3 (59.82-1.58) 77.6 (59.82-1.58)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.77 (at 1.58Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.191 , 0.225 0.201 , 0.231	Depositor DCC
R_{free} test set	2622 reflections (2.44%)	wwPDB-VP
Wilson B-factor (Å ²)	26.1	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3926	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, RY2, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.07	2/3772 (0.1%)	1.26	5/5121 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	189	VAL	N-CA	5.57	1.50	1.45
1	A	76	SER	CA-CB	-5.37	1.45	1.53

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	532	TYR	CB-CA-C	5.52	118.88	111.82
1	A	135	GLU	CA-C-N	5.42	126.11	119.99
1	A	135	GLU	C-N-CA	5.42	126.11	119.99
1	A	62	ARG	CA-C-N	5.26	124.97	119.92
1	A	62	ARG	C-N-CA	5.26	124.97	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3669	0	3721	26	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	6	0	8	4	0
3	A	12	0	0	1	0
4	A	4	0	6	3	0
5	A	235	0	0	4	0
All	All	3926	0	3735	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:276:LEU:HD21	1:A:281:MET:SD	2.38	0.64
1:A:252[B]:MET:HE1	1:A:434:LEU:HD23	1.82	0.62
1:A:294:SER:CB	3:A:2102:RY2:C	2.81	0.59
1:A:82:LEU:HD13	1:A:137:VAL:HG11	1.84	0.58
1:A:259:ILE:HD12	1:A:436:MET:HG2	1.86	0.58
4:A:2103:DMS:C1	5:A:2250:HOH:O	2.51	0.58
1:A:333:VAL:HG23	1:A:532:TYR:CZ	2.40	0.56
1:A:395[B]:ARG:H	2:A:2101:GOL:HO1	1.56	0.54
1:A:395[C]:ARG:H	2:A:2101:GOL:HO1	1.56	0.54
1:A:270:PRO:O	1:A:273:ARG:NH1	2.42	0.52
1:A:505:SER:OG	1:A:533:PHE:C	2.52	0.52
1:A:86:PRO:O	1:A:90:MET:HG2	2.09	0.52
4:A:2103:DMS:H11	5:A:2250:HOH:O	2.10	0.52
1:A:266:PRO:HG2	1:A:280:ALA:HA	1.94	0.50
4:A:2103:DMS:H13	5:A:2250:HOH:O	2.12	0.48
1:A:479:TYR:CE2	1:A:557:PRO:HA	2.49	0.47
1:A:265:GLN:HG2	5:A:2273:HOH:O	2.15	0.46
1:A:84:ALA:HA	1:A:144:GLY:O	2.16	0.46
1:A:252[A]:MET:HE3	1:A:434:LEU:HD23	1.98	0.46
1:A:505:SER:OG	1:A:534:GLY:N	2.48	0.46
1:A:186:LEU:O	1:A:216:GLY:HA3	2.15	0.46
1:A:454:ALA:O	1:A:458:GLN:HG3	2.17	0.45
1:A:83:TRP:CE3	1:A:122:PHE:CD1	3.05	0.44
1:A:311:LYS:HE2	1:A:313:SER:OG	2.17	0.44
1:A:155:TYR:OH	1:A:167:PHE:HA	2.19	0.42
1:A:85:ASN:O	1:A:86:PRO:C	2.62	0.42
1:A:488:ALA:HB2	1:A:506:LEU:HD11	2.02	0.42
1:A:533:PHE:O	1:A:537:VAL:HG12	2.20	0.42
1:A:505:SER:OG	1:A:534:GLY:HA2	2.20	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	482/537 (90%)	470 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/424 (92%)	384 (99%)	4 (1%)	68	49

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	175	ARG
1	A	274	ARG
1	A	492	SER
1	A	529	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	275	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	390	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	GOL	A	2101	-	5,5,5	0.11	0	5,5,5	0.46	0
4	DMS	A	2103	-	3,3,3	0.19	0	3,3,3	0.10	0
3	RY2	A	2102	1	12,13,15	0.86	1 (8%)	15,17,20	1.90	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	2101	-	-	4/4/4/4	-
3	RY2	A	2102	1	-	2/4/4/8	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2102	RY2	N1-N	2.01	1.36	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2102	RY2	N-C5-N3	-4.43	107.72	112.10
3	A	2102	RY2	C5-N3-N2	4.05	109.27	105.65
3	A	2102	RY2	C4-C6-C	2.08	122.67	121.41
3	A	2102	RY2	C5-N-N1	2.00	106.31	102.32

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	2101	GOL	O1-C1-C2-C3
2	A	2101	GOL	C1-C2-C3-O3
2	A	2101	GOL	O1-C1-C2-O2
2	A	2101	GOL	O2-C2-C3-O3
3	A	2102	RY2	C6-C4-C5-N
3	A	2102	RY2	C3-C4-C5-N

There are no ring outliers.

3 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	2101	GOL	4	0
4	A	2103	DMS	3	0
3	A	2102	RY2	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	478/537 (89%)	0.87	90 (18%) 3 6	12, 29, 69, 113	10 (2%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	532	TYR	6.9
1	A	530	ALA	6.3
1	A	57	ALA	5.9
1	A	276	LEU	5.8
1	A	536	LEU	5.6
1	A	560	LEU	5.5
1	A	533	PHE	5.1
1	A	146	TYR	5.1
1	A	493	VAL	4.9
1	A	385	GLY	4.7
1	A	118	ALA	4.5
1	A	328	TYR	4.3
1	A	191	GLY	4.3
1	A	561	PRO	4.0
1	A	58	ILE	3.9
1	A	83	TRP	3.8
1	A	502	ALA	3.8
1	A	122	PHE	3.7
1	A	491	VAL	3.7
1	A	142	VAL	3.7
1	A	189	VAL	3.7
1	A	124	TYR	3.4
1	A	123	ILE	3.4
1	A	139	ALA	3.4
1	A	126	VAL	3.4
1	A	527	PRO	3.4
1	A	528	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	114	ILE	3.3
1	A	145	VAL	3.3
1	A	213	ALA	3.2
1	A	212	ASN	3.2
1	A	148	ILE	3.2
1	A	534	GLY	3.1
1	A	529	LYS	3.1
1	A	143	PRO	3.0
1	A	120	ARG	3.0
1	A	92	ALA	3.0
1	A	243	GLY	3.0
1	A	86	PRO	2.9
1	A	275	ASN	2.9
1	A	79	VAL	2.9
1	A	137	VAL	2.9
1	A	144	GLY	2.9
1	A	501	ASN	2.9
1	A	138	ILE	2.9
1	A	129	LEU	2.9
1	A	240	LEU	2.9
1	A	245	LYS	2.9
1	A	278	PRO	2.8
1	A	559	ASN	2.8
1	A	531	GLY	2.8
1	A	77	THR	2.7
1	A	279	ALA	2.7
1	A	89	LEU	2.7
1	A	103	LEU	2.7
1	A	140	LEU	2.6
1	A	384	PRO	2.6
1	A	91	THR	2.6
1	A	147	SER	2.6
1	A	273	ARG	2.6
1	A	277	GLN	2.5
1	A	78	PRO	2.5
1	A	469	GLY	2.5
1	A	109	LEU	2.5
1	A	214	LYS	2.5
1	A	82	LEU	2.4
1	A	119	GLU	2.4
1	A	60	ALA	2.4
1	A	101	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	281	MET	2.3
1	A	514	THR	2.3
1	A	324	GLN	2.3
1	A	121	GLU	2.3
1	A	128	GLY	2.3
1	A	141	LYS	2.3
1	A	87	LYS	2.2
1	A	445	GLY	2.2
1	A	173	ARG	2.2
1	A	535	GLY	2.1
1	A	131	PRO	2.1
1	A	272	ASN	2.1
1	A	151	PHE	2.1
1	A	170	VAL	2.1
1	A	537	VAL	2.1
1	A	85	ASN	2.1
1	A	116	GLN	2.1
1	A	492	SER	2.1
1	A	132	GLU	2.0
1	A	396	LYS	2.0
1	A	244	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	2101	6/6	0.79	0.19	46,54,61,65	0
4	DMS	A	2103	4/4	0.86	0.20	63,69,74,74	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	RY2	A	2102	12/14	0.97	0.05	18,21,27,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.