



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:00 PM UTC

PDB ID : 1AU8 / pdb_00001au8
Title : HUMAN CATHEPSIN G
Authors : Medrano, F.J.; Bode, W.; Banbula, A.; Potempa, J.
Deposited on : 1997-09-12
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	224	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CATHEPSIN G.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1786	1096	370	310	10			

- Molecule 2 is N-(3-carboxypropanoyl)-L-valyl-N-[(1R)-5-amino-1-phosphonopentyl]-L-prolinamide (CCD ID: 0H8) (formula: C₁₉H₃₅N₄O₈P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	19	4	7	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: CATHEPSIN G

Chain A:  73% 22% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.23Å 63.46Å 80.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 1.90	Depositor
% Data completeness (in resolution range)	97.1 (6.00-1.90)	Depositor
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.186 , 0.251	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1955	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OH8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/1820	1.29	30/2455 (1.2%)

There are no bond length outliers.

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	190	ALA	N-CA-C	-6.75	101.44	110.55
1	A	231	VAL	N-CA-C	6.74	118.30	110.62
1	A	200	LEU	O-C-N	6.62	130.77	123.22
1	A	198	PRO	CB-CA-C	-6.10	101.90	110.63
1	A	71	HIS	N-CA-C	-5.98	104.10	112.12
1	A	220	GLY	N-CA-C	-5.69	107.46	115.32
1	A	116	ARG	O-C-N	-5.63	115.78	122.20
1	A	116	ARG	NE-CZ-NH2	5.49	124.14	119.20
1	A	97	ARG	NE-CZ-NH2	5.42	124.08	119.20
1	A	148	ARG	NE-CZ-NH2	5.38	124.04	119.20
1	A	167	GLN	O-C-N	-5.34	116.32	122.09
1	A	41	ARG	NE-CZ-NH2	5.34	124.00	119.20
1	A	87	ARG	NE-CZ-NH2	5.32	123.99	119.20
1	A	111	ARG	NE-CZ-NH2	5.32	123.99	119.20
1	A	20	ARG	NE-CZ-NH2	5.31	123.98	119.20
1	A	230	ARG	NE-CZ-NH2	5.30	123.97	119.20
1	A	90	ARG	NE-CZ-NH2	5.28	123.95	119.20
1	A	131	ARG	NE-CZ-NH2	5.28	123.95	119.20
1	A	48	ARG	NE-CZ-NH2	5.26	123.94	119.20
1	A	110	ARG	NE-CZ-NH2	5.26	123.93	119.20
1	A	179	ARG	NE-CZ-NH2	5.24	123.92	119.20
1	A	164	ARG	NE-CZ-NH2	5.24	123.91	119.20
1	A	86	ARG	NE-CZ-NH2	5.22	123.90	119.20
1	A	147	ARG	NE-CZ-NH2	5.21	123.89	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	113	ARG	NE-CZ-NH2	5.19	123.87	119.20
1	A	35	ILE	N-CA-C	5.17	115.21	107.51
1	A	178	ARG	NE-CZ-NH2	5.15	123.83	119.20
1	A	114	ARG	NE-CZ-NH2	5.14	123.82	119.20
1	A	23	ARG	NE-CZ-NH2	5.07	123.77	119.20
1	A	143	ARG	NE-CZ-NH2	5.05	123.74	119.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1786	0	1797	31	0
2	A	31	0	31	1	0
3	A	138	0	0	6	0
All	All	1955	0	1828	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:ASN:ND2	3:A:347:HOH:O	1.76	1.19
1:A:41:ARG:HB3	3:A:314:HOH:O	1.55	1.06
1:A:41:ARG:CB	3:A:314:HOH:O	2.16	0.89
1:A:131:ARG:HD2	1:A:131:ARG:H	1.55	0.71
1:A:185:ARG:HA	1:A:224:PRO:HG2	1.72	0.70
1:A:238:ILE:HG22	1:A:242:MET:HE2	1.74	0.70
1:A:169:LEU:HD13	1:A:175:TYR:HD2	1.56	0.69
1:A:163:GLN:HE21	1:A:167:GLN:HE22	1.43	0.66
1:A:113:ARG:O	1:A:113:ARG:HG3	1.97	0.64
1:A:41:ARG:HG3	3:A:378:HOH:O	2.00	0.61
1:A:128:GLU:HG2	1:A:210:HIS:HE1	1.69	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:GLN:HE21	1:A:167:GLN:NE2	2.03	0.55
1:A:41:ARG:NH1	3:A:367:HOH:O	2.39	0.55
1:A:238:ILE:CG2	1:A:242:MET:HE2	2.38	0.53
1:A:146:MET:SD	1:A:191:PHE:HZ	2.34	0.50
1:A:127:GLN:O	1:A:127:GLN:HG3	2.11	0.50
1:A:81:GLN:NE2	1:A:113:ARG:HG3	2.27	0.49
1:A:87:ARG:NH2	1:A:89:ILE:HD11	2.27	0.49
1:A:29:TYR:CG	1:A:121:VAL:HB	2.49	0.48
1:A:121:VAL:HG21	1:A:200:LEU:HD21	1.96	0.46
1:A:131:ARG:HD2	1:A:131:ARG:N	2.26	0.46
1:A:169:LEU:HD13	1:A:175:TYR:CD2	2.43	0.46
1:A:236:PRO:HG3	1:A:239:ARG:HH21	1.81	0.44
1:A:96:GLN:HB3	3:A:315:HOH:O	2.18	0.43
1:A:144:VAL:CG2	1:A:148:ARG:HG3	2.49	0.43
1:A:135:LEU:HD21	1:A:161:ARG:NH1	2.35	0.42
1:A:35:ILE:O	1:A:39:GLN:HB3	2.21	0.41
1:A:16:ILE:O	1:A:144:VAL:HA	2.21	0.40
1:A:57:HIS:NE2	2:A:1:OH8:HB3	2.36	0.40
1:A:35:ILE:HG23	1:A:60:GLY:HA3	2.03	0.40
1:A:45:PHE:HB3	1:A:198:PRO:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	222/224 (99%)	213 (96%)	8 (4%)	1 (0%)	24 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	37	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	191/191 (100%)	186 (97%)	5 (3%)	40	35

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	A	113	ARG
1	A	128	GLU
1	A	169	LEU
1	A	235	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	39	GLN
1	A	74	GLN
1	A	159	GLN
1	A	167	GLN
1	A	210	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	0H8	A	1	1	27,31,32	8.36	2 (7%)	34,41,44	3.01	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0H8	A	1	1	-	5/33/48/50	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	0H8	CA2-N2	-43.16	1.06	1.45
2	A	1	0H8	O2-C1	3.87	1.43	1.30

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1	0H8	O-C-N1	-15.24	94.01	121.38
2	A	1	0H8	CA-C-N1	6.06	133.52	118.65
2	A	1	0H8	CD-N1-C	-3.40	113.80	125.85
2	A	1	0H8	CA1-N1-C	3.02	132.08	121.45

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1	0H8	O-C-N1-CA1
2	A	1	0H8	C2-C3-C4-N

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	1	0H8	O-C-N1-CD
2	A	1	0H8	C2-C3-C4-O4
2	A	1	0H8	O2-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1	0H8	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.