



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 04:14 AM EDT

PDB ID : 9AUS / pdb_00009aus
Title : Crystal structure of loop-closed dumbbell RNA bridged by glycine
Authors : Radakovic, A.; Lewicka, A.; Todisco, M.; Aitken, H.R.M.; Weiss, Z.; Kim, S.;
Bannan, A.; Piccirilli, J.A.; Szostak, J.W.
Deposited on : 2024-02-29
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)
RNA backbone	3983	1030 (2.40-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	225	12% 89% 11%
1	H	225	16% 93% 7%
2	B	215	8% 93% 7%
2	L	215	27% 87% 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	22	5% 64% 27% 9%
3	R	22	68% 23% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8017 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab BL3-6 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	225	Total	C	N	O	S	0	0	0
			1678	1055	287	330	6			
1	A	225	Total	C	N	O	S	0	0	0
			1678	1055	287	330	6			

- Molecule 2 is a protein called Fab BL3-6 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	215	Total	C	N	O	S	2	0	0
			1643	1025	275	337	6			
2	B	215	Total	C	N	O	S	2	0	0
			1643	1025	275	337	6			

- Molecule 3 is a RNA chain called Loop-closed dumbbell RNA bridged by glycine.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	R	22	Total	C	N	O	P	0	0	0
			476	213	93	148	22			
3	C	22	Total	C	N	O	P	0	0	0
			477	213	93	149	22			

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	L	1	Total	O	S	0	0
			5	4	1		
4	L	1	Total	O	S	0	0
			5	4	1		
4	R	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is GLYCINE (CCD ID: GLY) (formula: C₂H₅NO₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	R	1	Total	C	N	O	0	0
			4	2	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	H	108	Total	O	0	0
			108	108		
6	L	53	Total	O	0	0
			53	53		
6	R	30	Total	O	0	0
			30	30		
6	A	102	Total	O	0	0
			102	102		
6	B	76	Total	O	0	0
			76	76		
6	C	19	Total	O	0	0
			19	19		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab BL3-6 heavy chain



- Molecule 1: Fab BL3-6 heavy chain




- Molecule 2: Fab BL3-6 light chain




- Molecule 2: Fab BL3-6 light chain





● Molecule 3: Loop-closed dumbbell RNA bridged by glycine



● Molecule 3: Loop-closed dumbbell RNA bridged by glycine



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.51Å 82.08Å 92.67Å 63.95° 88.69° 88.40°	Depositor
Resolution (Å)	46.07 – 2.07 46.07 – 2.07	Depositor EDS
% Data completeness (in resolution range)	91.1 (46.07-2.07) 83.5 (46.07-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.88 (at 2.07Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.221 , 0.254 0.221 , 0.254	Depositor DCC
R_{free} test set	2034 reflections (3.58%)	wwPDB-VP
Wilson B-factor (Å ²)	35.3	Xtriage
Anisotropy	0.292	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 28.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.033 for h,-k,-l 0.020 for -h,k,k-l 0.008 for -h,-k,-k+l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8017	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.96% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.18	0/1719	0.30	0/2339
1	H	0.17	0/1719	0.30	0/2339
2	B	0.09	0/1678	0.27	0/2277
2	L	0.09	0/1678	0.26	0/2277
3	C	0.17	0/535	0.42	0/831
3	R	0.30	0/534	0.48	1/831 (0.1%)
All	All	0.16	0/7863	0.31	1/10894 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	R	1	G	OP1-P-OP2	-6.17	101.09	119.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1678	0	1638	13	0
1	H	1678	0	1637	10	0
2	B	1643	0	1595	8	0
2	L	1643	0	1594	19	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	477	0	240	5	0
3	R	476	0	239	2	0
4	B	10	0	0	0	0
4	C	5	0	0	0	0
4	L	10	0	0	1	0
4	R	5	0	0	0	0
5	R	4	0	2	0	0
6	A	102	0	0	1	0
6	B	76	0	0	1	0
6	C	19	0	0	1	0
6	H	108	0	0	1	0
6	L	53	0	0	2	0
6	R	30	0	0	0	0
All	All	8017	0	6945	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:122:SER:O	6:L:401:HOH:O	2.09	0.70
2:L:38:GLN:HB2	2:L:48:LEU:HD11	1.78	0.66
2:B:22:ILE:HD12	2:B:103:THR:HG21	1.78	0.65
1:H:4:GLU:N	6:H:304:HOH:O	2.29	0.64
2:B:3:ILE:HD13	2:B:30:VAL:HG12	1.79	0.64
1:A:4:GLU:N	6:A:302:HOH:O	2.31	0.62
2:B:38:GLN:HB2	2:B:48:LEU:HD11	1.81	0.62
1:H:217:THR:HG22	1:A:127:SER:HB2	1.82	0.61
2:L:91:GLN:HE21	2:L:98:THR:HG22	1.66	0.59
2:L:40:LYS:HD3	2:L:85:ALA:HB2	1.85	0.58
1:A:43:ALA:HB3	1:A:46:LYS:HD2	1.86	0.58
2:B:121:PRO:HD3	2:B:133:VAL:HG22	1.87	0.57
2:L:138:ASN:ND2	2:L:139:ASN:OD1	2.40	0.55
1:A:92:GLU:OE1	1:A:92:GLU:N	2.39	0.54
1:H:131:PRO:HB3	1:H:157:TYR:HB3	1.91	0.53
2:B:80:GLN:NE2	6:B:407:HOH:O	2.42	0.52
1:H:43:ALA:HB3	1:H:46:LYS:HD3	1.92	0.52
2:L:185:ALA:O	2:L:189:LYS:HG2	2.10	0.52
1:H:92:GLU:OE1	1:H:92:GLU:N	2.38	0.51
2:L:97:SER:OG	6:L:402:HOH:O	2.20	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:L:146:LYS:HB3	2:L:198:THR:HB	1.93	0.51
3:C:17:G:H1'	3:C:21:A:H61	1.76	0.50
2:B:66:SER:OG	2:B:73:THR:OG1	2.30	0.49
2:L:49:ILE:HD13	2:L:55:LEU:HA	1.94	0.49
3:C:20:A:H1'	3:C:21:A:H5''	1.95	0.49
1:H:16:GLN:HG3	1:H:17:PRO:HD2	1.95	0.48
3:R:20:A:H4'	3:R:21:A:OP1	2.11	0.48
2:L:125:GLN:O	2:L:128:SER:OG	2.30	0.46
3:C:19:G:C6	3:C:21:A:C6	3.03	0.46
1:A:227:SER:OG	1:A:228:CYS:N	2.47	0.46
1:A:131:PRO:HB3	1:A:157:TYR:HB3	1.98	0.45
1:A:17:PRO:HD3	1:A:124:SER:C	2.42	0.45
1:H:212:HIS:HB3	1:H:217:THR:OG1	2.17	0.45
2:L:211:ASN:HB2	2:L:214:GLU:HB3	1.99	0.45
2:L:36:TRP:CZ3	2:L:89:CYS:HB3	2.53	0.44
1:H:130:GLY:HA3	1:H:217:THR:HG21	1.99	0.44
2:L:67:ARG:NH2	4:L:301:SO4:O3	2.41	0.43
2:B:211:ASN:O	2:B:214:GLU:HG2	2.19	0.43
2:L:211:ASN:H	2:L:214:GLU:CD	2.26	0.43
1:A:86:MET:HB3	1:A:89:LEU:HD21	2.01	0.43
2:B:108:LYS:HA	2:B:141:TYR:OH	2.19	0.43
1:H:86:MET:HB3	1:H:89:LEU:HD21	2.01	0.42
1:A:106:ARG:HB2	6:C:202:HOH:O	2.19	0.42
2:L:39:GLN:O	2:L:85:ALA:HB1	2.19	0.42
1:A:138:PRO:HG2	1:A:225:PRO:HG3	2.01	0.42
2:L:126:LEU:HD12	2:L:184:LYS:HG3	2.01	0.42
3:C:20:A:N3	3:C:21:A:H2'	2.34	0.42
3:R:5:G:O6	3:R:7:A:O2'	2.37	0.42
2:L:189:LYS:HA	2:L:189:LYS:HD3	1.90	0.41
2:L:214:GLU:O	2:L:215:CYS:HB2	2.21	0.41
1:H:15:VAL:HG11	1:H:89:LEU:HD13	2.02	0.41
1:A:213:LYS:HE2	1:A:213:LYS:HB3	1.83	0.41
3:C:17:G:H1'	3:C:21:A:N6	2.36	0.41
1:A:180:ALA:HA	1:A:190:LEU:HB3	2.03	0.40
2:L:36:TRP:CD2	2:L:74:LEU:HB2	2.56	0.40
1:A:94:THR:HG23	1:A:122:THR:HA	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/225 (99%)	216 (97%)	7 (3%)	0	100	100
1	H	223/225 (99%)	218 (98%)	5 (2%)	0	100	100
2	B	213/215 (99%)	207 (97%)	6 (3%)	0	100	100
2	L	213/215 (99%)	207 (97%)	6 (3%)	0	100	100
All	All	872/880 (99%)	848 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/186 (100%)	183 (98%)	3 (2%)	55	55
1	H	186/186 (100%)	184 (99%)	2 (1%)	65	68
2	B	190/190 (100%)	189 (100%)	1 (0%)	81	85
2	L	190/190 (100%)	189 (100%)	1 (0%)	81	85
All	All	752/752 (100%)	745 (99%)	7 (1%)	70	74

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	90	ARG
1	H	228	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	189	LYS
1	A	28	SER
1	A	125	SER
1	A	147	THR
2	B	144	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	42	GLN
1	H	183	GLN
2	L	39	GLN
1	A	85	GLN
1	A	87	ASN
2	B	80	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	C	21/22 (95%)	6 (28%)	1 (4%)
3	R	21/22 (95%)	4 (19%)	1 (4%)
All	All	42/44 (95%)	10 (23%)	2 (4%)

All (10) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	R	4	U
3	R	20	A
3	R	21	A
3	R	22	A
3	C	2	G
3	C	3	U
3	C	4	U
3	C	20	A
3	C	21	A
3	C	22	A

All (2) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
3	R	20	A
3	C	20	A

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GLY	R	102	3	3,3,4	0.61	0	1,2,4	0.24	0
4	SO4	B	301	-	4,4,4	0.24	0	6,6,6	0.08	0
4	SO4	B	302	-	4,4,4	0.24	0	6,6,6	0.08	0
4	SO4	R	101	-	4,4,4	0.23	0	6,6,6	0.08	0
4	SO4	L	301	-	4,4,4	0.23	0	6,6,6	0.09	0
4	SO4	C	101	-	4,4,4	0.24	0	6,6,6	0.07	0
4	SO4	L	302	-	4,4,4	0.23	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GLY	R	102	3	-	0/0/1/2	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	L	301	SO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/225 (100%)	0.90	26 (11%) 9 9	27, 40, 73, 116	1 (0%)
1	H	225/225 (100%)	1.06	37 (16%) 4 4	28, 40, 90, 132	1 (0%)
2	B	215/215 (100%)	1.05	18 (8%) 17 17	31, 46, 64, 114	2 (0%)
2	L	215/215 (100%)	1.38	58 (26%) 1 1	31, 52, 97, 128	2 (0%)
3	C	22/22 (100%)	0.53	1 (4%) 38 38	41, 91, 127, 135	0
3	R	22/22 (100%)	0.09	0 100 100	35, 47, 58, 67	0
All	All	924/924 (100%)	1.06	140 (15%) 5 5	27, 45, 92, 135	6 (0%)

All (140) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	185	ALA	5.2
1	H	202	GLY	4.9
2	L	192	VAL	4.9
2	B	1	SER	4.9
1	H	206	TYR	4.8
1	H	143	THR	4.7
1	H	200	SER	4.5
1	H	201	LEU	4.4
1	H	140	SER	4.3
2	L	190	HIS	4.3
2	L	212	ARG	4.3
1	A	145	GLY	4.2
2	L	154	ALA	4.2
2	B	24	CYS	4.0
2	L	210	PHE	3.9
1	H	150	LEU	3.9
1	H	203	THR	3.8
1	H	205	THR	3.5
2	L	188	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	H	137	ALA	3.4
2	L	215	CYS	3.3
2	L	193	TYR	3.3
1	H	223	VAL	3.3
1	H	227	SER	3.3
2	L	151	VAL	3.3
1	H	228	CYS	3.3
2	B	3	ILE	3.3
2	B	195	CYS	3.2
2	L	126	LEU	3.2
2	L	152	ASP	3.1
1	A	146	GLY	3.1
2	B	2	ASP	3.1
1	A	143	THR	3.0
2	L	189	LYS	3.0
2	L	191	LYS	3.0
1	H	152	CYS	3.0
2	L	131	ALA	3.0
2	L	2	ASP	3.0
1	A	202	GLY	3.0
2	L	184	LYS	3.0
1	H	207	ILE	3.0
1	H	225	PRO	3.0
2	B	215	CYS	2.9
1	H	56	PRO	2.9
2	L	211	ASN	2.9
1	A	170	ALA	2.9
1	A	201	LEU	2.9
2	L	107	ILE	2.9
2	B	214	GLU	2.9
2	L	104	LYS	2.9
2	L	187	TYR	2.9
2	L	213	GLY	2.9
1	A	15	VAL	2.8
2	L	194	ALA	2.8
2	L	12	LEU	2.8
2	L	98	THR	2.8
1	A	144	SER	2.7
1	H	224	GLU	2.7
2	L	84	PHE	2.7
2	B	79	LEU	2.7
1	H	4	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	142	SER	2.6
2	L	149	TRP	2.6
2	L	207	THR	2.6
1	A	106	ARG	2.6
1	A	228	CYS	2.6
2	L	21	THR	2.6
1	H	138	PRO	2.6
2	L	182	LEU	2.6
2	L	206	VAL	2.5
2	L	1	SER	2.5
2	L	78	SER	2.5
2	L	156	GLN	2.5
2	L	158	GLY	2.5
2	B	203	SER	2.5
2	L	180	LEU	2.5
2	B	127	LYS	2.5
1	H	196	VAL	2.5
2	B	123	ASP	2.5
1	H	139	SER	2.5
1	H	199	SER	2.5
2	L	130	THR	2.5
2	L	155	LEU	2.5
2	L	150	LYS	2.5
1	H	145	GLY	2.5
2	L	214	GLU	2.4
2	L	24	CYS	2.4
1	A	139	SER	2.4
2	L	147	VAL	2.4
1	A	227	SER	2.4
2	L	132	SER	2.4
2	L	157	SER	2.4
1	A	147	THR	2.4
1	H	170	ALA	2.4
2	L	20	VAL	2.4
2	L	197	VAL	2.4
1	H	226	LYS	2.4
1	A	141	LYS	2.3
2	B	213	GLY	2.3
1	H	127	SER	2.3
2	B	169	SER	2.3
2	L	127	LYS	2.3
1	H	204	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	107	ILE	2.2
1	A	56	PRO	2.2
2	B	207	THR	2.2
2	B	150	LYS	2.2
1	A	162	VAL	2.2
1	A	225	PRO	2.2
1	H	25	CYS	2.2
2	L	195	CYS	2.2
1	H	172	THR	2.2
1	H	146	GLY	2.2
1	A	211	ASN	2.2
1	H	141	LYS	2.1
1	H	208	CYS	2.1
1	A	140	SER	2.1
1	H	197	PRO	2.1
1	A	44	PRO	2.1
2	B	27	SER	2.1
2	L	161	GLN	2.1
2	L	205	PRO	2.1
1	A	226	LYS	2.1
2	L	118	ILE	2.1
2	L	30	VAL	2.1
2	L	204	SER	2.1
1	H	136	LEU	2.0
3	C	22	A	2.0
1	A	99	CYS	2.0
1	A	205	THR	2.0
2	L	120	PRO	2.0
1	A	207	ILE	2.0
1	A	4	GLU	2.0
2	L	133	VAL	2.0
2	L	15	SER	2.0
2	L	209	SER	2.0
1	H	91	ALA	2.0
1	H	182	LEU	2.0
2	B	126	LEU	2.0
2	L	146	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	C	101	5/5	0.58	0.13	130,131,136,137	0
4	SO4	L	302	5/5	0.65	0.12	94,95,101,125	0
4	SO4	R	101	5/5	0.72	0.13	60,64,72,75	0
4	SO4	B	301	5/5	0.73	0.10	83,84,100,116	0
4	SO4	B	302	5/5	0.78	0.09	67,71,88,88	0
5	GLY	R	102	4/5	0.78	0.12	63,64,65,66	0
4	SO4	L	301	5/5	0.84	0.08	66,67,69,71	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.