



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:19 PM UTC

PDB ID : 1AVB / pdb_00001avb
Title : ARCELIN-1 FROM PHASEOLUS VULGARIS L
Authors : Mourey, L.; Pedelacq, J.D.; Fabre, C.; Rouge, P.; Samama, J.P.
Deposited on : 1997-09-15
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	226	10% 87% 11% .
1	B	226	10% 84% 15% .
2	C	2	100%
2	D	2	100%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3868 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ARCELIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1756	1095	302	356	3			
1	B	226	Total	C	N	O	S	0	0	0
			1760	1098	303	356	3			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	D	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	113	Total	O	0	0
			113	113		
5	B	117	Total	O	0	0
			117	117		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ARCELIN-1




- Molecule 1: ARCELIN-1



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.65Å 92.57Å 67.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	33.71 – 1.90 33.71 – 1.90	Depositor EDS
% Data completeness (in resolution range)	94.9 (33.71-1.90) 97.0 (33.71-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.96 (at 1.89Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.208 , 0.242 0.219 , 0.250	Depositor DCC
R_{free} test set	2130 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	14.2	Xtriage
Anisotropy	0.907	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 64.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.60$, $\langle L^2 \rangle = 0.46$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3868	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9765e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1790	0.92	4/2432 (0.2%)
1	B	0.44	0/1794	0.92	3/2436 (0.1%)
All	All	0.45	0/3584	0.92	7/4868 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	87	PHE	N-CA-C	-6.96	97.90	109.24
1	A	129	ILE	N-CA-C	-6.92	98.47	108.36
1	B	87	PHE	N-CA-C	-6.72	98.73	109.76
1	B	129	ILE	N-CA-C	-6.23	99.45	108.36
1	B	64	SER	N-CA-C	-5.47	101.66	110.14
1	A	64	SER	N-CA-C	-5.11	101.82	109.95
1	A	49	SER	N-CA-C	5.06	118.55	112.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1756	0	1682	21	0
1	B	1760	0	1693	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	A	28	0	26	0	0
3	B	28	0	26	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	113	0	0	1	0
5	B	117	0	0	1	0
All	All	3868	0	3477	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:ALA:H	1:A:150:ASN:HD21	1.08	0.97
1:A:168:ARG:HH11	1:A:168:ARG:HG3	1.31	0.94
1:B:75:ALA:H	1:B:150:ASN:HD21	1.16	0.89
1:B:168:ARG:HH11	1:B:168:ARG:HG3	1.48	0.76
1:B:75:ALA:H	1:B:150:ASN:ND2	1.93	0.62
1:A:2:ASN:HB3	5:A:578:HOH:O	2.00	0.60
1:A:89:LEU:HD22	1:A:198:VAL:HG12	1.84	0.59
1:B:62:LEU:HD23	1:B:162:SER:HB2	1.85	0.58
1:A:80:ASN:O	1:A:210:THR:HG22	2.04	0.58
1:A:168:ARG:HG3	1:A:168:ARG:NH1	2.07	0.57
1:A:54:ILE:O	1:A:62:LEU:HD23	2.08	0.54
1:B:89:LEU:HD22	1:B:198:VAL:HG12	1.91	0.53
1:B:80:ASN:O	1:B:210:THR:HG22	2.10	0.52
1:A:75:ALA:H	1:A:150:ASN:ND2	1.91	0.51
1:B:2:ASN:HB3	5:B:585:HOH:O	2.11	0.51
1:B:59:ILE:O	1:B:59:ILE:HG22	2.12	0.49
1:A:137:ARG:HG3	1:A:137:ARG:HH11	1.76	0.49
1:B:7:ASN:HD22	1:B:220:SER:HB3	1.78	0.49
1:B:31:LEU:HD13	1:B:35:VAL:HG11	1.95	0.48
1:B:15:ASN:H	1:B:15:ASN:HD22	1.60	0.48
1:B:61:ASN:HD22	1:B:62:LEU:H	1.60	0.48
1:A:15:ASN:HD22	1:A:15:ASN:H	1.62	0.47
1:B:59:ILE:HG22	1:B:61:ASN:HB2	1.96	0.47
1:B:75:ALA:N	1:B:150:ASN:HD21	1.97	0.46
1:B:57:ARG:HG3	1:B:57:ARG:HH11	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:59:ILE:HG22	1:A:61:ASN:HB2	1.99	0.45
1:A:62:LEU:HD23	1:A:63:ALA:H	1.80	0.45
1:B:62:LEU:HD13	1:B:189:GLU:HB3	1.97	0.45
1:A:111:TYR:HA	1:A:135:SER:HB2	1.98	0.45
1:A:188:LEU:O	1:A:192:VAL:HG22	2.17	0.44
1:B:161:ASP:OD1	1:B:163:PRO:HD2	2.19	0.43
1:B:89:LEU:HD13	1:B:196:VAL:HG21	2.00	0.43
1:A:62:LEU:HD22	1:A:63:ALA:O	2.19	0.43
1:A:59:ILE:HG22	1:A:59:ILE:O	2.18	0.43
1:A:61:ASN:OD1	1:A:62:LEU:N	2.52	0.43
1:A:62:LEU:HD23	1:A:63:ALA:N	2.35	0.42
1:B:168:ARG:HG3	1:B:168:ARG:NH1	2.22	0.42
1:B:113:ARG:H	1:B:113:ARG:HG2	1.66	0.42
1:A:55:ASN:HD22	1:A:56:ASP:N	2.18	0.42
1:B:136:ILE:O	1:B:138:PRO:HD3	2.19	0.42
1:A:95:ARG:HG2	1:A:95:ARG:HH11	1.86	0.41
1:B:61:ASN:HD22	1:B:62:LEU:N	2.19	0.41
1:A:8:VAL:HG22	1:B:2:ASN:OD1	2.21	0.40
1:B:17:ILE:O	1:B:46:ALA:HA	2.22	0.40
1:B:171:LEU:HD23	1:B:171:LEU:C	2.47	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	218 (97%)	6 (3%)	0	100	100
1	B	224/226 (99%)	217 (97%)	7 (3%)	0	100	100
All	All	448/452 (99%)	435 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/199 (100%)	190 (96%)	8 (4%)	28	20
1	B	199/199 (100%)	192 (96%)	7 (4%)	32	24
All	All	397/398 (100%)	382 (96%)	15 (4%)	29	21

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	41	ASP
1	A	49	SER
1	A	62	LEU
1	A	79	GLU
1	A	80	ASN
1	A	168	ARG
1	A	207	LYS
1	B	9	GLU
1	B	15	ASN
1	B	89	LEU
1	B	135	SER
1	B	178	GLU
1	B	209	GLU
1	B	224	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	ASN
1	A	15	ASN
1	A	38	ASN
1	A	55	ASN
1	A	127	ASN
1	A	148	HIS
1	A	150	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	165	ASN
1	A	224	ASN
1	B	7	ASN
1	B	15	ASN
1	B	61	ASN
1	B	80	ASN
1	B	150	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.46	0	17,19,21	0.80	0
2	NAG	C	2	2	14,14,15	0.42	0	17,19,21	0.73	0
2	NAG	D	1	1,2	14,14,15	0.42	0	17,19,21	0.86	0
2	NAG	D	2	2	14,14,15	0.41	0	17,19,21	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C1-C2-N2-C7
2	C	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	A	500	1	14,14,15	0.47	0	17,19,21	1.08	2 (11%)
4	SO4	A	501	-	4,4,4	0.76	0	6,6,6	0.66	0
3	NAG	B	500	1	14,14,15	0.43	0	17,19,21	0.95	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	B	400	1	14,14,15	0.41	0	17,19,21	0.70	0
4	SO4	B	501	-	4,4,4	0.76	0	6,6,6	0.35	0
3	NAG	A	400	1	14,14,15	0.41	0	17,19,21	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	500	1	-	2/6/23/26	0/1/1/1
3	NAG	B	400	1	-	2/6/23/26	0/1/1/1
3	NAG	A	500	1	-	1/6/23/26	0/1/1/1
3	NAG	A	400	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	NAG	C1-O5-C5	2.59	115.66	112.19
3	A	500	NAG	C2-N2-C7	-2.52	119.52	122.90
3	B	500	NAG	C2-N2-C7	-2.32	119.80	122.90

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	400	NAG	O5-C5-C6-O6
3	A	400	NAG	O5-C5-C6-O6
3	B	400	NAG	C4-C5-C6-O6
3	A	400	NAG	C4-C5-C6-O6
3	A	500	NAG	O5-C5-C6-O6
3	B	500	NAG	C4-C5-C6-O6
3	B	500	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/226 (100%)	0.47	22 (9%) 13 14	8, 16, 45, 72	0
1	B	226/226 (100%)	0.54	23 (10%) 12 12	9, 16, 43, 67	0
All	All	452/452 (100%)	0.50	45 (9%) 12 13	8, 16, 45, 72	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	59	ILE	7.3
1	B	59	ILE	7.1
1	B	58	THR	7.1
1	A	61	ASN	5.8
1	B	62	LEU	5.0
1	B	226	ILE	4.7
1	A	62	LEU	4.6
1	B	78	ILE	4.3
1	B	61	ASN	4.3
1	A	58	THR	4.2
1	B	60	ASP	4.1
1	B	57	ARG	3.9
1	A	77	ASN	3.9
1	A	78	ILE	3.9
1	B	77	ASN	3.9
1	A	60	ASP	3.9
1	A	56	ASP	3.8
1	B	168	ARG	3.4
1	A	80	ASN	3.3
1	A	57	ARG	3.3
1	A	226	ILE	3.2
1	B	113	ARG	3.0
1	A	206	SER	3.0
1	A	49	SER	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	76	LYS	2.9
1	A	113	ARG	2.7
1	B	209	GLU	2.7
1	A	207	LYS	2.6
1	A	224	ASN	2.6
1	B	208	LYS	2.5
1	A	148	HIS	2.5
1	B	181	HIS	2.5
1	B	56	ASP	2.5
1	A	137	ARG	2.4
1	B	206	SER	2.4
1	B	207	LYS	2.4
1	A	109	THR	2.3
1	B	108	THR	2.3
1	A	168	ARG	2.2
1	B	109	THR	2.2
1	B	224	ASN	2.1
1	A	76	LYS	2.1
1	B	79	GLU	2.1
1	B	80	ASN	2.1
1	A	36	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.80	0.16	31,38,46,48	0
2	NAG	D	2	14/15	0.82	0.14	30,37,43,45	0
2	NAG	D	1	14/15	0.91	0.09	14,20,24,25	0
2	NAG	C	1	14/15	0.93	0.08	13,19,24,25	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	500	14/15	0.55	0.20	64,72,75,77	0
3	NAG	A	500	14/15	0.65	0.19	54,64,71,72	0
3	NAG	A	400	14/15	0.86	0.12	24,32,41,43	0
3	NAG	B	400	14/15	0.88	0.11	25,34,43,45	0
4	SO4	B	501	5/5	0.98	0.05	18,19,20,21	0
4	SO4	A	501	5/5	0.99	0.04	16,18,20,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.