



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:55 AM UTC

PDB ID : 8AV2 / pdb_00008av2
Title : Crystal structure for the FnIII module of mouse LEP-R in complex with the anti-LEP-R nanobody VHH-4.80
Authors : Verstraete, K.; Savvides, S.N.; Verschueren, K.G.; Tsirigotaki, A.
Deposited on : 2022-08-26
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	198	12% 88% 8% ••
1	B	198	11% 90% 9% •
2	C	141	2% 85% 6% 9%
2	D	141	5% 85% 5% • 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Leptin receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	192	Total	C	N	O	S	0	7	0
			1622	1052	272	291	7			
1	B	196	Total	C	N	O	S	0	6	0
			1637	1059	273	298	7			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	630	GLU	-	expression tag	UNP P48356
A	631	THR	-	expression tag	UNP P48356
A	632	GLY	-	expression tag	UNP P48356
A	668	GLN	ASN	engineered mutation	UNP P48356
A	672	SER	CYS	conflict	UNP P48356
A	698	GLN	ASN	engineered mutation	UNP P48356
A	726	GLN	ASN	engineered mutation	UNP P48356
B	630	GLU	-	expression tag	UNP P48356
B	631	THR	-	expression tag	UNP P48356
B	632	GLY	-	expression tag	UNP P48356
B	668	GLN	ASN	engineered mutation	UNP P48356
B	672	SER	CYS	conflict	UNP P48356
B	698	GLN	ASN	engineered mutation	UNP P48356
B	726	GLN	ASN	engineered mutation	UNP P48356

- Molecule 2 is a protein called anti-mLEPR VHH 4-80.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	129	Total	C	N	O	S	0	5	0
			1034	644	175	207	8			
2	D	128	Total	C	N	O	S	0	0	0
			984	614	164	199	7			


- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula:

C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	C	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0
4	D	1	Total O S 5 4 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	164	Total O 164 164	0	0
5	B	125	Total O 125 125	0	0
5	C	142	Total O 142 142	0	0
5	D	83	Total O 83 83	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Leptin receptor




- Molecule 1: Leptin receptor



- Molecule 2: anti-mLEPR VHH 4-80



- Molecule 2: anti-mLEPR VHH 4-80



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	53.19Å 85.88Å 144.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.42 – 1.75 25.42 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.42-1.75) 99.9 (25.42-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.01 (at 1.75Å)	Xtriage
Refinement program	BUSTER 2.10.4 (3-FEB-2022)	Depositor
R, R_{free}	0.193 , 0.233 0.191 , 0.228	Depositor DCC
R_{free} test set	3374 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.168	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5897	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CSO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/1661	0.95	1/2258 (0.0%)
1	B	0.65	0/1675	0.86	0/2276
2	C	0.73	0/1055	0.83	0/1426
2	D	0.66	1/1005 (0.1%)	0.84	1/1361 (0.1%)
All	All	0.68	1/5396 (0.0%)	0.88	2/7321 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	86	MET	SD-CE	-5.63	1.65	1.79

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	116	ASP	CA-CB-CG	5.80	118.41	112.60
1	A	761	ASP	N-CA-C	5.54	117.48	110.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1622	0	1617	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1637	0	1627	8	0
2	C	1034	0	979	6	0
2	D	984	0	927	5	0
3	A	28	0	26	0	0
3	B	28	0	26	0	0
4	A	25	0	0	0	0
4	B	10	0	0	0	0
4	C	5	0	0	0	0
4	D	10	0	0	1	0
5	A	164	0	0	0	0
5	B	125	0	0	0	0
5	C	142	0	0	0	1
5	D	83	0	0	0	0
All	All	5897	0	5202	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:37:ILE:HB	2:C:82[B]:VAL:HG21	1.85	0.58
1:B:804:LYS:HE2	1:B:824:ASN:HB3	1.88	0.54
1:B:738:ALA:HB2	1:B:813:PHE:HZ	1.74	0.53
2:C:85:GLN:HE21	2:C:87:ASN:HD21	1.58	0.52
1:B:644:TRP:CG	1:B:645:ARG:H	2.28	0.52
1:A:754:LEU:HD21	1:A:770:ILE:HD13	1.93	0.51
1:A:645[B]:ARG:HH21	1:A:729:PHE:HE1	1.57	0.51
1:B:638[A]:MET:SD	1:B:716:ASN:ND2	2.87	0.48
1:A:644:TRP:CG	1:A:645[A]:ARG:H	2.30	0.48
1:B:743[B]:SER:OG	1:B:755[B]:SER:OG	2.32	0.48
1:A:814:MET:SD	1:B:725:PHE:CE1	3.08	0.47
1:A:644:TRP:CG	1:A:645[B]:ARG:H	2.33	0.46
2:D:23:LEU:HG	2:D:86:MET:HE2	1.97	0.45
1:A:636[A]:VAL:HG22	1:A:718:LEU:HB3	1.99	0.45
2:C:86[A]:MET:HE2	2:C:89:LEU:HD21	2.00	0.43
2:C:5:VAL:HG13	2:C:30:PHE:CD1	2.54	0.43
2:C:86[B]:MET:HE1	2:C:128:VAL:HG21	2.01	0.43
2:D:110:ARG:NH2	4:D:201:SO4:O1	2.45	0.43
1:A:814:MET:HE2	1:A:814:MET:HB3	1.93	0.42
1:B:776:ASN:OD1	2:D:58:ASP:HB2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:7:LEU:HD21	2:D:30:PHE:HZ	1.85	0.42
2:D:21:LEU:HB2	2:D:86:MET:HE3	2.02	0.42
2:C:85:GLN:HE21	2:C:87:ASN:ND2	2.18	0.41
1:B:662:TRP:O	1:B:697:THR:HB	2.20	0.41
1:A:802:ILE:HD12	1:A:802:ILE:HA	1.97	0.40
1:A:745:TYR:HA	1:A:746:PRO:HD3	1.94	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:318:HOH:O	5:C:418:HOH:O[4_555]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/198 (98%)	185 (95%)	10 (5%)	0	100	100
1	B	199/198 (100%)	189 (95%)	10 (5%)	0	100	100
2	C	132/141 (94%)	130 (98%)	2 (2%)	0	100	100
2	D	126/141 (89%)	125 (99%)	1 (1%)	0	100	100
All	All	652/678 (96%)	629 (96%)	23 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	183/181 (101%)	175 (96%)	8 (4%)	25	7
1	B	185/181 (102%)	183 (99%)	2 (1%)	65	52
2	C	111/116 (96%)	111 (100%)	0	100	100
2	D	105/116 (90%)	105 (100%)	0	100	100
All	All	584/594 (98%)	574 (98%)	10 (2%)	55	36

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	638[A]	MET
1	A	638[B]	MET
1	A	656	ARG
1	A	669	ASP
1	A	671	LEU
1	A	718	LEU
1	A	754	LEU
1	A	773	LYS
1	B	656	ARG
1	B	699	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	C	6	GLN
2	C	87	ASN
2	D	8	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	751	1	3,6,7	0.59	0	1,6,8	0.04	0
1	CSO	B	751	1	3,6,7	0.81	0	1,6,8	2.46	1 (100%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	751	1	-	0/1/5/7	-
1	CSO	B	751	1	-	0/1/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	751	CSO	CB-CA-C	2.46	117.47	110.80

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	901	1	14,14,15	0.33	0	17,19,21	0.78	1 (5%)
3	NAG	A	902	1	14,14,15	0.26	0	17,19,21	0.38	0
4	SO4	A	904	-	4,4,4	0.33	0	6,6,6	0.20	0
4	SO4	D	201	-	4,4,4	0.32	0	6,6,6	0.33	0
4	SO4	B	903	-	4,4,4	0.30	0	6,6,6	0.25	0
3	NAG	B	902	1	14,14,15	0.32	0	17,19,21	0.54	0
3	NAG	A	901	1	14,14,15	0.29	0	17,19,21	0.71	1 (5%)
4	SO4	B	904	-	4,4,4	0.34	0	6,6,6	0.40	0
4	SO4	A	903	-	4,4,4	0.41	0	6,6,6	0.66	0
4	SO4	C	201	-	4,4,4	0.31	0	6,6,6	1.44	1 (16%)
4	SO4	A	905	-	4,4,4	0.28	0	6,6,6	0.43	0
4	SO4	A	906	-	4,4,4	0.34	0	6,6,6	0.12	0
4	SO4	D	202	-	4,4,4	0.30	0	6,6,6	0.25	0
4	SO4	A	907	-	4,4,4	0.35	0	6,6,6	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	901	1	-	0/6/23/26	0/1/1/1
3	NAG	A	902	1	-	0/6/23/26	0/1/1/1
3	NAG	B	902	1	-	0/6/23/26	0/1/1/1
3	NAG	A	901	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	201	SO4	O4-S-O2	-3.17	93.00	109.56
3	B	901	NAG	C1-O5-C5	2.62	115.70	112.19
3	A	901	NAG	C1-O5-C5	2.09	114.98	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	201	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	191/198 (96%)	0.31	24 (12%) 8 9	11, 31, 64, 91	7 (3%)
1	B	195/198 (98%)	0.49	21 (10%) 11 12	15, 37, 68, 88	6 (3%)
2	C	129/141 (91%)	-0.13	3 (2%) 61 67	11, 26, 49, 64	5 (3%)
2	D	128/141 (90%)	0.37	7 (5%) 30 35	25, 40, 63, 68	0
All	All	643/678 (94%)	0.29	55 (8%) 16 19	11, 34, 64, 91	18 (2%)

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	636[A]	VAL	6.4
1	A	671	LEU	5.6
1	A	666	THR	4.4
2	D	30	PHE	4.3
1	B	718	LEU	4.1
2	D	44	PRO	3.9
2	D	33	ASP	3.8
1	B	671	LEU	3.8
2	D	31	THR	3.7
2	D	5	VAL	3.7
1	B	634	VAL	3.6
2	D	32	LEU	3.4
1	A	826	PHE	3.3
1	A	760	PRO	3.3
1	A	798	ASN	3.2
1	A	747	LEU	3.2
1	B	717	SER	3.1
1	A	670	SER	3.1
1	A	672	SER	3.1
1	B	674	VAL	3.0
2	C	30	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	762	ASP	2.8
1	A	667	LYS	2.8
1	B	638[A]	MET	2.8
1	A	638[A]	MET	2.8
1	A	802	ILE	2.8
1	A	799	PHE	2.7
1	B	632	GLY	2.7
1	B	779	ASP	2.7
2	C	32	LEU	2.6
1	B	686	ASN	2.5
1	A	673	SER	2.5
1	A	763	TYR	2.5
1	B	633	ASP	2.4
1	A	800	ILE	2.4
1	B	760	PRO	2.4
1	A	668	GLN	2.4
1	B	758	LEU	2.3
1	A	796	HIS	2.3
1	B	796	HIS	2.3
1	B	802	ILE	2.3
1	A	665	LEU	2.3
1	B	761	ASP	2.3
1	B	670	SER	2.2
1	B	685	HIS	2.2
1	A	827	THR	2.2
2	C	132	SER	2.2
1	A	761	ASP	2.2
1	B	684	ALA	2.2
1	B	665	LEU	2.2
2	D	43	ALA	2.1
1	A	779	ASP	2.1
1	B	666	THR	2.0
1	B	800	ILE	2.0
1	A	674	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CSO	B	751	7/8	0.89	0.10	35,38,44,45	0
1	CSO	A	751	7/8	0.94	0.08	44,48,52,52	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	SO4	A	906	5/5	0.24	0.16	160,160,160,160	0
3	NAG	A	902	14/15	0.61	0.13	70,71,71,72	0
4	SO4	A	907	5/5	0.78	0.09	102,103,103,103	0
4	SO4	D	202	5/5	0.83	0.11	84,84,85,85	0
4	SO4	B	904	5/5	0.85	0.12	42,42,42,42	5
3	NAG	B	902	14/15	0.86	0.10	67,67,68,68	0
3	NAG	B	901	14/15	0.87	0.09	51,53,55,56	0
4	SO4	A	905	5/5	0.89	0.13	67,67,67,67	5
4	SO4	B	903	5/5	0.90	0.11	61,61,61,62	5
4	SO4	A	903	5/5	0.91	0.09	34,35,35,36	5
4	SO4	A	904	5/5	0.91	0.11	70,70,70,70	5
3	NAG	A	901	14/15	0.93	0.07	34,35,39,40	0
4	SO4	D	201	5/5	0.94	0.10	52,52,52,52	5
4	SO4	C	201	5/5	0.97	0.09	27,28,28,28	5

6.5 Other polymers [i](#)

There are no such residues in this entry.