



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:06 AM UTC

PDB ID : 8AVL / pdb_00008avl
Title : Superoxide dismutase SodFM2 from Bacteroides fragilis
Authors : Basle, A.; Barwinska-Sendra, A.; Sendra, K.M.; Waldron, K.
Deposited on : 2022-08-26
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

i

X-RAY DIFFRACTION

A.

the following graphic. The table shows the number of entries on which the scores are based.



Percentile relative to all X-ray structures

Percentile relative to X-ray structures of similar resolution

MetricWhole archive
(#Entries)

Similar resolution
 (#Entries, resolution range(Å))

 R_{free}

Clashscore

Ramachandran outliers

Sidechain outliers

RSRZ outliers

180053

190562

187476

187428

180081

4673 (1.60-1.60)

4931 (1.60-1.60)

4831 (1.60-1.60)

4830 (1.60-1.60)

4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Quality of chain

93%

7%

0.7

92%

8%

91%

3%

95%



5%

0.49%

50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	192	 94% 5% •
1	G	192	 91% 9% •
1	H	192	 93% 7%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 25126 atoms, of which 11290 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Fe].

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	192	Total	C	H	N	O	S	0	2	0
			2964	998	1417	260	287	2			
1	B	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	C	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	D	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	E	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	F	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	G	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			
1	H	192	Total	C	H	N	O	S	0	0	0
			2947	992	1409	260	284	2			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P53638
B	1	MET	-	initiating methionine	UNP P53638
C	1	MET	-	initiating methionine	UNP P53638
D	1	MET	-	initiating methionine	UNP P53638
E	1	MET	-	initiating methionine	UNP P53638
F	1	MET	-	initiating methionine	UNP P53638
G	1	MET	-	initiating methionine	UNP P53638
H	1	MET	-	initiating methionine	UNP P53638

- Molecule 2 is FE (III) ION (CCD ID: FE) (formula: Fe) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	D	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	F	1	Total Fe 1 1	0	0
2	G	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0

- Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (CCD ID: TRS) (formula: $C_4H_{12}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	N	O	
			18	4	10	1	3	


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	203	Total 203	O 203	0	0
4	B	222	Total 222	O 222	0	0
4	C	143	Total 143	O 143	0	0
4	D	146	Total 146	O 146	0	0
4	E	196	Total 196	O 196	0	0
4	F	213	Total 213	O 213	0	0
4	G	178	Total 178	O 178	0	0
4	H	206	Total 206	O 206	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Superoxide dismutase [Fe]

Chain A: 



- Molecule 1: Superoxide dismutase [Fe]

Chain B: 




- Molecule 1: Superoxide dismutase [Fe]

Chain C: 




- Molecule 1: Superoxide dismutase [Fe]

Chain D: 



- Molecule 1: Superoxide dismutase [Fe]

Chain E: 



- Molecule 1: Superoxide dismutase [Fe]

Chain F:

94%

5% •




• Molecule 1: Superoxide dismutase [Fe]

Chain G:

91%

9% •



• Molecule 1: Superoxide dismutase [Fe]

Chain H:

93%

7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.13Å 100.11Å 103.60Å 90.00° 91.80° 90.00°	Depositor
Resolution (Å)	103.55 – 1.60 103.55 – 1.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (103.55-1.60) 100.0 (103.55-1.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.60Å)	Xtriage
Refinement program	REFMAC 5.8.0352, REFMAC 5.8.0352	Depositor
R, R_{free}	0.133 , 0.204 0.133 , 0.204	Depositor DCC
R_{free} test set	9665 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.059	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for -h,l,k 0.000 for -h,-l,-k 0.020 for h,-k,-l	Xtriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	25126	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 69.76 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.5061e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TRS, FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/1598	0.89	0/2171
1	B	0.69	0/1583	0.93	1/2150 (0.0%)
1	C	0.67	0/1583	0.89	1/2150 (0.0%)
1	D	0.67	0/1583	0.87	0/2150
1	E	0.67	0/1583	0.87	0/2150
1	F	0.71	0/1583	0.89	1/2150 (0.0%)
1	G	0.69	0/1583	0.90	1/2150 (0.0%)
1	H	0.69	0/1583	0.86	0/2150
All	All	0.69	0/12679	0.89	4/17221 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1
1	G	0	1
All	All	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	43	PRO	N-CA-C	6.58	126.02	112.47
1	G	46	GLU	CB-CG-CD	5.76	122.39	112.60
1	B	152	LYS	CA-CB-CG	5.64	125.39	114.10
1	F	89	GLU	CA-CB-CG	5.27	124.64	114.10

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	47	TYR	Sidechain
1	G	100	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1547	1417	1502	14	0
1	B	1538	1409	1491	9	0
1	C	1538	1409	1491	9	0
1	D	1538	1409	1491	5	1
1	E	1538	1409	1491	7	0
1	F	1538	1409	1491	7	0
1	G	1538	1409	1491	15	0
1	H	1538	1409	1491	14	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	B	8	10	12	0	0
4	A	203	0	0	6	1
4	B	222	0	0	1	2
4	C	143	0	0	0	1
4	D	146	0	0	1	0
4	E	196	0	0	4	1
4	F	213	0	0	2	1
4	G	178	0	0	1	0
4	H	206	0	0	4	1
All	All	13836	11290	11951	72	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:21:GLN:NE2	4:A:301:HOH:O	1.94	1.00
1:F:181:GLU:HG3	4:F:373:HOH:O	1.69	0.92
1:B:1:MET:HE3	1:B:2:TYR:CE2	2.15	0.81
1:H:96:GLU:HG2	4:H:489:HOH:O	1.82	0.80
1:C:85:PRO:HD2	1:C:181:GLU:O	1.87	0.74
1:G:131:LYS:HE2	4:H:352:HOH:O	1.91	0.70
1:B:1:MET:HE1	1:B:42:VAL:HG21	1.83	0.60
1:A:152:LYS:HD2	4:A:354:HOH:O	2.00	0.60
1:G:131:LYS:HD2	1:H:100:ARG:NH1	2.17	0.60
1:A:99:LYS:HE2	1:A:105:PHE:H	1.66	0.60
1:G:131:LYS:HG2	1:H:93:LYS:HG2	1.83	0.60
1:A:99:LYS:HD3	1:A:104:SER:HA	1.84	0.59
1:G:131:LYS:CE	4:H:352:HOH:O	2.51	0.58
4:E:306:HOH:O	1:H:110:LYS:HE3	2.03	0.58
1:G:131:LYS:HB2	1:H:93:LYS:HG3	1.87	0.57
1:C:8:PRO:HD3	1:C:79:LEU:HD23	1.87	0.56
1:E:53:GLU:HB2	4:E:448:HOH:O	2.06	0.56
1:G:130:ASP:HB2	4:G:437:HOH:O	2.06	0.56
1:B:93:LYS:HE3	1:B:192:LEU:HD13	1.90	0.53
1:C:43:PRO:O	1:C:48:GLU:OE1	2.26	0.53
1:A:19:SER:OG	1:A:21:GLN:HG2	2.09	0.53
1:C:13:ALA:HB1	1:C:83:PRO:HG3	1.90	0.52
1:F:8:PRO:HD3	1:F:79:LEU:HD23	1.90	0.52
1:H:152:LYS:HD2	4:H:466:HOH:O	2.09	0.52
1:G:19:SER:OG	1:G:21:GLN:HG2	2.10	0.51
1:G:1:MET:HE3	1:G:3:GLU:OE2	2.11	0.51
1:F:8:PRO:HD3	1:F:79:LEU:CD2	2.41	0.51
1:G:131:LYS:HD2	1:H:100:ARG:HH12	1.75	0.51
1:G:131:LYS:CE	1:H:100:ARG:HH12	2.24	0.50
1:A:99:LYS:HE3	4:A:383:HOH:O	2.11	0.50
1:H:21:GLN:HG2	1:H:166:LEU:HD13	1.94	0.50
1:E:29:LYS:NZ	4:E:302:HOH:O	2.42	0.49
4:E:306:HOH:O	1:H:110:LYS:CE	2.60	0.49
1:A:73:ASN:HB3	1:A:124:TRP:CZ2	2.49	0.48
1:E:83:PRO:O	1:E:84:LYS:C	2.56	0.48
1:A:89[A]:GLU:CD	4:A:397:HOH:O	2.56	0.47
1:G:73:ASN:HB3	1:G:124:TRP:CZ2	2.50	0.47
1:C:73:ASN:HB3	1:C:124:TRP:CZ2	2.50	0.47
1:E:73:ASN:HB3	1:E:124:TRP:CZ2	2.51	0.46
1:A:99:LYS:HE2	1:A:105:PHE:N	2.28	0.45
1:B:73:ASN:HB3	1:B:124:TRP:CZ2	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:60:PRO:HD3	1:G:58:SER:O	2.15	0.45
1:D:73:ASN:HB3	1:D:124:TRP:CZ2	2.52	0.45
1:A:96:GLU:HB2	4:A:303:HOH:O	2.17	0.45
1:D:131:LYS:HG3	4:D:370:HOH:O	2.17	0.45
1:F:73:ASN:HB3	1:F:124:TRP:CZ2	2.52	0.45
1:C:83:PRO:O	1:C:84:LYS:C	2.61	0.44
1:F:16:PRO:O	4:F:301:HOH:O	2.21	0.44
1:H:73:ASN:HB3	1:H:124:TRP:CZ2	2.53	0.44
1:A:89[A]:GLU:HB2	1:A:90:PRO:HD2	2.00	0.44
1:B:77:TYR:O	1:B:80:GLN:HG2	2.18	0.42
1:G:77:TYR:O	1:G:80:GLN:HG2	2.20	0.42
1:D:38:LEU:HD23	1:D:38:LEU:HA	1.91	0.42
1:E:21:GLN:HG2	1:E:166:LEU:HD13	2.01	0.42
1:C:89:GLU:HB2	1:C:90:PRO:HD2	2.01	0.41
1:B:8:PRO:HD3	1:B:79:LEU:HD23	2.03	0.41
1:H:31:LEU:HD13	1:H:74:HIS:HB2	2.02	0.41
1:A:77:TYR:O	1:A:80:GLN:HG2	2.21	0.41
1:A:90:PRO:HG3	1:A:98:ILE:HD12	2.03	0.41
1:B:21:GLN:HG2	1:B:166:LEU:HD13	2.02	0.41
1:C:77:TYR:O	1:C:80:GLN:HG2	2.21	0.40
1:E:77:TYR:O	1:E:80:GLN:HG2	2.21	0.40
1:E:13:ALA:HB1	1:E:83:PRO:HG3	2.04	0.40
1:B:21:GLN:OE1	1:F:21:GLN:NE2	2.55	0.40
1:C:38:LEU:HD23	1:C:38:LEU:HA	1.95	0.40
1:G:31:LEU:HD13	1:G:74:HIS:HB2	2.04	0.40
1:A:89[A]:GLU:CG	4:A:397:HOH:O	2.70	0.40
1:F:77:TYR:O	1:F:80:GLN:HG2	2.22	0.40
1:G:131:LYS:HE2	1:H:93:LYS:HE2	2.03	0.40
1:B:29:LYS:HE3	4:B:462:HOH:O	2.21	0.40
1:D:77:TYR:O	1:D:80:GLN:HG2	2.21	0.40
1:H:1:MET:HE3	1:H:3:GLU:HG2	2.03	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:317:HOH:O	4:C:395:HOH:O[2_453]	2.07	0.13
4:B:429:HOH:O	4:F:436:HOH:O[2_453]	2.11	0.09
4:A:501:HOH:O	4:E:384:HOH:O[2_443]	2.16	0.04
1:D:16:PRO:O	4:H:301:HOH:O[2_544]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	192/192 (100%)	188 (98%)	4 (2%)	0	100	100
1	B	190/192 (99%)	186 (98%)	4 (2%)	0	100	100
1	C	190/192 (99%)	185 (97%)	5 (3%)	0	100	100
1	D	190/192 (99%)	186 (98%)	4 (2%)	0	100	100
1	E	190/192 (99%)	186 (98%)	3 (2%)	1 (0%)	24	10
1	F	190/192 (99%)	186 (98%)	4 (2%)	0	100	100
1	G	190/192 (99%)	186 (98%)	4 (2%)	0	100	100
1	H	190/192 (99%)	187 (98%)	3 (2%)	0	100	100
All	All	1522/1536 (99%)	1490 (98%)	31 (2%)	1 (0%)	48	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	84	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	162/160 (101%)	162 (100%)	0	100	100
1	B	160/160 (100%)	158 (99%)	2 (1%)	61	40
1	C	160/160 (100%)	158 (99%)	2 (1%)	61	40
1	D	160/160 (100%)	159 (99%)	1 (1%)	78	66

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	160/160 (100%)	160 (100%)	0	100	100
1	F	160/160 (100%)	159 (99%)	1 (1%)	78	66
1	G	160/160 (100%)	156 (98%)	4 (2%)	42	18
1	H	160/160 (100%)	160 (100%)	0	100	100
All	All	1282/1280 (100%)	1272 (99%)	10 (1%)	73	59

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1	MET
1	B	190	LYS
1	C	7	LEU
1	C	38	LEU
1	D	89	GLU
1	F	89	GLU
1	G	87	LYS
1	G	93	LYS
1	G	131	LYS
1	G	132	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	32	GLN
1	B	36	ASN
1	D	11	ASN
1	F	21	GLN
1	F	88	ASN
1	G	11	ASN
1	G	32	GLN
1	H	107	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 8 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	TRS	B	201	-	7,7,7	0.35	0	9,9,9	0.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	TRS	B	201	-	-	0/9/9/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	192/192 (100%)	-0.92	0 100 100	10, 21, 46, 75	2 (1%)
1	B	192/192 (100%)	-0.94	1 (0%) 87 90	14, 20, 43, 72	0
1	C	192/192 (100%)	-0.64	0 100 100	16, 29, 55, 83	0
1	D	192/192 (100%)	-0.86	0 100 100	15, 22, 48, 64	0
1	E	192/192 (100%)	-0.92	0 100 100	15, 22, 39, 68	0
1	F	192/192 (100%)	-0.90	0 100 100	14, 20, 44, 70	0
1	G	192/192 (100%)	-0.79	0 100 100	16, 23, 48, 66	0
1	H	192/192 (100%)	-0.90	0 100 100	15, 23, 42, 90	0
All	All	1536/1536 (100%)	-0.86	1 (0%) 92 92	10, 22, 46, 90	2 (0%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	MET	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	TRS	B	201	8/8	0.99	0.03	22,27,29,30	0
2	FE	B	202	1/1	1.00	0.06	24,24,24,24	0
2	FE	C	201	1/1	1.00	0.08	28,28,28,28	0
2	FE	D	201	1/1	1.00	0.06	23,23,23,23	0
2	FE	E	201	1/1	1.00	0.04	25,25,25,25	0
2	FE	F	201	1/1	1.00	0.05	23,23,23,23	0
2	FE	G	201	1/1	1.00	0.06	26,26,26,26	0
2	FE	H	201	1/1	1.00	0.06	24,24,24,24	0
2	FE	A	201	1/1	1.00	0.06	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FE B 202:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE D 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE E 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE F 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE G 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FE H 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE A 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.