



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:22 AM UTC

PDB ID : 2AWN / pdb_00002awn
Title : Crystal structure of the ADP-Mg-bound E. Coli MALK (Crystallized with ATP-Mg)
Authors : Lu, G.; Westbrook, J.M.; Davidson, A.L.; Chen, J.
Deposited on : 2005-09-01
Resolution : 2.30 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	381	
1	B	381	
1	C	381	
1	D	381	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose/maltodextrin import ATP-binding protein malK.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	330	2534	1605	445	472	12	0	0	0
1	B	374	2887	1825	515	534	13	0	0	0
1	C	344	2596	1642	466	476	12	0	0	0
1	D	301	2334	1475	412	435	12	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	372	ALA	-	cloning artifact	UNP P68187
A	373	SER	-	cloning artifact	UNP P68187
A	374	ALA	-	cloning artifact	UNP P68187
A	375	SER	-	cloning artifact	UNP P68187
A	376	HIS	-	cloning artifact	UNP P68187
A	377	HIS	-	cloning artifact	UNP P68187
A	378	HIS	-	cloning artifact	UNP P68187
A	379	HIS	-	cloning artifact	UNP P68187
A	380	HIS	-	cloning artifact	UNP P68187
A	381	HIS	-	cloning artifact	UNP P68187
B	372	ALA	-	cloning artifact	UNP P68187
B	373	SER	-	cloning artifact	UNP P68187
B	374	ALA	-	cloning artifact	UNP P68187
B	375	SER	-	cloning artifact	UNP P68187
B	376	HIS	-	cloning artifact	UNP P68187
B	377	HIS	-	cloning artifact	UNP P68187
B	378	HIS	-	cloning artifact	UNP P68187
B	379	HIS	-	cloning artifact	UNP P68187
B	380	HIS	-	cloning artifact	UNP P68187
B	381	HIS	-	cloning artifact	UNP P68187
C	372	ALA	-	cloning artifact	UNP P68187

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	373	SER	-	cloning artifact	UNP P68187
C	374	ALA	-	cloning artifact	UNP P68187
C	375	SER	-	cloning artifact	UNP P68187
C	376	HIS	-	cloning artifact	UNP P68187
C	377	HIS	-	cloning artifact	UNP P68187
C	378	HIS	-	cloning artifact	UNP P68187
C	379	HIS	-	cloning artifact	UNP P68187
C	380	HIS	-	cloning artifact	UNP P68187
C	381	HIS	-	cloning artifact	UNP P68187
D	372	ALA	-	cloning artifact	UNP P68187
D	373	SER	-	cloning artifact	UNP P68187
D	374	ALA	-	cloning artifact	UNP P68187
D	375	SER	-	cloning artifact	UNP P68187
D	376	HIS	-	cloning artifact	UNP P68187
D	377	HIS	-	cloning artifact	UNP P68187
D	378	HIS	-	cloning artifact	UNP P68187
D	379	HIS	-	cloning artifact	UNP P68187
D	380	HIS	-	cloning artifact	UNP P68187
D	381	HIS	-	cloning artifact	UNP P68187

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	44	Total	O	0	0
			44	44		
4	B	81	Total	O	0	0
			81	81		
4	C	36	Total	O	0	0
			36	36		
4	D	45	Total	O	0	0
			45	45		

4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.31Å 102.67Å 130.78Å 90.00° 90.81° 90.00°	Depositor
Resolution (Å)	37.28 – 2.30 37.28 – 2.30	Depositor EDS
% Data completeness (in resolution range)	85.0 (37.28-2.30) 94.5 (37.28-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.29Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.230 , 0.271 0.231 , 0.274	Depositor DCC
R_{free} test set	3843 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	52.4	Xtrriage
Anisotropy	0.390	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.023 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10669	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.34	3/2576 (0.1%)	1.50	27/3500 (0.8%)
1	B	0.75	3/2937 (0.1%)	1.20	29/3984 (0.7%)
1	C	0.71	3/2637 (0.1%)	1.32	23/3581 (0.6%)
1	D	1.15	4/2375 (0.2%)	1.27	24/3224 (0.7%)
All	All	1.01	13/10525 (0.1%)	1.32	103/14289 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	2
1	D	0	1
All	All	0	5

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	298	VAL	N-CA	48.38	2.06	1.46
1	A	299	ILE	N-CA	46.71	2.10	1.46
1	A	298	VAL	N-CA	39.75	1.96	1.46
1	C	297	ASP	N-CA	26.73	1.80	1.46
1	B	327	LEU	N-CA	24.86	1.78	1.46

The worst 5 of 103 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	297	ASP	CA-C-N	29.95	159.17	122.93
1	C	297	ASP	C-N-CA	29.95	159.17	122.93
1	A	294	ASP	N-CA-C	-29.22	74.07	112.93
1	A	298	VAL	CA-C-N	-27.56	80.79	121.85
1	A	298	VAL	C-N-CA	-27.56	80.79	121.85

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	293	SER	Peptide
1	B	326	ASN	Peptide
1	C	18	VAL	Mainchain
1	C	296	ALA	Peptide
1	D	297	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2534	0	2562	155	0
1	B	2887	0	2946	172	0
1	C	2596	0	2624	235	1
1	D	2334	0	2367	140	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	27	0	12	1	0
3	B	27	0	12	3	0
3	C	27	0	12	3	0
3	D	27	0	12	0	0
4	A	44	0	0	1	0
4	B	81	0	0	6	1
4	C	36	0	0	2	0
4	D	45	0	0	1	0
All	All	10669	0	10547	696	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 696 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:18:VAL:CG2	1:C:44:THR:HG21	1.31	1.58
1:C:296:ALA:CB	1:C:299:ILE:CG1	1.83	1.54
1:B:327:LEU:N	1:B:327:LEU:CA	1.78	1.45
1:C:297:ASP:CA	1:C:297:ASP:N	1.80	1.44
1:B:319:GLN:CA	1:B:326:ASN:HD21	1.29	1.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:102:LEU:CD2	4:B:765:HOH:O[2_556]	1.97	0.23

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/381 (85%)	294 (91%)	20 (6%)	10 (3%)	3	2
1	B	372/381 (98%)	339 (91%)	24 (6%)	9 (2%)	4	4
1	C	338/381 (89%)	290 (86%)	29 (9%)	19 (6%)	1	0
1	D	297/381 (78%)	263 (89%)	23 (8%)	11 (4%)	2	1
All	All	1331/1524 (87%)	1186 (89%)	96 (7%)	49 (4%)	2	1

5 of 49 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	236	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	293	SER
1	A	298	VAL
1	B	236	SER
1	B	323	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/323 (86%)	260 (94%)	17 (6%)	17	24
1	B	315/323 (98%)	297 (94%)	18 (6%)	18	27
1	C	277/323 (86%)	264 (95%)	13 (5%)	23	35
1	D	258/323 (80%)	239 (93%)	19 (7%)	13	17
All	All	1127/1292 (87%)	1060 (94%)	67 (6%)	18	26

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	228	ARG
1	D	275	VAL
1	D	350	GLU
1	B	183	LEU
1	B	144	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 53 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	331	GLN
1	C	163	ASN
1	D	276	GLN
1	B	332	ASN
1	C	27	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	401	2	28,29,29	1.44	4 (14%)	43,45,45	1.96	6 (13%)
3	ADP	C	403	2	28,29,29	1.20	3 (10%)	43,45,45	1.85	6 (13%)
3	ADP	B	402	2	28,29,29	1.57	4 (14%)	43,45,45	1.96	8 (18%)
3	ADP	D	404	2	28,29,29	1.28	2 (7%)	43,45,45	1.83	6 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	401	2	-	6/16/32/32	0/3/3/3
3	ADP	C	403	2	-	4/16/32/32	0/3/3/3
3	ADP	B	402	2	-	6/16/32/32	0/3/3/3
3	ADP	D	404	2	-	2/16/32/32	0/3/3/3

The worst 5 of 13 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	402	ADP	PA-O3A	-4.21	1.55	1.59
3	B	402	ADP	C2-N1	3.45	1.40	1.33
3	D	404	ADP	C2-N1	3.45	1.40	1.33
3	A	401	ADP	C2-N1	3.43	1.40	1.33
3	C	403	ADP	C2-N1	3.31	1.39	1.33

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	ADP	O5'-PA-O1A	-8.58	74.93	108.94
3	C	403	ADP	O5'-PA-O1A	-8.28	76.14	108.94
3	D	404	ADP	O5'-PA-O1A	-8.03	77.12	108.94
3	B	402	ADP	O5'-PA-O1A	-7.93	77.51	108.94
3	B	402	ADP	O2A-PA-O5'	-4.34	87.88	107.57

There are no chirality outliers.

5 of 18 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	402	ADP	PA-O3A-PB-O3B
3	C	403	ADP	PA-O3A-PB-O3B
3	D	404	ADP	PA-O3A-PB-O2B
3	C	403	ADP	C3'-C4'-C5'-O5'
3	A	401	ADP	O4'-C4'-C5'-O5'


There are no ring outliers.


3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	ADP	1	0
3	C	403	ADP	3	0
3	B	402	ADP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	330/381 (86%)	0.86	44 (13%) 7 8	39, 63, 107, 146	0
1	B	374/381 (98%)	0.50	20 (5%) 32 34	32, 52, 87, 119	0
1	C	344/381 (90%)	1.33	86 (25%) 2 2	38, 76, 130, 148	0
1	D	301/381 (79%)	0.84	37 (12%) 8 9	34, 64, 102, 115	0
All	All	1349/1524 (88%)	0.88	187 (13%) 6 7	32, 62, 113, 148	0

The worst 5 of 187 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	324	ARG	6.2
1	D	323	ILE	6.0
1	A	295	ILE	5.9
1	B	326	ASN	5.8
1	C	102	LEU	5.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	MG	C	503	1/1	0.35	0.40	109,109,109,109	0
3	ADP	C	403	27/27	0.78	0.38	85,110,116,117	27
2	MG	D	504	1/1	0.86	0.09	76,76,76,76	0
3	ADP	A	401	27/27	0.91	0.13	38,74,84,94	0
3	ADP	B	402	27/27	0.92	0.15	40,56,73,81	0
3	ADP	D	404	27/27	0.94	0.10	54,92,100,106	0
2	MG	B	502	1/1	0.96	0.10	48,48,48,48	0
2	MG	A	501	1/1	0.97	0.05	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around ADP A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.