



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:14 AM UTC

PDB ID : 3AWF / pdb_00003awf
Title : Crystal structure of Pten-like domain of Ci-VSP (236-576)
Authors : Matsuda, M.; Sakata, S.; Takeshita, K.; Suzuki, M.; Yamashita, E.; Okamura, Y.; Nakagawa, A.
Deposited on : 2011-03-19
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	
1	B	346	
1	C	346	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	C	1	-	X	-	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8081 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Voltage-sensor containing phosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	304	Total	C	N	O	S	0	5	0
			2464	1569	415	465	15			
1	B	308	Total	C	N	O	S	0	4	0
			2510	1596	429	470	15			
1	C	310	Total	C	N	O	S	0	4	0
			2516	1596	431	474	15			

There are 15 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	231	GLY	-	expression tag	UNP Q4W8A1
A	232	PRO	-	expression tag	UNP Q4W8A1
A	233	LEU	-	expression tag	UNP Q4W8A1
A	234	GLY	-	expression tag	UNP Q4W8A1
A	235	SER	-	expression tag	UNP Q4W8A1
B	231	GLY	-	expression tag	UNP Q4W8A1
B	232	PRO	-	expression tag	UNP Q4W8A1
B	233	LEU	-	expression tag	UNP Q4W8A1
B	234	GLY	-	expression tag	UNP Q4W8A1
B	235	SER	-	expression tag	UNP Q4W8A1
C	231	GLY	-	expression tag	UNP Q4W8A1
C	232	PRO	-	expression tag	UNP Q4W8A1
C	233	LEU	-	expression tag	UNP Q4W8A1
C	234	GLY	-	expression tag	UNP Q4W8A1
C	235	SER	-	expression tag	UNP Q4W8A1

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		
2	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	195	Total	O	0	0
			195	195		
4	B	176	Total	O	0	0
			176	176		
4	C	164	Total	O	0	0
			164	164		

- Molecule 1: Voltage-sensor containing phosphatase





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	131.56Å 177.44Å 50.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.17 – 1.99 44.17 – 1.99	Depositor EDS
% Data completeness (in resolution range)	98.8 (44.17-1.99) 98.8 (44.17-1.99)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.13 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0071	Depositor
R, R_{free}	0.179 , 0.223 0.179 , 0.225	Depositor DCC
R_{free} test set	4094 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	24.4	Xtriage
Anisotropy	0.053	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8081	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.44	8/2525 (0.3%)	1.14	4/3407 (0.1%)
1	B	1.33	5/2566 (0.2%)	1.15	6/3462 (0.2%)
1	C	1.33	6/2572 (0.2%)	1.16	6/3468 (0.2%)
All	All	1.37	19/7663 (0.2%)	1.15	16/10337 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	2
All	All	0	5

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	502	VAL	CA-CB	7.52	1.62	1.54
1	A	359	ILE	C-O	6.73	1.30	1.24
1	C	417	ARG	C-O	-6.14	1.16	1.24
1	C	562	ASP	C-O	-6.09	1.16	1.24
1	A	314	GLY	N-CA	6.03	1.49	1.44
1	C	464	ILE	CA-CB	-5.81	1.47	1.54
1	A	271	VAL	CA-CB	5.63	1.61	1.54
1	B	418	TYR	C-O	-5.52	1.17	1.24
1	A	410	VAL	C-O	-5.47	1.17	1.23
1	B	444	VAL	CA-CB	5.45	1.62	1.54
1	C	277	PRO	CA-C	5.41	1.57	1.52
1	C	393	TYR	C-O	-5.39	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	372	THR	CA-C	5.30	1.59	1.52
1	B	507	ILE	C-O	-5.16	1.18	1.24
1	A	362	HIS	C-O	-5.15	1.17	1.23
1	B	506	ASP	C-O	-5.09	1.17	1.24
1	C	392	GLU	C-O	-5.07	1.18	1.24
1	B	460	LEU	N-CA	5.06	1.52	1.45
1	A	527	PHE	N-CA	5.04	1.52	1.46

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	365	GLY	N-CA-C	-9.54	100.33	112.54
1	C	311	SER	N-CA-C	-9.35	100.45	112.23
1	B	433	LEU	CA-C-N	-9.28	110.83	120.38
1	B	433	LEU	C-N-CA	-9.28	110.83	120.38
1	B	367	LYS	N-CA-C	7.40	126.56	110.80
1	A	502	VAL	N-CA-CB	5.70	117.32	110.31
1	A	258	ASP	N-CA-C	-5.63	106.26	113.02
1	B	391	LEU	N-CA-C	5.42	117.61	111.11
1	A	468	ARG	CA-C-N	-5.34	114.28	122.62
1	A	468	ARG	C-N-CA	-5.34	114.28	122.62
1	C	354	ASP	CA-C-N	5.33	125.14	119.28
1	C	354	ASP	C-N-CA	5.33	125.14	119.28
1	B	480	TYR	CA-CB-CG	5.31	123.45	113.90
1	C	287	ASN	CA-C-N	5.29	125.55	119.83
1	C	287	ASN	C-N-CA	5.29	125.55	119.83
1	C	267	VAL	N-CA-C	-5.29	105.24	111.00

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	362	HIS	Peptide
1	B	362	HIS	Peptide
1	B	366	GLY	Peptide
1	C	362	HIS	Peptide
1	C	571	SER	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2464	0	2417	23	0
1	B	2510	0	2461	18	0
1	C	2516	0	2459	22	0
2	A	15	0	0	0	0
2	B	15	0	0	0	0
2	C	20	0	0	0	0
3	C	6	0	8	2	0
4	A	195	0	0	4	0
4	B	176	0	0	1	0
4	C	164	0	0	2	0
All	All	8081	0	7345	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:466:SER:O	1:B:469:GLN:HG2	1.77	0.83
1:A:546[A]:GLU:HG2	4:A:578:HOH:O	1.81	0.79
1:C:572:ASP:OD1	1:C:572:ASP:C	2.29	0.76
1:C:424:LYS:HD3	1:C:535:LEU:HD22	1.69	0.73
1:A:421:TYR:CD2	1:A:535[B]:LEU:HD22	2.24	0.72
1:C:405:ASP:HB3	1:C:408:GLN:HG3	1.78	0.64
1:B:572:ASP:OD1	1:B:572:ASP:C	2.40	0.63
1:B:313:ARG:NH1	4:B:229:HOH:O	2.32	0.62
1:A:456:ASN:C	1:A:456:ASN:HD22	2.08	0.62
1:C:341:LEU:HD12	1:C:433:LEU:HD13	1.83	0.60
1:A:380:GLU:OE1	1:A:426:LYS:HE3	2.01	0.60
1:B:424:LYS:HD3	1:B:535:LEU:HD22	1.84	0.59
1:B:341:LEU:HD12	1:B:433:LEU:HD13	1.84	0.59
1:C:303:LYS:HD3	1:C:356:ASP:CB	2.34	0.58
1:C:366:GLY:HA2	4:C:641:HOH:O	2.05	0.57
1:C:303:LYS:HD3	1:C:356:ASP:HB3	1.89	0.55
1:C:478:GLU:HG3	1:C:480:TYR:HB2	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:ASN:HB3	1:A:292:VAL:CG2	2.38	0.54
1:C:545:ARG:HB3	3:C:1:GOL:H12	1.89	0.53
1:A:333:ASN:OD1	1:A:508:LYS:HE3	2.10	0.51
1:A:367:LYS:HD3	1:A:398:ARG:CZ	2.40	0.51
1:C:516:LYS:HE3	1:C:516:LYS:H	1.75	0.51
1:B:303:LYS:HD3	1:B:356:ASP:HB2	1.93	0.51
1:A:421:TYR:CG	1:A:535[B]:LEU:HD22	2.46	0.51
1:B:341:LEU:CD1	1:B:433:LEU:HD13	2.41	0.50
1:B:367:LYS:HG3	1:B:368:GLY:N	2.25	0.50
1:A:437:LYS:HB2	1:A:534:SER:HA	1.94	0.50
1:A:276:PHE:HB2	1:A:364:LYS:HD3	1.94	0.50
1:C:549:ASP:O	1:C:550:ASN:HB2	2.12	0.49
1:A:367:LYS:HG3	1:A:411:GLU:HG3	1.94	0.49
1:C:270:HIS:HA	1:C:358:VAL:HG12	1.94	0.49
1:A:405:ASP:HB3	1:A:408:GLN:HG2	1.94	0.49
1:A:366:GLY:HA2	4:A:39:HOH:O	2.13	0.49
1:C:475:LYS:HB2	1:C:480:TYR:HB3	1.95	0.48
1:A:348:LYS:HD2	1:A:380:GLU:OE2	2.14	0.48
1:C:367:LYS:HG2	1:C:398:ARG:CZ	2.44	0.47
1:A:456:ASN:ND2	1:A:458:SER:H	2.13	0.47
1:A:330:ASP:OD1	1:A:468:ARG:NH2	2.47	0.47
1:C:303:LYS:HD3	1:C:356:ASP:HB2	1.96	0.47
1:C:511:PHE:O	1:C:526:PRO:HD2	2.15	0.46
1:B:494[A]:THR:HG22	1:B:495[A]:CYS:N	2.30	0.46
1:B:367:LYS:HG3	1:B:411:GLU:HG3	1.97	0.46
1:A:485:GLN:HG3	4:A:675:HOH:O	2.17	0.45
1:B:301:PRO:O	1:B:302:ASP:HB2	2.17	0.45
1:C:550:ASN:HB2	1:C:551:PRO:HD3	1.99	0.45
1:B:445:THR:HG23	1:B:494[B]:THR:HG22	1.98	0.44
1:B:494[A]:THR:CG2	1:B:495[A]:CYS:N	2.81	0.44
1:C:545:ARG:HB3	3:C:1:GOL:C1	2.47	0.44
1:B:287:ASN:HB3	1:B:292:VAL:CG2	2.48	0.43
1:A:518:LEU:HD12	1:A:519:PRO:HD2	2.00	0.43
1:C:437:LYS:HD3	1:C:439:LEU:HD21	2.00	0.43
1:A:397:ARG:HE	1:A:397:ARG:HB3	1.62	0.43
1:C:301:PRO:O	1:C:302:ASP:HB2	2.18	0.43
1:C:253:ARG:NH1	1:C:256:ARG:HH22	2.17	0.42
1:A:278:SER:O	1:A:288:PRO:HA	2.20	0.41
1:A:437:LYS:HE2	1:A:437:LYS:HB3	1.56	0.41
1:B:443:GLY:HA2	1:B:495[B]:CYS:O	2.20	0.41
1:A:459:ASP:OD1	1:A:513:SER:OG	2.34	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:LEU:HD21	1:B:425:ILE:HD11	2.02	0.41
1:C:408:GLN:HG2	4:C:665:HOH:O	2.19	0.41
1:A:469:GLN:NE2	4:A:683:HOH:O	2.54	0.41
1:B:541[B]:VAL:HG13	1:B:570:PHE:HE1	1.86	0.40
1:B:448:ALA:HB3	1:B:564:PHE:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/346 (88%)	294 (97%)	8 (3%)	1 (0%)	36	35
1	B	306/346 (88%)	297 (97%)	8 (3%)	1 (0%)	36	35
1	C	308/346 (89%)	302 (98%)	5 (2%)	1 (0%)	36	35
All	All	917/1038 (88%)	893 (97%)	21 (2%)	3 (0%)	36	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	550	ASN
1	B	550	ASN
1	C	550	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/307 (90%)	270 (98%)	5 (2%)	51	58
1	B	279/307 (91%)	270 (97%)	9 (3%)	34	35
1	C	279/307 (91%)	264 (95%)	15 (5%)	20	17
All	All	833/921 (90%)	804 (96%)	29 (4%)	32	32

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	309	LEU
1	A	426	LYS
1	A	456	ASN
1	A	502	VAL
1	A	516	LYS
1	B	257	LYS
1	B	278	SER
1	B	294	ARG
1	B	398	ARG
1	B	433	LEU
1	B	467	GLU
1	B	480	TYR
1	B	518	LEU
1	B	572	ASP
1	C	253	ARG
1	C	257	LYS
1	C	279	SER
1	C	294	ARG
1	C	313	ARG
1	C	364	LYS
1	C	367	LYS
1	C	398	ARG
1	C	426	LYS
1	C	433	LEU
1	C	437	LYS
1	C	502	VAL
1	C	516	LYS
1	C	520	ARG
1	C	572	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	428	ASN
1	A	456	ASN
1	B	332	HIS
1	B	408	GLN
1	B	499	ASN
1	C	332	HIS
1	C	499	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1	-	4,4,4	0.26	0	6,6,6	0.61	0
2	SO4	C	10	-	4,4,4	0.23	0	6,6,6	0.36	0
2	SO4	C	7	-	4,4,4	0.35	0	6,6,6	0.88	0
2	SO4	B	2	-	4,4,4	0.50	0	6,6,6	0.48	0
2	SO4	C	3	-	4,4,4	0.16	0	6,6,6	0.50	0
3	GOL	C	1	-	5,5,5	0.76	0	5,5,5	2.30	4 (80%)
2	SO4	B	5	-	4,4,4	0.18	0	6,6,6	0.79	0
2	SO4	C	4	-	4,4,4	0.28	0	6,6,6	0.38	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	9	-	4,4,4	0.33	0	6,6,6	0.42	0
2	SO4	B	8	-	4,4,4	0.53	0	6,6,6	1.02	0
2	SO4	A	6	-	4,4,4	0.29	0	6,6,6	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	1	-	-	4/4/4/4	-

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	GOL	C3-C2-C1	-2.95	100.97	111.80
3	C	1	GOL	O2-C2-C3	2.48	119.45	109.18
3	C	1	GOL	O3-C3-C2	2.37	121.04	110.38
3	C	1	GOL	O1-C1-C2	-2.14	100.76	110.38

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	GOL	O1-C1-C2-O2
3	C	1	GOL	C1-C2-C3-O3
3	C	1	GOL	O1-C1-C2-C3
3	C	1	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/346 (87%)	-0.13	3 (0%) 79 79	13, 23, 40, 56	5 (1%)
1	B	308/346 (89%)	-0.01	7 (2%) 61 60	11, 26, 42, 53	4 (1%)
1	C	310/346 (89%)	0.09	9 (2%) 53 52	10, 28, 45, 55	4 (1%)
All	All	922/1038 (88%)	-0.02	19 (2%) 63 63	10, 25, 43, 56	13 (1%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	404	GLY	4.0
1	C	280	GLY	3.5
1	C	365	GLY	3.5
1	C	253	ARG	3.3
1	A	404	GLY	2.9
1	B	366	GLY	2.8
1	B	469	GLN	2.8
1	B	399	THR	2.7
1	B	450	GLN	2.6
1	A	257	LYS	2.6
1	C	279	SER	2.4
1	A	399	THR	2.4
1	C	522	TYR	2.2
1	B	254	ARG	2.2
1	B	365	GLY	2.2
1	C	353	SER	2.2
1	B	363	CYS	2.2
1	C	572	ASP	2.1
1	C	302	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	9	5/5	0.78	0.13	71,71,74,74	0
2	SO4	C	4	5/5	0.80	0.15	84,85,86,86	0
2	SO4	B	5	5/5	0.89	0.12	56,56,58,62	0
2	SO4	B	8	5/5	0.91	0.12	33,36,44,46	0
2	SO4	C	7	5/5	0.92	0.11	29,38,41,44	0
2	SO4	C	10	5/5	0.92	0.08	57,60,61,61	0
2	SO4	C	3	5/5	0.95	0.12	48,50,53,53	0
3	GOL	C	1	6/6	0.95	0.09	21,30,34,35	0
2	SO4	A	1	5/5	0.98	0.06	34,34,37,38	0
2	SO4	A	6	5/5	0.98	0.06	28,32,35,37	0
2	SO4	B	2	5/5	0.99	0.04	23,25,28,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.