



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 06:40 AM UTC

PDB ID : 4AWZ / pdb_00004awz
Title : Crystal Structure of the Mobile Metallo-beta-Lactamase AIM-1 from *Pseudomonas aeruginosa*: Insights into Antibiotic Binding and the role of Gln157
Authors : Leiros, H.-K.S.; Borra, P.S.; Brandsdal, B.O.; Edvardsen, K.S.W.; Spencer, J.; Walsh, T.R.; Samuelsen, O.
Deposited on : 2012-06-06
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	% 77% 11% • 11%
1	B	303	% 77% 10% • 12%
1	C	303	% 78% 8% • 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6952 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called METALLO-BETA-LACTAMASE AIM-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	0	8	0
			2053	1277	368	395	13			
1	B	267	Total	C	N	O	S	0	8	0
			2049	1279	369	389	12			
1	C	269	Total	C	N	O	S	0	6	0
			2038	1270	366	390	12			

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		
2	B	1	Total	Ca	0	0
			1	1		
2	C	1	Total	Ca	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	2	Total 2	Zn 2	0	0
4	C	2	Total 2	Zn 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	328	Total 328	O 328	0	0
5	B	243	Total 243	O 243	0	0
5	C	229	Total 229	O 229	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: METALLO-BETA-LACTAMASE AIM-1



• Molecule 1: METALLO-BETA-LACTAMASE AIM-1



• Molecule 1: METALLO-BETA-LACTAMASE AIM-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	79.47 Å 79.47 Å 229.27 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.80 20.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.8 (20.00-1.80) 98.9 (20.00-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 1.80 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.139 , 0.173 0.149 , 0.181	Depositor DCC
R_{free} test set	3799 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	11.9	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 44.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.058 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6952	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.85% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.14	11/2101 (0.5%)	0.96	6/2877 (0.2%)
1	B	1.33	11/2097 (0.5%)	0.99	4/2867 (0.1%)
1	C	1.49	10/2086 (0.5%)	0.95	8/2856 (0.3%)
All	All	1.33	32/6284 (0.5%)	0.97	18/8600 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
All	All	0	3

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	252	ALA	C-N	29.55	1.72	1.33
1	B	1227	ASP	C-N	26.73	1.61	1.33
1	C	76[A]	ASP	C-N	24.70	1.68	1.33
1	C	76[B]	ASP	C-N	24.70	1.68	1.33
1	B	277	ALA	C-N	23.31	1.66	1.33
1	A	1227	ASP	C-N	17.52	1.49	1.33
1	C	57	THR	C-N	17.16	1.57	1.33
1	C	227	ASP	C-N	16.77	1.58	1.33
1	C	1227	ASP	C-N	15.00	1.48	1.33
1	B	45[A]	LEU	C-N	14.51	1.52	1.33
1	B	45[B]	LEU	C-N	14.51	1.52	1.33
1	A	76	ASP	C-N	14.09	1.55	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	227	ASP	C-N	13.34	1.53	1.33
1	C	45	LEU	C-N	12.37	1.49	1.33
1	A	57	THR	C-N	12.29	1.50	1.34
1	A	45[A]	LEU	C-N	12.16	1.49	1.33
1	A	45[B]	LEU	C-N	12.16	1.49	1.33
1	A	252	ALA	C-N	11.39	1.48	1.33
1	C	87	THR	C-N	10.84	1.46	1.34
1	B	252	ALA	C-N	10.79	1.47	1.33
1	B	76	ASP	C-N	10.21	1.49	1.33
1	C	277	ALA	C-N	9.88	1.47	1.33
1	A	227	ASP	C-N	9.79	1.48	1.33
1	B	166	PRO	C-N	9.42	1.47	1.33
1	C	166	PRO	C-N	8.22	1.45	1.33
1	B	57	THR	C-N	7.68	1.44	1.34
1	B	87	THR	C-N	6.92	1.43	1.34
1	B	191	ALA	C-N	6.69	1.44	1.34
1	A	166	PRO	C-N	6.66	1.42	1.33
1	A	277	ALA	C-N	6.18	1.45	1.34
1	A	191	ALA	C-N	5.25	1.42	1.34
1	A	87	THR	C-N	5.01	1.39	1.34

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	277	ALA	O-C-N	-19.97	98.25	122.81
1	A	1227	ASP	O-C-N	-9.71	108.26	122.43
1	C	30	ARG	N-CA-C	9.71	121.80	111.03
1	A	1227	ASP	CA-C-N	9.44	133.49	122.35
1	A	1227	ASP	C-N-CA	9.44	133.49	122.35
1	A	76	ASP	O-C-N	-8.20	111.14	122.46
1	C	1227	ASP	O-C-N	-7.64	111.27	122.43
1	C	252	ALA	O-C-N	-7.63	111.93	122.46
1	C	1227	ASP	CA-C-N	7.36	131.39	122.63
1	C	1227	ASP	C-N-CA	7.36	131.39	122.63
1	B	1227	ASP	O-C-N	-6.69	113.14	122.37
1	C	191	ALA	O-C-N	-6.39	115.39	123.06
1	A	76	ASP	CA-C-N	6.14	133.48	122.06
1	A	76	ASP	C-N-CA	6.14	133.48	122.06
1	B	252	ALA	O-C-N	-5.27	115.25	122.46
1	B	277	ALA	N-CA-C	-5.14	103.36	110.35
1	C	45	LEU	CA-C-N	-5.12	113.59	120.87
1	C	45	LEU	C-N-CA	-5.12	113.59	120.87

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1227	ASP	Mainchain
1	B	45[A]	LEU	Mainchain
1	C	1227	ASP	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2053	0	1967	26	0
1	B	2049	0	1975	21	0
1	C	2038	0	1956	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	2	0	0	0	0
5	A	328	0	0	8	2
5	B	243	0	0	9	3
5	C	229	0	0	4	1
All	All	6952	0	5898	68	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:252:ALA:C	1:C:254:LEU:N	1.72	1.48
1:C:76[A]:ASP:C	1:C:78:ALA:N	1.68	1.47
1:A:57:THR:C	1:A:66[B]:CYS:N	1.73	1.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:76[B]:ASP:C	1:C:78:ALA:N	1.87	1.31
1:B:170:ASN:HB2	5:B:4058:HOH:O	1.68	0.93
1:A:170[B]:ASN:ND2	5:A:4200:HOH:O	2.02	0.91
1:B:45[A]:LEU:HD12	1:B:45[A]:LEU:N	1.97	0.78
1:B:174[B]:LEU:HD22	5:B:4141:HOH:O	1.82	0.78
1:A:300:ARG:NH2	5:A:4309:HOH:O	2.15	0.78
1:C:300:ARG:NH2	5:C:4218:HOH:O	2.14	0.77
1:A:45[A]:LEU:HD12	1:A:45[A]:LEU:N	2.00	0.77
1:A:300:ARG:HD2	5:A:4316:HOH:O	1.88	0.73
1:C:249:ARG:NH2	5:C:4181:HOH:O	2.23	0.71
1:A:300:ARG:CZ	1:B:281:ALA:O	2.41	0.67
1:C:82:LEU:HD21	1:C:128:LEU:HD13	1.80	0.64
1:C:252:ALA:C	1:C:254:LEU:CA	2.69	0.63
1:A:45[A]:LEU:HD12	1:A:45[A]:LEU:H	1.65	0.61
1:B:50:GLY:HA2	1:B:209[A]:GLU:HG2	1.81	0.61
1:A:107:GLU:CG	5:A:4106:HOH:O	2.50	0.60
1:B:45[A]:LEU:HD12	1:B:45[A]:LEU:H	1.65	0.59
1:C:75:SER:H	1:C:80:HIS:HD2	1.53	0.57
1:B:229:ARG:HD3	5:B:4176:HOH:O	2.04	0.56
1:C:191:ALA:O	1:C:249:ARG:NH1	2.39	0.56
1:A:147:LYS:HD3	1:A:171:ILE:HD12	1.88	0.56
1:B:130:LYS:NZ	5:B:4099:HOH:O	2.29	0.56
1:A:300:ARG:HD3	5:A:4317:HOH:O	2.06	0.54
1:B:80:HIS:HE1	5:B:4066:HOH:O	1.92	0.52
1:C:107:GLU:CD	1:C:107:GLU:H	2.18	0.52
1:A:107:GLU:HG3	5:A:4106:HOH:O	2.09	0.51
1:A:45[A]:LEU:N	1:A:45[A]:LEU:CD1	2.73	0.51
1:B:75:SER:H	1:B:80:HIS:HD2	1.58	0.51
1:C:140:ALA:HA	1:C:143[B]:ILE:HD12	1.94	0.50
1:B:170:ASN:ND2	5:B:4137:HOH:O	2.43	0.50
1:C:80:HIS:HE1	5:C:4057:HOH:O	1.94	0.50
1:A:75:SER:H	1:A:80:HIS:HD2	1.60	0.49
1:B:174[B]:LEU:HD13	1:B:175:ALA:O	2.12	0.49
1:C:47:LYS:NZ	5:C:4027:HOH:O	2.42	0.49
1:B:45[A]:LEU:N	1:B:45[A]:LEU:CD1	2.69	0.48
1:A:80:HIS:HE1	5:A:4104:HOH:O	1.97	0.48
1:B:68:ILE:HG21	1:B:86:ALA:HA	1.95	0.47
1:C:140:ALA:HA	1:C:143[A]:ILE:HG13	1.98	0.45
1:A:49:TYR:OH	1:A:212:ASP:OD2	2.32	0.45
1:A:114:PHE:CZ	1:A:146:LEU:HD11	2.52	0.44
1:C:252:ALA:CA	1:C:254:LEU:N	2.71	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:36:ALA:HB1	1:A:66[B]:CYS:HB2	2.00	0.44
1:B:32:CYS:O	1:B:276:ARG:NH1	2.49	0.43
1:B:105:ARG:HD3	1:B:108:ASP:OD2	2.16	0.43
1:B:110:ARG:NH2	5:B:4076:HOH:O	2.49	0.43
1:A:68:ILE:HG21	1:A:86:ALA:HA	1.99	0.43
1:A:32[B]:CYS:HB3	1:A:66[B]:CYS:HB3	2.01	0.43
1:A:82:LEU:HD21	1:A:128:LEU:HD13	2.00	0.43
1:B:256[A]:CYS:O	1:B:287:THR:HA	2.19	0.43
1:B:170:ASN:HB2	5:B:4135:HOH:O	2.20	0.42
1:B:298:ARG:HD3	5:B:4229:HOH:O	2.20	0.42
1:A:45[A]:LEU:CD1	1:A:54:TYR:HB3	2.49	0.42
1:C:270:TRP:HA	1:C:270:TRP:CE3	2.54	0.42
1:C:76[B]:ASP:O	1:C:78:ALA:N	2.38	0.42
1:C:257:ASP:OD1	1:C:285:MET:HE3	2.19	0.42
1:C:223:THR:HB	1:C:225:ILE:HG13	2.03	0.41
1:A:164:VAL:HG23	5:A:4168:HOH:O	2.20	0.41
1:A:177:ASP:OD1	1:A:177:ASP:N	2.52	0.41
1:C:226:SER:HB2	1:C:228:PHE:HB3	2.03	0.41
1:A:256[A]:CYS:O	1:A:287:THR:HA	2.21	0.40
1:C:76[A]:ASP:C	1:C:78:ALA:CA	2.76	0.40
1:A:40:ASP:O	1:A:66[B]:CYS:N	2.54	0.40
1:B:129:GLN:NE2	1:B:170:ASN:HB3	2.36	0.40
1:C:256[A]:CYS:O	1:C:287:THR:HA	2.21	0.40
1:A:214:ARG:NH1	1:A:285:MET:SD	2.95	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:4052:HOH:O	5:C:4140:HOH:O[1_565]	1.79	0.41
5:A:4115:HOH:O	5:B:4079:HOH:O[6_555]	1.85	0.35
5:A:4173:HOH:O	5:B:4045:HOH:O[6_545]	2.13	0.07

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	275/303 (91%)	263 (96%)	12 (4%)	0	100	100
1	B	273/303 (90%)	262 (96%)	10 (4%)	1 (0%)	30	19
1	C	273/303 (90%)	261 (96%)	12 (4%)	0	100	100
All	All	821/909 (90%)	786 (96%)	34 (4%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	281	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/229 (92%)	206 (98%)	4 (2%)	50	41
1	B	208/229 (91%)	202 (97%)	6 (3%)	37	25
1	C	207/229 (90%)	203 (98%)	4 (2%)	50	41
All	All	625/687 (91%)	611 (98%)	14 (2%)	51	34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45[A]	LEU
1	A	45[B]	LEU
1	A	107	GLU
1	A	311	THR
1	B	30	ARG
1	B	45[A]	LEU
1	B	45[B]	LEU
1	B	107	GLU
1	B	181	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	276	ARG
1	C	45	LEU
1	C	107	GLU
1	C	143[A]	ILE
1	C	143[B]	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	HIS
1	A	157	GLN
1	B	80	HIS
1	C	80	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	4
1	B	2
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	76[B]:ASP	C	78:ALA	N	1.87
1	A	57:THR	C	66[B]:CYS	N	1.73
1	C	252:ALA	C	254:LEU	N	1.72
1	C	76[A]:ASP	C	78:ALA	N	1.68
1	B	277:ALA	C	281:ALA	N	1.66
1	B	1227:ASP	C	2227:VAL	N	1.61
1	C	57:THR	C	66[A]:CYS	N	1.60

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/303 (88%)	-0.66	2 (0%) 84 85	3, 10, 22, 37	11 (4%)
1	B	267/303 (88%)	-0.57	4 (1%) 72 72	4, 10, 22, 36	12 (4%)
1	C	269/303 (88%)	-0.49	4 (1%) 72 72	4, 12, 26, 40	8 (2%)
All	All	805/909 (88%)	-0.57	10 (1%) 76 77	3, 10, 24, 40	31 (3%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	30	ARG	5.4
1	B	30	ARG	4.3
1	C	28	ALA	4.0
1	A	211	ASP	3.4
1	C	211	ASP	2.9
1	C	31	GLY	2.9
1	B	209[A]	GLU	2.5
1	B	211	ASP	2.4
1	B	281	ALA	2.3
1	A	210	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	C	3228	1/1	0.99	0.03	12,12,12,12	1
3	MG	A	3229	1/1	0.99	0.04	10,10,10,10	0
3	MG	B	3229	1/1	0.99	0.03	11,11,11,11	0
3	MG	C	3229	1/1	0.99	0.06	11,11,11,11	0
2	CA	A	3228	1/1	1.00	0.06	12,12,12,12	1
2	CA	B	3228	1/1	1.00	0.03	12,12,12,12	1
4	ZN	A	3230	1/1	1.00	0.01	9,9,9,9	0
4	ZN	A	3231	1/1	1.00	0.04	13,13,13,13	0
4	ZN	B	3230	1/1	1.00	0.01	9,9,9,9	0
4	ZN	B	3231	1/1	1.00	0.07	17,17,17,17	0
4	ZN	C	3230	1/1	1.00	0.01	8,8,8,8	0
4	ZN	C	3231	1/1	1.00	0.01	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.